

# ERRATA

## Corrections to published RIFE reports

	Page, Section	Comment															
RIFE-1 1995	38, Section 16.2	Last but one sentence, replace 1994 with 1995.															
	39, Section 16.4	1 <sup>st</sup> sentence, 2 <sup>nd</sup> paragraph, replace 1994 with 1995.															
	45, Table 1	Replace <sup>241</sup> Am Sellafield (sea pipelines) limit of 1.3 TBq with 0.3 TBq. Replace <sup>60</sup> Co Harwell (pipeline) percentage of 1.5 with 6.9.															
	74, Table 16 99, Table 33(a)	The following activity in soil data were reported as being Bq kg <sup>-1</sup> (dry) whilst they should have been reported as Bq kg <sup>-1</sup> (wet). All data are averages unless stated.															
		<table border="1"> <thead> <tr> <th>Site/location</th> <th><sup>210</sup>Po</th> <th><sup>238</sup>Pu</th> <th><sup>239+240</sup>Pu</th> </tr> </thead> <tbody> <tr> <td>Sellafield (Table 16)</td> <td>64</td> <td></td> <td></td> </tr> <tr> <td>Aldermaston (Table 33(a))</td> <td></td> <td>0.0091</td> <td>0.36</td> </tr> <tr> <td>max</td> <td></td> <td>0.016</td> <td>0.56</td> </tr> </tbody> </table>	Site/location	<sup>210</sup> Po	<sup>238</sup> Pu	<sup>239+240</sup> Pu	Sellafield (Table 16)	64			Aldermaston (Table 33(a))		0.0091	0.36	max		0.016
Site/location	<sup>210</sup> Po	<sup>238</sup> Pu	<sup>239+240</sup> Pu														
Sellafield (Table 16)	64																
Aldermaston (Table 33(a))		0.0091	0.36														
max		0.016	0.56														
	99, Table 33(a)	The concentration of <sup>137</sup> Cs in clay at Outfall (Pangbourne) was 12±0.15 Bq kg <sup>-1</sup> (dry)															
	133, Appendix 3	The average consumption rates of nuts and offal by 10 year old children were 1.5 kg y <sup>-1</sup> . The consumption of whelks at Sellafield by group E (Whitehaven commercial) was 11 kg y <sup>-1</sup> .															
	138, Appendix 6	The values of t <sub>f</sub> and t <sub>s</sub> were 0. The transfer factors for beef offal ( <sup>241</sup> Pu) and lamb ( <sup>241</sup> Pu) were 2 10 <sup>-2</sup> and 4 10 <sup>-4</sup> respectively.															
RIFE-2 1996	32, Section 8.1	lines 8-11. Replace with “In 1996 no fragments of spent fuel were found on the public beach at Dounreay. Thirteen small fragments were found with caesium-137 activities in the range 10 <sup>5</sup> -10 <sup>8</sup> Bq (these activities were measured by the operator). They were all found on the Dounreay foreshore which although a public area is largely inaccessible. A”															
	58, Table 2	Replace <sup>35</sup> S Oldbury limit of 0.8 TBq with 0.75 TBq. Replace <sup>41</sup> Ar Trawsfynydd limit of 350 TBq with 3500 TBq.															

85, Table 16  
87, Table 18  
91, Table 20(a)  
95, Table 21  
119, Table 41

The following activity in soil data were reported as being Bq kg<sup>-1</sup> (dry) whilst they should have been reported as Bq kg<sup>-1</sup> (wet). All data are averages unless stated.

Site/location	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U
Drigg (Table 16)	8.3	0.28	7.4
Ravenglass (Table 18)	16	0.56	15
Springfields (Table 20(a))	49	2.3	45
Capenhurst (Table 21)	9.8	0.36	10
Derby (Table 41)	44	1.7	43

Table 47

This was omitted in error. The data are attached.

Table 47. Radioactivity in plants near landfill sites, 1996

Sampling location	Material	No of samples	Mean radioactivity concentration (dry)*, Bq kg <sup>-1</sup>							
			<sup>3</sup> H	<sup>14</sup> C	<sup>90</sup> Sr	<sup>125</sup> I	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>238</sup> Pu	<sup>239+240</sup> Pu
Beddingham Lewes, East Sussex	Grass	4	<40 ±18	130 ±28	1.8 ±0.1	<0.19	<0.61	<0.54 ±0.30	<0.00099 ±0.00037	0.0067 ±0.0012
Cilgwyn Quarry, Gwynedd	"	4	<30	360 ±55	3.0 ±0.2	<0.63	<0.69	<5.2 ±0.9	<0.0095	0.018 ±0.005
Lyndown, Devon	"	4	<28	150 ±30	2.4 ±0.2	<1.3 ±0.2	<0.60	<0.62 ±0.17	<0.0010	<0.0024 ±0.0009
Witton, Cheshire	"	4	<38	130 ±33	0.76 ±0.12	<1.1 ±0.3	<0.59	<0.63	<0.0013	0.0021 ±0.0016

\* Results are available for other artificial nuclides detectable by gamma spectrometry  
All such results are less than the limit of detection

RIFE-3  
1997

19, Table 1.1

Replace beta, tritium and <sup>60</sup>Co Devonport (sewer) discharges with 1.97 10<sup>-6</sup>, 2.22 10<sup>-6</sup>, 5.60 10<sup>-7</sup> TBq respectively.

Replace alpha and beta limit and percentage Greenwich with 4.44 10<sup>-3</sup> TBq and <1 respectively.

21, Table 1.2

Replace tritium Winfrith limit with 5 TBq.

38, Section 3.6.5

1<sup>st</sup> paragraph. Reference to factor of 0.85 millisievert per milligray should be ICRP (1996b).

70, Table 4.10

72, Table 4.12

81, Table 4.16

121, Table 9.1

The following activity in soil data were reported as being Bq kg<sup>-1</sup> (dry) whilst they should have been reported as Bq kg<sup>-1</sup> (wet). All data are averages unless stated.

Site/location	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U
Drigg (Table 4.10)	9.9	0.37	9.5
Ravenglass (Table 4.12)	18	0.60	16
Springfields (Table 4.12)	31	1.5	30
Capenhurst (Table 4.16)	9.5	0.40	9.5
Derby (Table 9.1)	27	0.97	24

90, Section 6.3

The maximum dose due to gaseous disposals was received by adults.

161, Appendix 4

The 1 year old child dose coefficient for <sup>99</sup>Tc was 4.80 10<sup>-9</sup>.

RIFE-4  
1998

Page, Section	Comment																
70, Table 4.12	The concentrations of total Cs and $^{144}\text{Ce}$ in ovine muscle (max) were 0.61 and $<1.8 \text{ Bq kg}^{-1}$ (wet) respectively. No value for $^{155}\text{Eu}$ is available.																
75, Table 4.15(a) 77, Table 4.16 116, Table 9.1	The following activity in soil data were reported as being $\text{Bq kg}^{-1}$ (dry) whilst they should have been reported as $\text{Bq kg}^{-1}$ (wet). All data are averages unless stated.																
	<table border="1"> <thead> <tr> <th>Site/location</th> <th><math>^{234}\text{U}</math></th> <th><math>^{235}\text{U}</math></th> <th><math>^{238}\text{U}</math></th> </tr> </thead> <tbody> <tr> <td>Springfields (Table 4.15(a))</td> <td>72</td> <td>3.0</td> <td>68</td> </tr> <tr> <td>Capenhurst (Table 4.16)</td> <td>7.9</td> <td>0.30</td> <td>7.4</td> </tr> <tr> <td>Derby (Table 9.1)</td> <td>31</td> <td>0.93</td> <td>26</td> </tr> </tbody> </table>	Site/location	$^{234}\text{U}$	$^{235}\text{U}$	$^{238}\text{U}$	Springfields (Table 4.15(a))	72	3.0	68	Capenhurst (Table 4.16)	7.9	0.30	7.4	Derby (Table 9.1)	31	0.93	26
Site/location	$^{234}\text{U}$	$^{235}\text{U}$	$^{238}\text{U}$														
Springfields (Table 4.15(a))	72	3.0	68														
Capenhurst (Table 4.16)	7.9	0.30	7.4														
Derby (Table 9.1)	31	0.93	26														
96, Table 6.4(a)	The concentration of $^{241}\text{Am}$ in mud at Paddy's Hole was $<1.0 \text{ Bq kg}^{-1}$ (dry). No measurement of $^{239/240}\text{Pu}$ was made.																
125, Section 11.1	Last but one paragraph. The estimated dose was 0.094 mSv.																
131, Section 11.8	Last paragraph, first sentence. Replace 1997 with 1998.																

RIFE-5  
1999

71, Table 4.15(a) 73, Table 4.16 118, Table 9.1	The following activity in soil data were reported as being $\text{Bq kg}^{-1}$ (dry) whilst they should have been reported as $\text{Bq kg}^{-1}$ (wet). All data are averages unless stated.																
	<table border="1"> <thead> <tr> <th>Site/location</th> <th><math>^{234}\text{U}</math></th> <th><math>^{235}\text{U}</math></th> <th><math>^{238}\text{U}</math></th> </tr> </thead> <tbody> <tr> <td>Springfields (Table 4.15(a)) max</td> <td>180</td> <td>15</td> <td>200</td> </tr> <tr> <td>Capenhurst (Table 4.16) max</td> <td>12</td> <td>0.46</td> <td>12</td> </tr> <tr> <td>Derby (Table 9.1) max</td> <td>34</td> <td>1.3</td> <td>31</td> </tr> </tbody> </table>	Site/location	$^{234}\text{U}$	$^{235}\text{U}$	$^{238}\text{U}$	Springfields (Table 4.15(a)) max	180	15	200	Capenhurst (Table 4.16) max	12	0.46	12	Derby (Table 9.1) max	34	1.3	31
Site/location	$^{234}\text{U}$	$^{235}\text{U}$	$^{238}\text{U}$														
Springfields (Table 4.15(a)) max	180	15	200														
Capenhurst (Table 4.16) max	12	0.46	12														
Derby (Table 9.1) max	34	1.3	31														
112, Section 8.2	The second sentence of paragraph three states that "the duck and tide washed pasture pathways gave doses of 0.032 and 0.009 mSv $\text{y}^{-1}$ respectively." The dose due to the duck pathway should read 0.042 mSv $\text{y}^{-1}$ . The value for tide washed pasture is correct.																
123, Table 10.2	The concentration of $^{14}\text{C}$ in grass from Billingham was 960 $\text{Bq kg}^{-1}$ (wet).																
162, Table A1.2	The Dounreay (Fast Reactor) data were duplicated.																

RIFE-6  
2000

31, Section 3.5	It was stated that the dose limits do not apply to natural radionuclides. This sentence should be deleted.												
75, Table 4.16 124, Table 9.1	The following activity in soil data were reported as being $\text{Bq kg}^{-1}$ (dry) whilst they should have been reported as $\text{Bq kg}^{-1}$ (wet). All data are averages unless stated.												
	<table border="1"> <thead> <tr> <th>Site/location</th> <th><math>^{234}\text{U}</math></th> <th><math>^{235}\text{U}</math></th> <th><math>^{238}\text{U}</math></th> </tr> </thead> <tbody> <tr> <td>Capenhurst (Table 4.16) max</td> <td>8.5</td> <td>0.35</td> <td>8.4</td> </tr> <tr> <td>Derby (Table 9.1) max</td> <td>24</td> <td>0.96</td> <td>23</td> </tr> </tbody> </table>	Site/location	$^{234}\text{U}$	$^{235}\text{U}$	$^{238}\text{U}$	Capenhurst (Table 4.16) max	8.5	0.35	8.4	Derby (Table 9.1) max	24	0.96	23
Site/location	$^{234}\text{U}$	$^{235}\text{U}$	$^{238}\text{U}$										
Capenhurst (Table 4.16) max	8.5	0.35	8.4										
Derby (Table 9.1) max	24	0.96	23										

Page, Section	Comment
155, Table 12.1	Target date for project 'Tritium and carbon-14 in seafood' should have been March 2003.
166, Table A1.1	Discharges of tritium from Devonport (pipeline) given as 0.87 TBq should have been 0.087 TBq.
168, Table A1.2	Sellafield Discharge limits of alpha and beta activity should have been 0.00196 and 0.328 TBq. Percentage of limit for alpha and beta activity should have been 4.0 and <1. Discharges of tritium and <sup>14</sup> C from Sellafield given as 213 and 2.58 TBq should have been 355 and 2.94 TBq. Relevant percentages given as 15 and 30 should have been 25 and 34.
RIFE-7 2001 71, Table 4.8 80, Table 4.15(a) 93, Table 5.2(a) 122, Table 7.3 127, Table 8.2(a) 130, Table 9.1	The following activity in soil data were reported as being Bq kg <sup>-1</sup> (dry) whilst they should have been reported as Bq kg <sup>-1</sup> (wet). All data are averages unless stated.

Site/location	<sup>60</sup> Co	<sup>106</sup> Ru	<sup>125</sup> Sb	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U	<sup>241</sup> Am
Sellafield (Table 4.8)	<0.80	<3.1	<1.1		80				5.8
max	1.2				97	9.3	0.34	9.1	6.0
Springfields (Table 4.15(a))						95	4.6	89	
max									
Harwell (Table 5.2(a))	<0.40			<0.40	2.9				
Featherstone position A (Table 7.3)						9.5	0.41	9.0	
Featherstone position B (Table 7.3)						7.3	0.34	7.5	
Cardiff (Table 8.2(a))				<0.33	5.6				
max				<0.40	6.5				
Derby (Table 9.1)						18	0.80	18	
max						30	1.3	29	

176, Table A1.1 Discharges of Alpha for Hunterston 'A' given as 0.14 TBq should have been 1.4 10<sup>-5</sup>TBq. The % of limit given as 350 should have been <1.

181, Table A1.2 Dungeness 'A' discharge limit and % of limit for tritium should have been 3 and 23 respectively.

RIFE-8  
2002  
59, Table 4.1 Two tritium results were omitted.  
The data are attached.

Table 4.1. Beta/gamma radioactivity in fish from the Irish Sea vicinity and further afield, 2002

Location	Material	No. of sampling observations	<sup>3</sup> H
Liverpool Bay	Flounder	2	<25
Mersey estuary	Flounder	2	<25

79, Table 4.14  
82 Table 4.17  
128, Table 7.1(a)  
138, Table 8.2(a)

The following activity in soil data were reported as being  $\text{Bq kg}^{-1}$  (dry) whilst they should have been reported as  $\text{Bq kg}^{-1}$  (wet). All data are averages unless stated.

Site/location	$^{60}\text{Co}$	$^{106}\text{Ru}$	$^{125}\text{Sb}$	$^{134}\text{Cs}$	$^{137}\text{Cs}$	$^{234}\text{U}$	$^{235}\text{U}$	$^{238}\text{U}$
Sellafield (Table 4.14)	<0.80	<2.3	<1.2	68				
max	1.0	<2.7	<1.4	82				
Drigg (Table 4.17)								
max						6.9	0.30	6.5
Aldermaston (Table 7.1(a))								
max						8.7	0.35	8.3
Cardiff (Table 8.2(a))				<0.30	6.4			
max					8.1			

102, Figure 6.1

An incorrect bar in Figure 6.1 for Bradwell (2002) is corrected below




Figure 6.1. Caesium-137 concentration in sediments near nuclear power stations

RIFE-1 - RIFE-8  
1995-2002

Urenco Capenhurst have reassessed atmospheric discharges of uranium; the reassessed discharges are listed in Table E1.

Table E1. Reassessed atmospheric discharges of uranium from Urenco Capenhurst

Year	Original reported discharge TBq	Reassessed discharge TBq
1993	$1.74 \times 10^{-9}$	$2.41 \times 10^{-7}$
1994	$6.74 \times 10^{-9}$	$2.63 \times 10^{-7}$
1995	$2.69 \times 10^{-8}$	$2.75 \times 10^{-7}$
1996	$1.11 \times 10^{-7}$	$8.23 \times 10^{-7}$
1997	$6.80 \times 10^{-8}$	$4.90 \times 10^{-7}$
1998	$6.87 \times 10^{-8}$	$1.87 \times 10^{-6}$
1999	$8.15 \times 10^{-8}$	$1.01 \times 10^{-6}$
2000	$9.64 \times 10^{-8}$	$8.72 \times 10^{-7}$
2001	$1.20 \times 10^{-7}$	$9.77 \times 10^{-7}$
2002	$1.16 \times 10^{-7}$	$6.01 \times 10^{-7}$

RIFE-9  
2003

82, Table 3.15  
138 Table 6.1(a)  
141, Table 6.3(a)  
151, Table 7.3(a)  
157, Table 8.1(a)

The following activity in soil data were reported as being Bq kg<sup>-1</sup> (dry) whilst they should have been reported as Bq kg<sup>-1</sup> (wet). All data are averages unless stated.

Site/location	<sup>60</sup> Co	<sup>106</sup> Ru	<sup>125</sup> Sb	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>154</sup> Eu	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U	<sup>241</sup> Am
Sellafield (Table 3.15)	<0.90	<3.3	<1.2	<0.40	75	<0.50				5.9
max	1.6	<4.2	<1.6		89	<0.60	11	0.54	10	7.7
Aldermaston (Table 6.1(a))							11	0.48	11	
max										
Derby (Table 6.3(a))							47	1.6	40	
max										
Cardiff (Table 7.3(a))				<0.40	8.8					
max					11					
Drigg (Table 8.1)							6.7	0.26	6.7	
max										

185, Table 9.12

Some data were incorrect. The amended version of the table is attached.

Table 9.12. Concentrations of radionuclides in rainwater and air 2003

Location	Sample	No. of sampling observations	Mean radioactivity concentration <sup>a</sup> in rainwater and air									
			<sup>3</sup> H <sup>7</sup>	Be	<sup>90</sup> Sr <sup>b</sup>	<sup>137</sup> Cs	<sup>210</sup> Pb	<sup>210</sup> Po	<sup>228</sup> Th	Gross alpha <sup>b</sup>	Gross beta <sup>b</sup>	
Ceredigion Aberporth	Rainwater	12	<2.4	<1.6		<0.053	0.10			*		
	Air	4		0.0022		<0.00000052	0.00017			*		
Co. Down Conlig	Rainwater	4		<1.5		<0.022	*			*		
	Air	4		0.0022		<0.00000063	0.00015			*		
Dumfries and Galloway Eskdalemuir	Rainwater	4	4	<2.7	1.2		<0.0098	0.094		*		
	Air	4		0.0018		<0.00000043	0.00013			*		
North Yorkshire Dishforth	Rainwater	4		<2.2		<0.039	*			*		
	Air	4		0.0016		<0.00000055	0.00014			*		
Oxfordshire Chilton	Rainwater	12		<1.5	<0.00064	<0.032	0.32			*	0.074	0.17
	Air	13		0.0018		<0.00000034	0.00027	<0.000014		*		
Shetland Lerwick	Rainwater	4		1.6		<0.017	*			*		
	Air	4		0.0015		<0.00000052	0.00010			*		
Suffolk Orfordness	Rainwater	4	<2.2	<2.4		<0.048	*			5.2		
	Air	4		0.0022		<0.00000053	0.00020			*		

\* Not detected by the method used

<sup>a</sup> Bq l<sup>-1</sup> for rainwater and Bq kg<sup>-1</sup> for air

<sup>b</sup> Annual bulk analysis

187, Table 9.14

The concentration of <sup>210</sup>Po in Cornwall, River Fowey was <0.0098 Bq l<sup>-1</sup>.

Table 9.16. Estimates of maximum radiation exposure from radionuclides in drinking water, 2003<sup>a</sup>

Country	Exposure, mSv		
	Man-made radionuclides <sup>b</sup>	Natural radionuclides <sup>c</sup>	All radionuclides
England	<0.001	0.028	0.028
Northern Ireland	<0.001	0.026	0.026
Scotland	<0.001	<sup>d</sup>	<sup>d</sup>
Wales	<0.001	0.027	0.027

<sup>a</sup> The maximum dose is selected for each nuclide group from data for individual sampling locations.  
 Many estimates of dose are based on concentration results at limits of detection.

<sup>b</sup> Including tritium

<sup>c</sup> Including carbon-14

<sup>d</sup> Analysis of natural radionuclides was not undertaken

214, Table A1.2

The data shown for Faslane are a duplication of the data for Rosyth and were included in error.

RIFE-10  
2004

75, Table 3.7

The entry for Haverigg should read 0.087.

45, Figure 3.8

An incorrect bar in Figure 3.8 for Americium discharge is corrected below:




Figure 3.8. Americium-241 and liquid discharge from Sellafield and concentrations in cod\*, lobsters and winkles near Sellafield (\* estimated in 2004 due to lack of availability of cod)

87, Table 3.15

151 Table 6.1(a)

154, Table 6.3(a)

166, Table 7.3(a)

173, Table 8.1(a)

The following activity in soil data were reported as being Bq kg<sup>-1</sup> (dry) whilst they should have been reported as Bq kg<sup>-1</sup> (wet). All data are averages unless stated.

Site/location	<sup>60</sup> Co	<sup>106</sup> Ru	<sup>125</sup> Sb	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U
Sellafield (Table 3.15)	<0.43	<1.4	<0.73					
max	0.80	<1.5	<0.80			16	0.64	15
Aldermaston (Table 6.1(a))								
max						7.8	0.29	7.2
Derby (Table 6.3(a))								
max						27	0.94	23
Cardiff (Table 7.3(a))				<0.47	7.1			
max				<0.50	7.7			
Drigg (Table 8.1)								
max						11	0.42	11

Page, Section	Comment
223, Table A1.1	The % annual limit for <sup>106</sup> Ru discharge at Sellafield was 7% (not 70%).
246, Table A5.1	Some dose per unit intake values were missing for 1 yr old. These were:

Table A5.1. Dosimetric data

Radionuclide	Dose per unit intake by inhalation using ICRP-60 methodology (Sv Bq <sup>-1</sup> )
Sr-90 <sup>†</sup>	1.2E-07
Zr-95 <sup>†</sup>	2.1E-08
Ba-140 <sup>†</sup>	2.6E-08
Pb-210 <sup>†</sup>	4.0E-06
Th-228 <sup>†</sup>	1.4E-04
U-238	9.4E-06

<sup>†</sup> Energy and dose per unit intake data include the effects of radiations of short-lived daughter products

RIFE-11  
2005

72, Table 3.3a	Footnote 'd' showed an incorrect value. It should have read: <sup>d</sup> The concentration of <sup>237</sup> Np was 0.00035 Bq kg <sup>-1</sup>
112, Table 4.3a	Column headings should have read: <sup>239</sup> Pu+ <sup>240</sup> Pu <sup>241</sup> Pu
140, Table 5.5a	The result of <0.13 for <sup>241</sup> Am in the <i>Fucus vesiculosus</i> samples from Pilot Station was incorrectly put into the <sup>239</sup> Pu+ <sup>240</sup> Pu column.
206, Figures 9.5 and 9.6	Incorrect units were shown. The correct units were mBq l <sup>-1</sup> .
225, Table 9.15	Incorrect headings in the top part of the table. Should have been as below:

Table 9.15. Concentrations of radionuclides in sources of drinking water in England and Wales, 2005

Location	Sample source	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>					
			<sup>3</sup> H	<sup>40</sup> K	<sup>90</sup> Sr	<sup>137</sup> Cs	<sup>210</sup> Po	
Wales								
Gwynedd	Cwm Ystradlyn Treatment Works	4	<4.0	<0.020	0.0036	0.0018	<0.010	
Mid-Glamorgan	Llwyn-on Reservoir	4	<4.0	<0.045	0.0030	<0.0010	<0.013	
Powys	Elan Valley Reservoir	4	<4.0	<0.050	0.0040	0.00090	<0.010	

248, Table A1.2	Sellafield discharge limits for alpha and beta should have been 8.90 10 <sup>-5</sup> and 0.00174 TBq respectively.
251, Table A1.2	Aldermaston Tritium discharge and % limit should have been 14.1 and 8.3 respectively.

RIFE 8-11  
2002-2005

Concentrations in sediments	For sediment samples with unusually high water contents it was discovered in 2007 that the resulting sample bulk densities were outside the instrument calibration range. Following investigations a correction factor has been calculated and this has been applied to the affected data from 2002-2005 and the new results are reported here in Table E2.
-----------------------------	---



These amendments do not significantly affect any assessments, charts or statements in the relevant RIFE reports.

Table E2. Amended concentrations of radionuclides in sediment, 2002-2005

Year	Site	Location	No. of sampling observations	Mean radioactivity concentration (dry), Bq kg <sup>-1</sup>						
				<sup>57</sup> Co	<sup>60</sup> Co	<sup>65</sup> Zn	<sup>95</sup> Zr	<sup>95</sup> Nb	<sup>106</sup> Ru	<sup>125</sup> Sb
2002	Aldermaston	Reading (Kennet)	4							
		Stream draining south	4							
	Bradwell	Maldon	2		<3.4					
		Waterside	2		<4.0					
	Capenhurst	Rossmore (4.3 km downstream)	2							
	Cardiff	Canal	2							
		West of pipeline	2							
	Devonport	Lopwell	2		<3.7					
	Dungeness	Pilot Sands	2		<0.90					
	Harwell	Appleford	4		<0.60					
Day's Lock		4		<0.50						
Sellafield	Caerhun	2		<3.3		<9.6	<7.7	<23	<9.2	
2003	Aldermaston	Reading (Kennet)	4							
		Aldermaston	4							
	Amersham	Outfall (Grand Union Canal)	3	<0.30	<1.1	<1.5				
	Bradwell	Waterside	2		<2.0					
		Canal	1							
	Derby	River Derwent (downstream)	4		<1.0					
Devonport	Lopwell	2		<2.5						
2004	Aldermaston	Reading (Kennet)	4							
		Aldermaston	4							
		Stream draining south	4							
	Amersham	Upstream of outfall (Grand Union Canal)	2	<6.4	<1.8	<4.1				
	Sellafield	Caerhun	2		<1.6		<4.5	<2.2	<12	<13
2005	Aldermaston	Reading (Kennet)	4							
	Amersham	Upstream of outfall (Grand Union Canal)	2	<5.3	<1.6	<3.6				
	Cardiff	Canal	2							
		Lydebank Brook	4		<1.7					
	Harwell	Appleford	4		<2.5					
		Caerhun	2		<2.6		<8.8	<6.8	<20	<20
Trawsfynydd	Bailey Bridge	2		<8.3					<44	

Year	Site	Location	No. of sampling observations	Mean radioactivity concentration (dry), Bq kg <sup>-1</sup>							
				<sup>125</sup> I	<sup>131</sup> I	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>144</sup> Ce	<sup>154</sup> Eu	<sup>155</sup> Eu	<sup>241</sup> Am
2002	Aldermaston	Reading (Kennet)	4				7.3				<1.9
		Stream draining south	4				<5.1				<1.2
	Bradwell	Maldon	2			6.5	80				<4.0
		Waterside	2			3.9	59				<13
	Capenhurst	Rossmore (4.3 km downstream)	2				<4.4				
	Cardiff	Canal	2	<0.80			2.4				
		West of pipeline	2	<3.1			33				
	Devonport	Lopwell	2				7.7				
	Dungeness	Pilot Sands	2				<0.90				<1.6
	Harwell	Appleford	4				<13				
Day's Lock		4				6.0					
Sellafield	Caerhun	2			<3.4	430	<25	<7.3	<8.0	75	
2003	Aldermaston	Reading (Kennet)	4				8.0				<1.6
		Aldermaston	4				6.3				<2.7
	Amersham	Outfall (Grand Union Canal)	3	<1.0	<550		<2.1				
	Bradwell	Waterside	2				35				<2.7
		Canal	1	<1.4			16				
	Devonport	Lopwell	2				<10				
2004	Aldermaston	Reading (Kennet)	4				5.4				<1.1
		Aldermaston	4				<3.9				<1.3
		Stream draining south	4				<2.8				1.6
	Amersham	Upstream of outfall (Grand Union Canal)	2	<0.80	<1.4		10				
	Sellafield	Caerhun	2	<1.5		<1.5	220	<5.7	<7.3	<3.1	51
2005	Aldermaston	Reading (Kennet)	4				<3.9				6.5
	Amersham	Upstream of outfall (Grand Union Canal)	2	<1.0	<9.1		6.2				
	Cardiff	Canal	2	<1.8			9.1				
		Lydebank Brook	4				9.0				
	Harwell	Appleford	4				<11				
		Caerhun	2			<2.5	230	<9.3	<12	<5.3	59
Trawsfynydd	Bailey Bridge	2			<4.2	920				76	

	Page, Section	Comment
RIFE-11 2005	270, Table A7.2B	Trawsfynydd, should read... Prenatal children of occupants over sediment 0.008 Direct radiation, gamma dose rate over sand/stone
	70, Table 2.17	The concentration of <sup>241</sup> Am in winkles at Drigg should have been 29.
RIFE-12 2006	103, Section 4 Key points	Line 22 second column replace with • At Dungeness, dose from gaseous discharges increased.
	187, Figure 8.5	The range in the key should have been 2 to 8.
	234, Table A4.2B	Trawsfynydd, should read... Prenatal children of fish consumers 0.013 Fish, gamma dose rate over sediment, <sup>90</sup> Sr
	127, Table 4.5a	The <sup>210</sup> Po and <sup>210</sup> Pb results are the wrong way round for South Gare winkles. <sup>210</sup> Po should be 11 and <sup>210</sup> Pb should be 0.46 Bq kg <sup>-1</sup>
RIFE-13 2007	153, Table 5.1	Derby, the total exposure and exposure from intakes of sediment and water should have been <0.005 mSv.
	161, Section 6 Key points	Line 17 second column should read... • The total dose of 0.008...
	236, Table A4.2B	Trawsfynydd, should read... Adult fish consumers 0.014 Fish, gamma dose rate over sediment, <sup>90</sup> Sr, <sup>137</sup> Cs, <sup>241</sup> Am
	239, Appendix 5	Line 3 first column should read... ... indicated that it was likely there would be no adverse impact
	12, Figure S1	Both bars for Bradwell should be the same height. The bar for exposures due to liquid wastes is wrong.
RIFE-14 2008	33, Section 2	Springfields, doses to the public Lines 1 & 2 second column should read... ...pathways from gaseous discharges were less than 0.005mSv which was less than 0.5 per cent...
	51, Figure 2.22	The bar for Whitehaven in 2008 should have been the same height as the bar for 2007
	109, Section 4	Gaseous discharges and terrestrial monitoring Line 28, first column should read... The results of monitoring for 2008...

Page, Section                      Comment

RIFE-14  
2008

167, Table 6.3a                      Results for Cardiff East WWTW should have been:

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>			
			Organic			
			<sup>3</sup> H <sup>e</sup>	<sup>3</sup> H	<sup>3</sup> H <sup>f</sup>	14 <sup>c</sup>
<b>Terrestrial samples</b>						
Crude effluent	Cardiff East WWTW	3E	<150	<220	82	<11
Final effluent	Cardiff East WWTW	3E	<60	<70	80	<11
Sludge pellets	Cardiff East WWTW	3E		76000		740
Solids from crude effluent	Cardiff East WWTW	3E		<7500		<1800

225, Table A2.2                      Sellafield (sea pipelines) Tritium discharge limit should have read 2 10<sup>4</sup>

236, Table A4.2B                      Trawsfynydd, should read...  
Adult fish consumers              0.010      Fish, gamma dose rate over sediment, <sup>90</sup>Sr, <sup>137</sup>Cs, <sup>241</sup>Am

RIFE-15  
2009

233, Table A2.1                      MoD Coulport under reported discharges for the end of 2009. The <sup>3</sup>H discharge for 2009 should have been 3.40 E-03 TBq.

249, Table A4.2B                      Trawsfynydd, should read...  
Adult fish consumers              0.012      Fish, gamma dose rate over sediment, <sup>90</sup>Sr, <sup>137</sup>Cs, <sup>241</sup>Am

RIFE-16  
2010

30, Table 1.2B                      Trawsfynydd, should read...  
Adult fish consumers              0.012      Fish, gamma dose rate over sediment, <sup>90</sup>Sr, <sup>137</sup>Cs, <sup>241</sup>Am

37, Section 2                      Line 13, paragraph 3, second column should read...  
The dose to wildfowlers and farmers from exposure over salt marsh was 0.032 mSv, which was less than 4 per cent of the dose limit for members of the public of 1 mSv. The small decrease in dose from 0.036 mSv (in 2009) was due to lower gamma dose rates over marsh in 2010.

100, Section 3                      The graph in Figure 3.2 is missing 2010 data. The data for 2010 is shown in Figure 3.2 RIFE 17

122, Section 4                      Line 7, paragraph 1, first column should read...  
An increase in the fish and crustacean consumption rates has been observed, together with a decrease in the mollusc and occupancy rates, in comparison with those of the previous survey reported in 2006.

	Page, Section	Comment
RIFE-16 2010	Appendix 1, Annex 2	Table X2.2 Sellafield Group N winkle consumption should have said 15kg y <sup>-1</sup> (not 18 kg y <sup>-1</sup> )
RIFE-17 2011	52, Section 2	On Figure 2.14 the year labels from 2004 to 2011 were underneath the bar chart incorrectly and should have been one place to the right, as shown in RIFE 18.
	61, Section 2	Springfields 'Source specific doses' last entry on the table should read: 'Consumers of locally grown food' not 'Infant consumers of locally grown food'
	209, Section 9	Line 7, paragraph 7, should read: Tritium concentrations in the western English Channel were also very low (Figure 9.7).
	240, Appendix 2	Third entry on the table – Capenhurst (Urenco UK) the discharge limits (annual equivalent) <sup>a</sup> Bq column should have read: Uranium           7.50E+06 Other Alpha       2.40E+06 Technetium-99    1.00E+08 Others             2.25E+09
RIFE-14-17 2011	CD, Appendix 1	Table X2.2 Sellafield Q – Ravenglass nature warden assessment, the ingestion and inhalation rates of sediment have been incorrect, they should have read:  RIFE-14 3.1 10 <sup>-3</sup> kg y <sup>-1</sup> mud by inadvertant ingestion 5.6 10 <sup>-5</sup> kg y <sup>-1</sup> mud by resuspension and inhalation  RIFE-15 3.4 10 <sup>-3</sup> kg y <sup>-1</sup> mud by inadvertant ingestion 6.3 10 <sup>-5</sup> kg y <sup>-1</sup> mud by resuspension and inhalation  RIFE-16 3.4 10 <sup>-3</sup> kg y <sup>-1</sup> mud by inadvertant ingestion 6.3 10 <sup>-5</sup> kg y <sup>-1</sup> mud by resuspension and inhalation  RIFE-17 3.4 10 <sup>-3</sup> kg y <sup>-1</sup> mud by inadvertant ingestion 6.3 10 <sup>-5</sup> kg y <sup>-1</sup> mud by resuspension and inhalation
RIFE-18 2012	134, Table 4.1	Hinkley Point. These are small changes to the total dose and source specific dose shown below. The apply to relevant points of text, tables (S, 1.2, 1.3, 1.4 and 4.1) and figures (1.1, 4.1 and 6.2).

Site	Exposed population <sup>a</sup>	Exposure, mSv per year					
		Total	Fish and shellfish	Other local food	External radiation from intertidal areas or the shoreline	Gaseous plume related pathways	Direct radiation from site
<b>Total dose – all sources</b>	<b>Adult occupants over sediment</b>	<b>0.013</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>0.012</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>
Source specific doses	Seafood consumers	0.018	<0.005	–	0.017	–	–

Page, Section	Comment									
RIFE-18 2012	240, Appendix 2 Third entry on the table – Sellafield – the discharges during 2012 (Bq and % of annual limit) columns and should have read: <table border="1" style="margin-left: 40px;"> <tr> <td>Beta</td> <td>1.03E+09</td> <td>2.5</td> </tr> <tr> <td>Antimony-125</td> <td>3.20E+09</td> <td>11</td> </tr> <tr> <td>Caesium-137</td> <td>1.59E+08</td> <td>2.7</td> </tr> </table>	Beta	1.03E+09	2.5	Antimony-125	3.20E+09	11	Caesium-137	1.59E+08	2.7
Beta	1.03E+09	2.5								
Antimony-125	3.20E+09	11								
Caesium-137	1.59E+08	2.7								
41, Figure 2.3	The River Ribble houseboat dose rate datum for 2012 (figure 2.3, RIFE-18) was plotted incorrectly, it is shown corrected in Figure 2.4 in RIFE-19									
134, Table 2.18	Sellafield. These are small changes to the total dose shown below. They apply to relevant points of text, tables (1.2 and 2.18) and figure 2.6.									

Exposed population <sup>a</sup>	Exposure, mSv per year							
	Total	Seafood (nuclear industry discharges)	Seafood (other discharges)	Other local food	External radiation from intertidal areas, river banks or fishing gear	Intakes of sediment and water	Gaseous plume related pathways	Direct radiation from site
<b>Total dose – maximum effect of gaseous release and direct radiation sources</b>								
<b>Infant root vegetable consumers</b>	<b>0.011</b>	–	–	<b>0.011</b>	–	–	–	–

196, Table 7.7 Discharge data reported previous to RIFE-18 classified as Oil & Gas (Offshore) should have been classified as Oil & Gas (Onshore). This has been corrected for RIFE-18 onwards.