

# Radioactivity in Food and the Environment, 2015





ENVIRONMENT AGENCY  
FOOD STANDARDS AGENCY  
FOOD STANDARDS SCOTLAND  
NATURAL RESOURCES WALES  
NORTHERN IRELAND ENVIRONMENT AGENCY  
SCOTTISH ENVIRONMENT PROTECTION AGENCY

# **Radioactivity in Food and the Environment, 2015**

**RIFE – 21**

October 2016



This report was compiled by the Centre for Environment, Fisheries and Aquaculture Science on behalf of the Environment Agency, Food Standards Agency, Food Standards Scotland, Natural Resource Wales, Northern Ireland Environment Agency and the Scottish Environment Protection Agency.



Printed on paper made from a minimum 75% de-inked post-consumer waste.

Front cover photograph: Sizewell Power Stations. Published by kind permission of Cefas.

Inside cover photograph: Reproduced with kind permission of Public Health England

© Crown Copyright, 2016

Requests for the reproduction of materials contained in this report and for other information should be addressed to:

- in England and Wales,  
Radiological Monitoring and Assessment Team of the Environment Agency ([enquiries@environment-agency.gov.uk](mailto:enquiries@environment-agency.gov.uk)),  
Chemical Safety Division of the Food Standards Agency ([radiation@foodstandards.gsi.gov.uk](mailto:radiation@foodstandards.gsi.gov.uk))  
or Natural Resources Wales ([enquiries@naturalresourceswales.gov.uk](mailto:enquiries@naturalresourceswales.gov.uk))
- in Scotland,  
the Radioactive Substances Unit of SEPA ([radiologicalmonitoring@sepa.org.uk](mailto:radiologicalmonitoring@sepa.org.uk)) or the Food Standards Scotland ([will.munro@fss.scot](mailto:will.munro@fss.scot)) and
- in Northern Ireland,  
the Industrial Pollution and Radiochemical Inspectorate of NIEA ([IPRI@daera-ni.gov.uk](mailto:IPRI@daera-ni.gov.uk))

CONTENTS.....	3
LIST OF TABLES.....	5
LIST OF FIGURES .....	7
PREFACE .....	9
TECHNICAL SUMMARY.....	10
<b>1. Introduction .....</b>	<b>17</b>
1.1 Purpose and scope of the monitoring programmes.....	17
1.2 Summary of doses.....	18
1.2.1 The assessment process.....	18
1.2.2 Total dose results for 2015.....	19
1.2.3 Total dose trends.....	20
1.2.4 Source specific dose results for 2015.....	20
1.2.5 Protecting the environment.....	20
1.2.6 Food irradiation.....	23
1.3 Sources of radiation exposure.....	23
1.3.1 Radioactive waste disposal from nuclear licensed sites.....	23
1.3.2 International agreements, the UK Discharge Strategy and new nuclear power stations.....	25
1.3.3 Managing radioactive liabilities in the UK.....	26
1.3.4 Solid radioactive waste disposal at sea.....	29
1.3.5 Other sources of radioactivity.....	29
<b>2. Nuclear fuel production and reprocessing .....</b>	<b>37</b>
2.1 Capenhurst, Cheshire.....	38
2.2 Springfields, Lancashire.....	40
2.3 Sellafield, Cumbria.....	43
2.3.1 Doses to the public.....	44
2.3.2 Gaseous discharges.....	49
2.3.3 Liquid discharges.....	50
2.3.4 Monitoring of unusual pathways.....	64
2.4 Windscale, Cumbria.....	65
<b>3. Research establishments.....</b>	<b>93</b>
3.1 Dounreay, Highland.....	93
3.2 Harwell, Oxfordshire.....	97
3.3 Winfrith, Dorset.....	98
3.4 Minor sites.....	100
3.4.1 Culham, Oxfordshire.....	100
3.4.2 Imperial College Reactor Centre, Ascot, Berkshire.....	101
<b>4. Nuclear power stations.....</b>	<b>109</b>
4.1 Berkeley, Gloucestershire and Oldbury, South Gloucestershire.....	110
4.2 Bradwell, Essex.....	111
4.3 Dungeness, Kent.....	115
4.4 Hartlepool, County Durham.....	116
4.5 Heysham, Lancashire.....	117
4.6 Hinkley Point, Somerset.....	118
4.7 Sizewell, Suffolk.....	119
4.8 Chapelcross, Dumfries and Galloway.....	120
4.9 Hunterston, North Ayrshire.....	121
4.10 Torness, East Lothian.....	123
4.11 Trawsfynydd, Gwynedd.....	124
4.12 Wylfa, Isle of Anglesey.....	125

<b>5. Defence establishments .....</b>	<b>155</b>
5.1 Aldermaston, Berkshire .....	155
5.2 Barrow, Cumbria.....	156
5.3 Derby, Derbyshire .....	156
5.4 Devonport, Devon .....	157
5.5 Faslane and Coulport, Argyll and Bute.....	158
5.6 Holy Loch, Argyll and Bute .....	159
5.7 Rosyth, Fife .....	159
5.8 Vulcan NRTE, Highland.....	160
<b>6. Radiochemical production .....</b>	<b>167</b>
6.1 Grove Centre, Amersham, Buckinghamshire.....	167
6.2 Maynard Centre, Cardiff .....	168
<b>7. Industrial and landfill sites .....</b>	<b>177</b>
7.1 Low Level Waste Repository near Drigg, Cumbria.....	177
7.2 Other landfill sites .....	178
7.3 Metals Recycling Facility, Lillyhall, Cumbria .....	179
7.4 Phosphate processing, Whitehaven, Cumbria .....	180
7.5 Aberdeen .....	181
7.6 Dalgety Bay, Fife.....	182
7.7 Former RAF Kinloss, Moray.....	183
7.8 Other non-nuclear sites .....	183
<b>8. Regional monitoring .....</b>	<b>191</b>
8.1 Channel Islands.....	191
8.2 Isle of Man.....	191
8.3 Northern Ireland.....	192
8.4 Overseas sources .....	192
8.5 General diet .....	193
8.6 Milk .....	193
8.7 Crops.....	195
8.8 Airborne particulate, rain, freshwater and groundwater .....	195
8.9 Seawater surveys.....	196
<b>9. References.....</b>	<b>215</b>
<b>APPENDIX 1. Sampling, measurement, presentation and assessment methods and data .....</b>	<b>227</b>
<b>APPENDIX 2. Disposals of radioactive waste.....</b>	<b>228</b>
<b>APPENDIX 3. Abbreviations and glossary.....</b>	<b>237</b>
<b>APPENDIX 4. Research in support of the monitoring programmes .....</b>	<b>240</b>
<b>APPENDIX 5. Disposal of dredge material from Maryport Harbour, Cumbria .....</b>	<b>241</b>

# List of Tables

Abbreviated Title	Number .....	Page
<b>Technical summary</b>		
UK total dose from all sources	S.....	12
<b>Introduction</b>		
Direct radiation from nuclear licensed sites	1.1.....	31
UK total dose from all sources - details	1.2.....	32
Trends in total dose	1.3.....	34
UK source specific doses	1.4.....	35
<b>Nuclear fuel production and reprocessing</b>		
Radiation exposure - Capenhurst and Springfields	2.1.....	66
Capenhurst	2.2.....	67
Springfields	2.3.....	69
Terrestrial foodstuffs near Sellafield	2.4.....	72
Sellafield - fish beta/gamma	2.5.....	73
Sellafield - shellfish beta/gamma	2.6.....	75
Sellafield - seafood transuranics	2.7.....	77
Sellafield - marine sediment	2.8.....	79
Sellafield - gamma radiation dose rates	2.9.....	82
Sellafield - beta radiation dose rates on fishing gear	2.10.....	86
Sellafield - beta radiation dose rates on sediment	2.11.....	86
Sellafield - aquatic plants	2.12.....	87
Terrestrial foodstuffs near Ravenglass	2.13.....	89
Sellafield - surface water	2.14.....	89
Sellafield - road drains	2.15.....	90
Radiation exposure - Sellafield, Irish Sea groups	2.16.....	90
Radiation exposure - Sellafield	2.17.....	91
<b>Research establishments</b>		
Radiation exposure - research	3.1.....	102
Dounreay	3.2.....	103
Harwell	3.3.....	106
Winfrith	3.4.....	107
Culham	3.5.....	108
<b>Nuclear power stations</b>		
Radiation exposure - power stations	4.1.....	127
Berkeley and Oldbury	4.2.....	129
Bradwell	4.3.....	130
Dungeness	4.4.....	134
Hartlepool	4.5.....	136
Heysham	4.6.....	138
Hinkley Point	4.7.....	140
Sizewell	4.8.....	142
Chapelcross	4.9.....	143
Hunterston	4.10.....	146
Torness	4.11.....	149
Trawsfynydd	4.12.....	152
Wylfa	4.13.....	153
<b>Defence establishments</b>		
Radiation exposure - defence	5.1.....	161
Aldermaston	5.2.....	162
Other defence sites	5.3.....	164

**Radiochemical production**

Radiation exposure - radiochemical	6.1.....	173
Amersham	6.2.....	174
Cardiff	6.3.....	175

**Industrial and landfill sites**

Radiation exposure - industrial and landfill	7.1.....	184
Low Level Waste Repository near Drigg	7.2.....	185
Landfill Scotland	7.3.....	186
Landfill England and Wales	7.4.....	186
Landfill East Northamptonshire	7.5.....	187
Landfill Aberdeenshire	7.6.....	187
Whitehaven	7.7.....	188
Non-nuclear gaseous discharges	7.8.....	188
Non-nuclear liquid discharges	7.9.....	189
River Forth and Clyde	7.10.....	190

**Regional monitoring**

Channel Islands	8.1.....	200
Dairies	8.2.....	202
Isle of Man	8.3.....	202
Northern Ireland	8.4.....	203
Diet	8.5.....	206
Air and rain	8.6.....	207
Freshwater Scotland	8.7.....	208
Freshwater England and Wales	8.8.....	209
Freshwater Northern Ireland	8.9.....	209
Radiation exposure - drinking water	8.10.....	210
Groundwater	8.11.....	210
Seawater	8.12.....	211
Sediments	8.13.....	213

# List of Figures

Abbreviated Title	Number .....	Page
<b>Technical summary</b>		
UK <i>total dose</i> from all sources	S.....	11
<b>Introduction</b>		
Dose assessment approach	1.1.....	19
UK <i>total doses</i>	1.2.....	21
UK source specific doses	1.3.....	22
UK sources of waste	1.4.....	24
Potential sites for new nuclear power stations	1.5.....	27
<b>Nuclear fuel production and reprocessing</b>		
Nuclear fuel production and reprocessing – <i>total doses</i>	2.1.....	39
Capenhurst - technetium-99 in grass	2.2.....	39
Springfields - monitoring locations	2.3.....	41
Springfields - <i>total doses</i> and external gamma doses	2.4.....	41
Springfields - discharge and monitoring trends	2.5.....	43
Sellafield - <i>total doses</i> from all sources	2.6.....	46
Sellafield - <i>total doses</i> (gaseous and direct radiation sources)	2.7.....	47
Sellafield - radioactivity in milk	2.8.....	50
Sellafield - technetium-99 in seaweed (historic)	2.9.....	51
Sellafield - technetium-99 in seaweed (recent)	2.10.....	52
Sellafield - monitoring locations in Cumbria	2.11.....	53
Sellafield - monitoring locations	2.12.....	54
Sellafield - carbon-14 in seafood	2.13.....	55
Sellafield - cobalt-60 in seafood	2.14.....	55
Sellafield - technetium-99 in seafood	2.15.....	55
Sellafield - caesium-137 in seafood	2.16.....	56
Sellafield - plutonium-239+240 in seafood	2.17.....	56
Sellafield - americium-241 in seafood	2.18.....	56
Sellafield - caesium-137 in mud	2.19.....	58
Sellafield - plutonium in mud	2.20.....	58
Sellafield - cobalt-60 in mud	2.21.....	58
Sellafield - americium-241 in mud	2.22.....	59
Sellafield - Irish Sea sediment concentrations	2.23.....	60
Sellafield - Irish Sea - dose rates	2.24.....	61
<b>Research establishments</b>		
Research establishments - <i>total doses</i>	3.1.....	95
Dounreay - monitoring locations and seaweed	3.2.....	96
Dounreay - monitoring locations (local)	3.3.....	96
Thames sites - monitoring locations	3.4.....	99
Harwell - liquid discharges	3.5.....	99
Winfrith - monitoring locations	3.6.....	100
Winfrith – liquid discharges	3.7.....	101
<b>Nuclear power stations</b>		
Power Stations - <i>total doses</i> from all sources	4.1.....	112
Caesium-137 in marine sediments	4.2.....	113
Trawsfynydd - caesium-137 in sediments	4.3.....	126
<b>Defence</b>		
Aldermaston - liquid discharges	5.1.....	157
Devonport - liquid discharges	5.2.....	159

**Radiochemical production**

Cardiff - <i>total doses</i> from all sites in Severn Estuary	6.1.....	170
Cardiff - monitoring locations	6.2.....	171
Cardiff - tritium in sediments	6.3.....	172
Cardiff - tritium in seafood	6.4.....	172
Cardiff - carbon-14 in seafood	6.5.....	172

**Industrial and landfill sites**

Landfill monitoring locations	7.1.....	179
Whitehaven - polonium-210 in winkles	7.2.....	181
Whitehaven - polonium-210 in crabs	7.3.....	181
Whitehaven - <i>total doses</i> to seafood consumers	7.4.....	182

**Regional monitoring**

Northern Ireland - monitoring locations	8.1.....	193
Northern Ireland - sediment	8.2.....	194
Chernobyl - caesium-137 in freshwater fish	8.3.....	194
Drinking water monitoring locations	8.4.....	196
Irish Sea - caesium-137 in seawater	8.5.....	197
English Channel - caesium-137 in seawater	8.6.....	197
Irish Sea - tritium in seawater	8.7.....	197
Bristol Channel - tritium in seawater	8.8.....	197
English Channel - tritium in seawater	8.9.....	198
Caesium-137 in seawater	8.10.....	199

# Preface

This report covers sampling and analysis carried out in 2015 for the UK-wide monitoring programmes of the Food Standards Agency, Environment Agency, Scottish Environment Protection Agency, Food Standards Scotland, Natural Resources Wales and the Northern Ireland Environment Agency. The monitoring programmes conducted by these agencies are independent of, and also used as a check on, the nuclear site operators' programmes.

In December 2015, the UK's last remaining Magnox reactor at Wylfa ceased power generation and entered the first stages of decommissioning. This report includes the results of monitoring which continue to take place around Magnox sites as they undergo decommissioning as well as the Advanced Gas Cooled Reactors and Pressurised Water Reactor which continue to generate power. This report

also includes monitoring at sites involved in nuclear fuel production and reprocessing, research establishments, defence establishments, radiochemical production and certain industrial and landfill sites.

The results of these monitoring programmes are used to assess the dose received by members of the public in the vicinity of nuclear licensed sites and industrial and landfill sites. These dose assessments utilize information on diets and occupancy habits of people living near nuclear licensed sites gathered during habit surveys. In 2015 habit surveys were carried out in Bradwell, Harwell, Sizewell, Sellafield in England, and at Chapelcross and Rosyth in Scotland. Radiation doses to people living around nuclear licensed sites from authorised releases of radioactivity were well below the UK national and European limit of 1 millisievert per year in 2015.

# Technical summary

The technical summary is divided into sections to highlight the five main topics within the report. These are:

- Radiation exposures (doses) to people living around UK nuclear licensed sites
- Radioactivity concentrations in samples collected around UK nuclear licensed sites
- External dose rates measured around UK nuclear licensed sites, as a result of exposure to radiation from sediments, etc.
- UK site incidents and non-routine surveys
- Radiation exposures and radioactivity concentrations at other locations remote from UK nuclear licensed sites

## Radiation exposure around UK nuclear licensed sites

In this report we make an assessment of doses to the public near nuclear licensed sites using the results of monitoring of radioactivity in food and the environment, supplemented by modelling where appropriate. The assessments use radionuclide concentrations, dose rates and information on the habits of people living near the sites. Changes in the doses received by people can occur from year to year and are mostly caused by variations in radionuclide concentrations and external dose rates. However, in some years, doses are affected by changes in people's habits, in particular the food they eat, which is reported in habits surveys. The dose quantity presented in this summary is known as the '*total dose*' and is made up of contributions from all sources of radioactivity from man-made processes. Source specific dose assessments are also performed in some cases to provide additional information and as a check on the *total dose* assessment method. *Total dose* is confirmed as a robust measure of exposure.

Figure S and Table S show the assessed *total doses* in 2015 due to the combined effects of authorised/permitted waste discharges and direct radiation for those people most exposed to radiation near all major nuclear licensed sites in the UK. In this report, doses to individuals are determined for those people most exposed to radiation. These results are for comparison with legal limits. The method of calculation involves an assessment for the 'representative person'. This term has the same meaning as 'average member of the critical group' which was used in earlier reports. In this report the term 'representative person' is sometimes shortened to 'person'. Such a person is a hypothetical construct for dose assessment purposes. Reports prior to the one for 2013 referred to an average dose to individuals in a group of people rather than to a single person. The doses are equivalent and comparable.

In 2015, radiation doses from authorised/permitted releases of radioactivity, to adults and children living around nuclear licensed sites, remained well below the UK national and European limit of 1 millisievert (mSv, a measure of dose) per year (see Appendix 3 for explanation of dose units).


The locations where the public received the highest doses in 2015 were the same as those in 2014. They were near Sellafield (0.42 mSv), Amersham (0.14 mSv) and Capenhurst (0.13 mSv). The doses received near Capenhurst and Amersham were dominated by direct radiation from sources on the sites.

In 2015, the representative person in the vicinity of the Sellafield site was a mollusc consumer. The person also consumed significant quantities of other seafood. Their dose of 0.42 mSv includes a contribution of 0.35 mSv from the past discharges from the former phosphate processing plant at Whitehaven and 0.078 mSv related to the discharges of artificial radionuclides by the nuclear industry. The equivalent local seafood consumer in 2014 received a dose of 0.22 mSv (including a contribution of 0.15 mSv and 0.068 mSv related to the former phosphate processing plant and the nuclear industry, respectively). The increase in dose at Sellafield was largely due to an increase in polonium-210 concentrations in locally caught lobsters and crabs. Polonium-210 concentrations in lobster and crab samples continued to be within or close to the expected range due to natural sources in 2015. The largest contribution to dose to seafood consumers in the vicinity of Sellafield was from polonium-210, from the former phosphate processing plant at Whitehaven.

The highest dose near Sellafield was mostly due to historical liquid discharges. The maximum dose at Sellafield for the representative person most affected by pathways related to gaseous discharge and direct radiation sources was 0.008 mSv in 2015, similar to the value for 2014 of 0.009 mSv. The person was a 10 year-old child in 2015 and the dominant contribution to their dose was direct radiation from the site.

In Scotland, the representative person who received the highest dose from authorised releases of radioactivity consumed fish, shellfish and wildfowl on the Dumfries and Galloway coast. The dose in 2015 was 0.038 mSv. Most of this was due to the effects of past discharges from the Sellafield site.

In Wales, the representative person who received the highest dose from authorised releases of radioactivity consumed locally grown food at Trawsfynydd. The dose in 2015 was 0.016 mSv.



**Figure 5.** Total doses in the UK due to radioactive waste discharges and direct radiation, 2015 (Exposures at Sellafield, Whitehaven and Drigg receive a significant contribution to the dose from technologically enhanced naturally occurring radionuclides from previous non-nuclear industrial operations)

**Summary Table S. Total doses due to all sources at major UK sites, 2015<sup>a</sup>**

Establishment	Exposure, mSv <sup>b</sup> per year	Contributors <sup>c</sup>
<b>Nuclear fuel production and processing</b>		
Capenhurst	0.13	Direct radiation
Springfields	0.050	Gamma dose rate over sediment
Sellafield <sup>e</sup>	0.42	Crustaceans, molluscs, <sup>210</sup> Po
<b>Research establishments</b>		
Dounreay	0.010	Gamma dose rate over sediment
Harwell	0.017	Direct radiation
Winfrith	0.014	Direct radiation
<b>Nuclear power stations</b>		
Berkeley and Oldbury	<0.005	Milk, <sup>14</sup> C, <sup>35</sup> S <sup>d</sup>
Bradwell	0.017	Direct radiation
Chapelcross	0.022	Milk, <sup>90</sup> Sr, <sup>241</sup> Am <sup>d</sup>
Dungeness	0.014	Direct radiation
Hartlepool	0.022	Direct radiation, gamma dose rate over sediment
Heysham	0.023	Fish, gamma dose rate over sediment, molluscs, <sup>239/240</sup> Pu, <sup>241</sup> Am
Hinkley Point	0.016	Gamma dose rate over sediment
Hunterston	0.025	Milk, <sup>90</sup> Sr, <sup>241</sup> Am <sup>d</sup>
Sizewell	0.021	Direct radiation
Torness	0.020	Direct radiation
Travsfynydd	0.014	Direct radiation, milk, <sup>241</sup> Am
Wylfa	0.013	Direct radiation
<b>Defence establishment</b>		
Aldermaston and Burghfield	<0.005	Milk, <sup>3</sup> H <sup>d</sup> , <sup>137</sup> Cs <sup>d</sup> , <sup>234</sup> U, <sup>238</sup> U
Barrow	0.051	Gamma dose rate over sediment
Derby	<0.005	Water, <sup>60</sup> Co <sup>d</sup>
Devonport	<0.005	Fish, <sup>14</sup> C, <sup>131</sup> I <sup>d</sup> , <sup>241</sup> Am <sup>d</sup>
Faslane	<0.005	Cattle meat, <sup>137</sup> Cs, <sup>241</sup> Am <sup>d</sup>
Rosyth	0.006	Gamma dose rate over sediment
<b>Radiochemical production</b>		
Amersham	0.14	Direct radiation
Cardiff	<0.005	Gamma dose rate over sediment
<b>Industrial and landfill</b>		
LLWR near Drigg <sup>e</sup>	0.42	Crustaceans, molluscs, <sup>210</sup> Po
Whitehaven <sup>e</sup>	0.42	Crustaceans, molluscs, <sup>210</sup> Po

<sup>a</sup> Includes the effects of waste discharges and direct radiation from the site. May also include the far-field effects of discharges of liquid waste from Sellafield

<sup>b</sup> Committed effective dose calculated using methodology of ICRP-60 to be compared with the dose limit of 1 mSv

<sup>c</sup> Pathways and radionuclides that contribute more than 10% of the total dose. Some radionuclides are reported as being at the limits of detection

<sup>d</sup> The assessed contribution is based on data at limits of detection

<sup>e</sup> The doses from man-made and naturally occurring radionuclides were 0.078 and 0.35 mSv respectively. The source of man-made radionuclides was Sellafield; naturally occurring ones were from the phosphate processing works near Sellafield at Whitehaven. Minor discharges of radionuclides were also made from the LLWR near Drigg site into the same area

## Habits surveys near UK nuclear licensed sites

In 2015, the regular programmes of habits surveys around nuclear licensed sites continued. These give site-specific information on diets and occupancy habits of people near nuclear licensed sites. Surveys were carried out at Bradwell, Harwell, Sizewell and Sellafield in England and at Chapelcross and Rosyth in Scotland. The findings were used to confirm the adequacy of current monitoring programmes or strengthen and update them with a better representation of relevant pathways, and to improve the assessment of doses to members of the public near nuclear licensed sites.

## Radioactivity concentrations in samples collected around UK nuclear licensed sites

This section summarises any changes in concentrations of radioactivity in food or the environment, given in becquerels per kilogramme ( $\text{Bq kg}^{-1}$ ) or becquerels per litre ( $\text{Bq l}^{-1}$ ).

A revised UK Radioactive Discharge Strategy was published in 2009. This describes how the UK will implement the commitments in the OSPAR Radioactive Substances Strategy (RSS) on radioactive discharges to the marine environment of the North-East Atlantic. The UK Strategy has resulted in substantial reductions in radioactive discharges and in nuclear licensed sites producing action plans to further reduce discharges. From a regulatory perspective, the Environment Agency, Natural Resources Wales and the Scottish Environment Protection Agency (SEPA) have continued to support the Strategy and in 2015, the Environment Agency and Natural Resources Wales issued new authorisations/permits, or varied existing ones, at five sites, Cardiff, Harwell, Low Level Waste Repository (near Drigg), Sellafield and Winfrith, resulting in one or more of: strengthened conditions, reduced limits or new routes for disposing of radioactive waste.

There were no major variations in environmental concentrations of radioactivity in 2015 compared to those in 2014.

During the past decade, discharges from GE Healthcare Limited at Cardiff have continued to decline. This has led to a downward trend in concentrations of tritium in fish and molluscs near the site.

During 2015, discharges of technetium-99 from Sellafield continued to be low, following the introduction of abatement technology in previous years. Discharges are expected to remain low in the future. Technetium-99 from Sellafield can be detected in the Irish Sea, in Scottish waters and in the North Sea. However, concentrations of

technetium-99 have shown a strong trend downward from their peak in 2003.

Marine sediment samples are a useful indicator of trends in the environment. People who spend time on beaches can be exposed to radiation through the radionuclide content of the sediments. Near Sellafield, the environmental concentrations of most radionuclides have declined substantially over the last 25 years. Small increases in plutonium isotopes and americium-241 have been observed in mud samples from the Ravenglass estuary near Sellafield. However, these have had little or no effect on radiation exposures.

On occasion, the effects of non-nuclear sites discharge are detected at low levels by the routine monitoring programme for nuclear licensed sites. In 2015, iodine-131 was detected in marine samples at several nuclear licensed sites. The source of the iodine-131 is not known with certainty but a likely cause was the therapeutic use of this radionuclide in local hospitals. The concentrations were of low radiological significance.

An enhanced monitoring programme at Bradwell was undertaken in 2015 in response to local views whilst onsite treatment of Intermediate Level Waste was being undertaken. The results show no change in levels of radioactivity in the environment.

## Dose rates from around UK nuclear licensed sites

Sediments in intertidal areas can make a significant contribution to the total radiation exposure of members of the public. For this reason, external doses are recorded by measuring dose rates. These 'external doses' are included in the assessment of doses to the public where they are higher than background levels. Background levels are subtracted in dose assessments.

There were no major changes in external dose rates in intertidal areas in 2015 compared with 2014. At most locations, the external dose rates were close to background levels. Levels were higher in some estuaries near Sellafield (up to twice the background rate) and in the Ribble Estuary.

## UK nuclear licensed site incidents and non-routine surveys

During 2015, as a result of an ongoing programme of monitoring by the operator, radioactive items (particles\*, including contaminated pebbles/stones) from Sellafield were detected on Cumbrian coastline beaches and removed (339 in financial year 2015/16). Public Health England (PHE) has provided advice that the overall health risks for beach users from radioactive objects on beaches near Sellafield are very low and significantly lower than other risks that people accept when using the beaches. A programme of work is in place to address the remaining uncertainties in the origins, fate and effects of the particles with the aim of returning the work to a routine monitoring programme only by about 2018.

At Dounreay, the comprehensive beach monitoring programme continued for fragments of irradiated nuclear fuel (particles) and further particles were recovered from local beaches. Fishing restrictions under the Food and Environment Protection Act (FEPA) 1985 are still in force.

'Special' (or *ad hoc*) sampling related to nuclear licensed site operation is undertaken at sites when the need arises, for example when increases in discharges are reported. No such need arose in 2015.

## Radiation doses and levels at other locations

Additional monitoring was undertaken in the UK and surrounding seas to study the effects of (i) overseas incidents, (ii) non-nuclear sites and (iii) regional variation in levels of radioactivity across the UK.

## Overseas incidents

The accident at Fukushima Dai-ichi nuclear power station in Japan in March 2011 resulted in significant quantities of radioactivity being released to air and sea.

Controls on food imports from Japan took two forms, and these controls continued in 2015. The European Commission (EC) implemented controls on the import of food and feed originating in or consigned from Japan. All food and feed imported from Japan had to be certified by the Japanese authorities, with the exception of certain alcoholic beverages and (since March 2014) tea. From January 2016, only certain foods specified in the controls

---

\* "Particle" is a term used in this report which encompasses discrete radioactive items which can range in radioactivity concentration, size and origin. "Particles" include radioactive scale, fragments of irradiated nuclear fuel, incinerated waste materials, typically less than 2mm in diameter, and larger radioactive artefacts (e.g. dials) and stones which have radioactive contamination on their surface. Particles are not physically the same at each of the sites mentioned, but can be compared according to the hazard posed.

continue to require certification. In addition, a percentage of Japanese imports into the European Union (EU) were monitored at ports of entry. None of the imports to the UK in 2015 have contained activity exceeding the maximum permissible levels; most results have been below the limits of detection. The public doses received due to the imports were of negligible radiological significance.

Direct monitoring of fish in relation to the effects of the 1986 Chernobyl accident ceased in upland lakes in 2014 as part of the overall review of the FSA's monitoring programme. All restrictions on moving, selling and slaughtering sheep in upland areas of the UK were removed in 2012.

Food imported into the UK may contain radioactive contamination from Chernobyl and other known or unknown sources. A monitoring system is in place to detect radioactivity in consignments. In 2015, no significant radioactivity was detected at entry points and there was no need to introduce food safety controls on any consignments.

## Non-nuclear sites

In the past, liquid slurry containing thorium and uranium was discharged into the Irish Sea from a phosphate plant near Whitehaven in Cumbria. This was a practice that generated what is sometimes known as 'Technologically enhanced Naturally Occurring Radioactive Material' (TNORM). Where discharges of TNORM occur, this can lead to an increase in the concentrations of naturally occurring radionuclides in the environment. This site stopped operating at the end of 2001, was decommissioned in 2002 and the plant has subsequently been demolished. Concentrations of naturally occurring radionuclides in fish and shellfish near Whitehaven have been found to be higher than the maximum expected ranges due to natural sources. Concentrations of natural radionuclides have declined in the last 10 years, so that in recent years the concentrations were very close to natural background, making any increase due to the past discharges difficult to distinguish. Estimates of the concentrations of naturally occurring radionuclides in seafood caused by past discharges from the site have been made by subtracting the expected natural concentration of these radionuclides in UK seafood from the measured levels. Polonium-210, which is naturally occurring, is present in some seafood samples at slightly above background levels. Polonium-210 (and lead-210) are important radionuclides in that small changes in levels above background significantly influence the dose contribution from these radionuclides and similarly the value of the estimated dose. The representative person in the area who consumed large amounts of seafood was estimated to receive a dose of 0.42 mSv, with about 80 per cent from polonium-210. The dose includes a contribution from the effects of discharges from the adjacent sites at

Sellafield and, to a much lesser extent, at the Low Level Waste Repository (LLWR) near Drigg.

Concentrations of tritium were found in leachate from some landfill sites, at levels that were of very low radiological significance. There are several disposal routes for radioactive waste to landfill that could contain tritium, for example, from hospitals and industrial sites, and due to disposals of gaseous tritium light devices (such as fire exit signs).

Work to address the radioactive contamination at Dalgety Bay is ongoing. Public protection measures have been established and these were maintained during 2015 and into 2016. A continuing monthly beach monitoring and particle recovery programme was begun in 2012 by a contractor working on behalf of the MoD. The fence demarcating the area where the highest activity particles were discovered remains in place, as well as the information signs advising the public of the contamination and precautions to be taken. In addition, the FEPA Order issued by the FSA in Scotland (now FSS) prohibiting the collection of seafood from the Dalgety Bay area remains in force. Following the publication of the risk assessment together with the appropriate persons report in 2013, COMARE recommended that effective remediation of the affected area is undertaken as soon as is possible. The MoD has progressed with addressing the contamination by initially publishing its Outline Management Options Appraisal Report in January 2014 followed by the publication in July 2014 of its broad management strategy and timescale for implementation of its preferred management option. PHE, at the request of SEPA, has provided advice on target levels of radioactive contamination for Dalgety Bay following any remediation of the affected area.

Further details can be found in Section 7.6 of this report and on the Radioactive Substances pages of SEPA's website ([www.sepa.org.uk](http://www.sepa.org.uk)). As work in this area is ongoing, an update will be provided in next year's RIFE report.

## Regional monitoring

Monitoring artificial radioactivity in Northern Ireland showed that consumer doses were all less than 1 per cent of the annual limit of 1 mSv for members of the public. A survey on the Channel Islands confirmed that doses due to discharges from the French reprocessing plant at La Hague and other local sources were less than 1 per cent of the limit.

Food in people's general diet and sources of public drinking water were analysed across the United Kingdom. Results showed that artificial radionuclides only contributed a small

proportion of the total public radiation dose in people's general diet.

The distribution of radionuclides in coastal seas away from nuclear licensed sites continues to be monitored. This supports the UK's marine environmental policies and international treaty commitments. Government research vessels are used in the sampling programme and the results have been used to show trends in the quality of the UK's coastal seas. These surveys, together with the results of monitoring at nuclear licensed sites, contribute to the data collected by the OSPAR Commission. They also help to measure progress towards the UK Governments' objectives for improving the state of the marine environment.

Disposal of dredge material from harbours and other areas is licensed under the Marine and Coastal Access Act (MCAA), 2009. In 2015, a request was received by the Marine Management Organisation (MMO) for additional licenses for the disposal of approximately 80,000 tonnes of dredged material at sea from Maryport in Cumbria. A specific assessment of doses to workers and members of the public was conducted for the disposal of the dredge material. The combined dose (from artificial and naturally-occurring radionuclides) to individual members of the crew and public was within the IAEA *de minimis* criteria of 0.010 mSv per year

## The monitoring programmes and further research

The monitoring programmes in this report involved four specialist laboratories working together, each with rigorous quality assurance procedures, and a wide range of sample collectors throughout the United Kingdom. They were organised by the environment agencies, FSA and FSS and they are independent of the industries discharging radioactive wastes. The programmes include monitoring on behalf of the Scottish Government, Channel Island States, the Department of Agriculture Environment and Rural Affairs (formerly, the Department of the Environment Northern Ireland), the Department of Business, Energy and Industrial Strategy (BEIS) (formerly the Department of Energy and Climate Change (DECC)), Defra, the Isle of Man Government, Natural Resources Wales (NRW) and the Welsh Government. Overall, around 10,000 analyses and dose rate measurements were completed in 2015.

The results of the analysis of food samples collected near nuclear licensed sites in England and Wales are published on FSA's website ([www.food.gov.uk](http://www.food.gov.uk)). More information about all programmes described in this report is available from the sponsoring agencies. Their contact details can be found on the inside front and back covers of this report.



# 1. Introduction

This section (i) describes the purpose and scope of the UK monitoring programmes for radioactivity in food and the environment, (ii) provides a summary of the key results in terms of radiation exposures at each major industrial site in 2015 and (iii) gives an overview of the main sources of radiation in a regulatory context.

## 1.1 Purpose and scope of the monitoring programmes

In England and Wales, FSA conducts food monitoring, whilst the Environment Agency carries out environmental and dose rate monitoring\*. In Scotland, SEPA carries out food, environmental and dose rate monitoring, working closely with FSS on its programme, and in Northern Ireland this is carried out by the Northern Ireland Environment Agency (NIEA). Surveillance of imports through points of entry continued in 2015. The regular programme of monitoring of drinking water, air and rain continued on behalf of BEIS (formerly DECC), NIEA and the Scottish Government. The FSA and SEPA (as part of the joint SEPA/FSS monitoring programme) also carry out UK monitoring of milk and canteen meals that are collected remotely from nuclear licensed sites. The marine environment of the whole of the British Isles away from nuclear licensed sites is monitored for Defra.

FSA was given responsibility for food safety throughout the UK under the Food Standards Act 1999. The Food Standards Act 1999 was amended by the Food (Scotland) Act 2015 and responsibility for food safety in Scotland was transferred to the newly formed FSS on 1st April 2015. The Environment Agency, NRW, NIEA and SEPA, referred to together as the environment agencies in this report, are responsible for environmental protection in England, Wales, Northern Ireland and Scotland, respectively. The Environment Agency and NRW regulate radioactive waste disposal under the Environmental Permitting (England and Wales) Regulations 2010 (EPR 10), (United Kingdom - Parliament, 2010a), whilst in Scotland and Northern Ireland, SEPA and NIEA regulate radioactive waste disposal under the Radioactive Substances Act 1993 (RSA 93) (United Kingdom - Parliament, 1993). The Environment Agency and SEPA also have broader responsibilities under the Environment Act 1995 (United Kingdom - Parliament, 1995a) for environmental protection and determining general concentrations of pollution in the environment.

\* *The Environment Agency has an agreement with NRW to undertake some specific activities on its behalf in Wales including some environmental monitoring and aspects of radioactive substances regulation.*

### Key points

- The RIFE report represents collaboration by the environment agencies, FSA and FSS across the UK, independent of industry
- Provides an open check on food safety and the public's exposure to radiation in conformity with the EU Basic Safety Standards Directive
- Monitoring programme results support the UK meeting its international treaty obligations
- Dose results are summarised for major industrial sites; all doses were below the legal limit in 2015

The monitoring programmes have several purposes. Ongoing monitoring helps to establish the long-term trends in concentrations of radioactivity over time within the vicinity of, and at distance from, nuclear licensed sites. The results are also used to confirm the safety of the food chain. Monitoring the environment provides indicators of radionuclide dispersion around each nuclear site. Environmental and food results are used to assess dose to the public to confirm that the controls and conditions placed in the authorisations/permits provide the necessary protection and to ensure compliance with statutory dose limits. Most of the monitoring carried out and presented in this report concerns the local effects of discharges from nuclear licensed sites in the UK. Other work includes monitoring of food imports from Japan to confirm that controls on the effects of the Fukushima Dai-ichi accident in 2011 by the Japanese authorities are working adequately. Monitoring of food and the environment remote from nuclear licensed sites is also carried out, giving information on background concentrations of radionuclides; these data are reported to the EC. Guidance on planning and implementing routine environmental programmes has been published (Environment Agency, FSA and SEPA, 2010).

The RIFE report and the associated monitoring programmes conform to the requirements in Article 66 of the Euratom Directive laying down basic safety standards for protection against the dangers arising from exposure to ionising radiation (EC, 2014). Specifically, it provides estimates of doses to members of the public from authorised practices and enables such results to be made available to stakeholders.

FSA completed a public consultation exercise following the risk-based review into the way it monitors radioactivity in food in June 2013 (FSA, 2012a and 2013). This has

resulted in a revised monitoring programme taking effect in 2014 with reductions in sampling and analysis of some foods representing a very low radiological risk. An interim review, following the first year of the new programme, was presented to the FSA's open Board Meeting on the 16 March 2016. More information can be found at: <http://www.food.gov.uk/about-us/our-board/meetings/2016/010116/board-meeting-agenda-16-march-2016>.

In 2012/2013 SEPA undertook a review of its environmental radioactivity monitoring programme coordinated through SEPA's Environmental Radioactivity Monitoring Task Team (ERM-TT). The ERM-TT includes representatives from SEPA, Scottish Government, PHE, FSA, FSS and Scottish Natural Heritage. The review took into account new habits data, recent environmental monitoring data and the joint guidance published by SEPA, FSA and the Environment Agency concerning planning and implementing routine environmental radiological monitoring programmes (Environment Agency, FSA and SEPA, 2010). Minor modifications were made to the programme which came into effect on 1st January 2014.

In 2014, the Environment Agency began a review of their environmental monitoring around nuclear sites. The review is to ensure the programmes are appropriate and are consistent with advice in the joint Agency technical guidance (Environment Agency, FSA and SEPA, 2010). The majority of the review was completed during 2015, with some further minor follow up work continuing into 2016. The overall outcome was an adjustment and consolidation of the monitoring around some sites. The Environment Agency's monitoring programme for 2016 will be changed to reflect the review outcomes.

Appendix 1 is in a file accompanying the main report. It gives details of methods of sampling and analysis and explains how results are interpreted in terms of public radiation exposures. A report of trends in monitoring data and doses for 2004 – 2008 has been published (Environment Agency, FSA, NIEA and SEPA, 2010b). A summary of the assessment approach and current trends in doses are given in Section 1.2.

The analytical science for the monitoring programmes was carried out by a number of UK laboratories, including those listed below. These laboratories also carried out most of the sample collection for the programmes.

- Centre for Environment, Fisheries & Aquaculture Science (Cefas)
- Environmental Scientifics Group (ESG)
- Public Health England (PHE)
- LGC Limited (formerly Laboratory of the Government Chemist)

## 1.2 Summary of doses

### 1.2.1 The assessment process

The majority of the monitoring was carried out to check the effects of discharges from nuclear and non-nuclear operators on people's food and their environment. The results are used to assess doses to the public that can then be compared with the relevant dose limits. The dose assessments are retrospective in that they apply to 2015, using monitoring results for that year. The radioactivity concentrations and dose rates reported include the combined radiological impact of all discharges made up to the time of sampling.

In this report, two main types of retrospective dose assessment are made (see Figure 1.1). The first type of assessment is more complete in considering the effects of gaseous and liquid discharges of radioactive waste together and additionally includes exposure to direct radiation from nuclear licensed sites. It gives an estimate of *total dose* to people around the nuclear licensed sites and *total dose* is presented as the primary dose quantity. Direct radiation can be a significant contributor to dose close to operating power stations or where radioactive materials are stored. The regulation of direct radiation is the responsibility of the Office for Nuclear Regulation (ONR). Operators of nuclear licensed sites provide estimates of direct radiation doses to ONR which are made available for use in these assessments (Table 1.1). The *total dose* assessments use recent habit survey data which has been profiled using an agreed method (Camplin *et al.*, 2005).

The second type of assessment focuses on specific sources and their associated pathways. It serves as a check on the adequacy of the *total dose* method and offers additional information for key pathways. The sum of the doses from specific sources does not give the same result as the assessment of *total dose* from all sources. This is because the assessment methods use different ways of defining the most exposed people.

Both types of assessment consider the people in the population who are most exposed to radiation. These results are for comparison with legal limits. The method of calculation involves an assessment for the 'representative person'.

The calculated doses are compared with the dose limit for members of the public of 1 mSv per year. Dose assessments for exposure to skin are also made at some sites and compared with the relevant skin dose limit. The approaches used are for relatively widespread contamination in food and the environment where the probability of encounter/consumption is certain. These methods are not appropriate for exposure to small radioactive particles where the chance of encounter is a relevant factor to be considered (Dale *et al.*, 2008). All dose limits are based on recommendations made by the

Primary purpose	Assess dose from main sources of exposure at each site for comparison with 1 mSv limit			
Types of assessment	Total dose	Source specific dose		
Sources considered	Gaseous discharges Liquid discharges Direct radiation from site	Gaseous discharges	Liquid discharges	Direct radiation (dose estimates provided by ONR)
Habits data e.g. food consumption rates or occupancy of beaches	Define usage of pathways relating to all sources at site	Define usage of pathways relating to gaseous discharges at site	Define usage of pathways relating to liquid discharges at site	
Monitoring data	Collate monitoring data for relevant pathways e.g. radionuclide concentrations in food or dose rates on beaches	Collate monitoring data for relevant pathways e.g. radionuclide concentrations in food	Collate monitoring data for relevant pathways e.g. radionuclide concentrations in food or dose rates on beaches	
Dose calculations	Calculate dose from all sources to individuals who may represent those most exposed	Calculate dose from gaseous discharges to people representing those most exposed	Calculate dose from liquid discharges to people representing those most exposed	
	Select the highest dose for the person representing the most exposed			
Dose quantity	Total dose	Dose from gaseous discharges	Dose from liquid discharges	Dose from direct radiation

**Figure 1.1.** The dose assessment process for major nuclear sites

ICRP (ICRP, 1991) and are consistent with EU legislation (EC, 2014).

An additional comparison can be made with doses from natural radioactivity. The UK average is 2.2 mSv per year, with a range across counties from 1.5 mSv per year to 7 mSv per year (Watson *et al.*, 2005).

Collective doses are beyond the scope of this report. They are derived using modelling techniques. The EC has published an assessment of individual and collective doses from reported discharges from nuclear power stations and reprocessing sites for the gaseous and liquid waste disposals in the years 2004 to 2008 (Jones *et al.*, 2013).

Radiation exposures to some specific groups of workers are included in the assessment of doses from nuclear licensed sites. These are workers who may be exposed incidentally, but do not work specifically with ionising radiation. These include fishermen, farmers, sewage workers, nature wardens, etc. It is appropriate to compare their doses to the dose limit for members of the public (Allott, 2005). Doses to workers who are involved with ionising radiation and receive a dose from their work should be assessed as part of their employment.

## 1.2.2 Total dose results for 2015

The results of the assessment for each site are summarised in Table 1.2 (see also Figure S and Table S in the Technical Summary). These data are presented in three parts. The representative person receiving the highest doses from the pathways predominantly relating to gaseous discharges

and direct radiation are shown in part A and those for liquid discharges in part B. Occasionally, the people receiving the highest doses from all pathways and sources are different from those in A and B. Therefore, this case is presented in part C. The major contributions to dose are provided. The use of radionuclide concentrations reported at the limits of detection provide an upper estimate of doses calculated for pathways based on these measurements. The full output from the assessment for each site can be provided by contacting one of the agencies listed on the inside cover of the report.

In all cases, doses estimated for 2015 were less than the limit of 1 mSv for members of the public. The people most affected from gaseous discharges and direct radiation varied from site to site but the dominant pathway was often direct radiation where it was applicable. The people most affected from liquid discharges were generally adult consumers of seafood or occupants over contaminated sediments.

The representative person, who received the highest *total dose*, consumed molluscan shellfish at high-rates (and other seafood) near Sellafield. The *total dose* (from all sources) at this site is combined with the effects of all local sources, including specifically the effects of historic discharges of natural radionuclides from the former phosphate processing plant at Whitehaven and (to a lesser extent) the effects of discharges from LLWR (near Drigg). The representative persons, receiving the next highest *total doses*, were local inhabitants living near the Capenhurst and Amersham sites; these doses were almost entirely due to direct radiation from the sites.

### 1.2.3 Total dose trends

A time-series of *total dose* from 2004 - 2015 is shown in Figure 1.2 (Table 1.3 gives numerical values). Many sites showed a downward trend in *total dose* over this period. Changes in direct radiation dominated the inter-annual variation at most of the power station sites, and small fluctuations in external dose rates had relatively large effects at some sites where high rates of intertidal occupancy were recorded. Following the cessation of power production by Magnox reactors at Dungeness and Sizewell, the effect has been a reduction in direct radiation at these sites. The most significant trend in *total dose* due to discharges of waste was for high-rate consumers of seafood on the Cumbrian coast near Sellafield, Whitehaven and the LLWR near Drigg. In this case, the overall downward trend in *total dose* broadly followed the general downward trend in concentrations of naturally occurring and artificial radionuclides from non-nuclear and nuclear sources, respectively. Year to year changes were also influenced by changes in consumption and occupancy characteristics of local people and the natural variability in radionuclide concentrations in food and the environment. Indeed, in 2015, the dose to these people increased due to changes in concentrations of polonium-210 in local seafood.

At Cardiff, there has been an overall downward trend in *total dose* which is partly due to reductions in discharges of tritium and carbon-14 to sea. The recent increase in *total dose* at this site was due to higher carbon-14 concentrations in milk in 2013. The *total dose* observed at Dounreay in recent years has decreased from the peak value in 2008 due to changes in caesium-137 concentrations in game meat and the type of game meat sampled. The reduction in *total dose* at Heysham, Hinkley Point and Springfields was largely due to findings from new habits surveys in 2011, 2010 and 2012 respectively.

### 1.2.4 Source specific dose results for 2015

The results of the source specific assessments for the main industrial sites in the UK are summarised in Table 1.4 and Figure 1.3. The focus for these assessments is the effect of gaseous or liquid waste discharges, unlike that for *total dose* which also includes all sources including the effect of direct radiation.

The most significant exposures were found at the LLWR near Drigg, at Sellafield and at Whitehaven where seafood consumption dominated, and at Barrow and at Springfields where external exposure on houseboats dominated. At the LLWR near Drigg, at Sellafield and at Whitehaven the majority of the dose was from the legacy of historical discharges from Sellafield and from non-nuclear industrial operations resulting in technologically enhanced levels of natural radionuclides. The most important pathways and radionuclides at each site were similar to those found

for *total dose* if the effect of direct radiation is taken into account. At Barrow and Springfields the dose was largely due to activity in sediments beneath houseboats, also from historical discharges from Sellafield.

Although some source specific doses were estimated to be higher than *total doses*, the reasons for this are understood and relate to the different assumptions of the two assessment methodologies. The assumptions used for source specific assessments are conservative with respect to adding together the effects of consumption of different foods. The assumptions used for *total dose* assessments are more realistic and the results confirm the adequacy of the *total dose* approach of assessment. Radiation doses to adults and children, calculated using the source specific method, were all found to be well below the national and European limit of 1 mSv per year.


### 1.2.5 Protecting the environment

The main focus of this report is on the protection of people, but the protection of wildlife and the environment is also relevant. ICRP in its 2007 recommendations concluded that there is a need for a systematic approach for the radiological assessment of non-human species to support the management of radiation effects in the environment (ICRP, 2007). In pursuit of this aim, ICRP has considered the use of a set of Reference Animals and Plants (RAPs) (ICRP, 2008) and have published their aims in terms of environmental protection, that is (i) prevention or reduction of the frequency of deleterious radiation effects on biota to a level where they would have a negligible impact on the maintenance of biological diversity, (ii) the conservation of species and the health and status of natural habitats, communities and ecosystems (ICRP, 2014).


In the UK, the current legislative measures relevant to the protection of wildlife from radiation are the Birds and the Habitats Directives, on the conservation of wild birds (CEC, 2009) and on the conservation of natural habitats and wild flora and fauna (CEC, 1992), and are implemented through the Habitats regulations.

Under the Habitats Regulations, the Environment Agency, NRW and SEPA review new and existing authorisations/permits to ensure that they do not have an adverse effect on the integrity of Natura 2000 habitat sites. When a new authorisation/permit to discharge or dispose of radioactive waste is issued, or one is varied, the applicant is required to make an assessment of the potential impact of the discharges on reference organisms that represent species which may be adversely affected.

The Environment Agency has assessed the dose rates to reference organisms and feature species for authorised discharges under RSA 93 and, since April 2010, EPR10 (Environment Agency, 2009a). Environmental concentrations were predicted using appropriate dispersion models and the data were used to calculate the dose rates.



**Figure 1.2.** Total doses around the UK's nuclear sites due to radioactive waste discharges and direct radiation (2004-2015). (Exposures at Sellafield/Whitehaven/LLWR receive a significant contribution to the dose from technologically enhanced naturally occurring radionuclides from previous non-nuclear industrial operations)



**Figure 1.3.** Source specific doses in the UK, 2015 (Exposures at Whitehaven and Sellafield receive a significant contribution to the dose from technologically enhanced naturally occurring radionuclides from previous non-nuclear industrial operations)

The assessment concluded that, for all but two of the habitat sites, dose rates to the worst affected organisms were less than an agreed threshold of 40  $\mu\text{Gy h}^{-1}$ . Hence, there was no significant impact on the integrity of these habitat sites. The two habitat sites, with the potential for dose rates to the worst affected organism to be greater than the agreed threshold, were the Drigg coast and the Ribble and Alt Estuaries. A detailed assessment has been carried out for the Drigg coast using monitoring data and this confirmed there was no indication of significant impact from ionising radiation on the sand dune biota (Wood *et al.*, 2008). A detailed assessment was also carried out for the Ribble and Alt estuaries using monitoring data and taking into account new discharge limits for the Springfields site which came into force in 2008 (Environment Agency, 2009b). This assessment concluded that the dose rate to the worst affected organism was less than the agreed threshold and hence there was no significant impact on the integrity of this habitat site.

SEPA has carried out a Pressures and Impacts Assessment from radioactive substances on Scotland's water environment. The study concluded that there was no adverse impact on the aquatic environment as a result of authorised discharges of radioactive substances, although it recognised that there may be a need for further data to support this conclusion. A report of the study is available from SEPA. SEPA has included a specific habitats assessment in any new authorisation granted by the agency.

### 1.2.6 Food irradiation

Food irradiation is a processing technique where food is exposed to ionising radiation in a controlled manner. Irradiation may be used to eliminate or reduce food-borne pathogenic organisms, extend shelf life by delaying food from rotting or developing mould, and prevent certain food products from ripening, germinating or sprouting. Irradiation may also be used as a phytosanitary measure to rid imported plants or plant products of organisms which may be harmful to domestic flora. The ionising radiation produces free radicals, which interact within the food to produce the desired effect. It does not make the food radioactive. The ionising radiation is either generated by machine, as is the case for electron beams or x-rays, or produced by the radioactive decay of caesium-137 or cobalt-60 (both unstable isotopes whose decay produces gamma radiation).

Food irradiation has been permitted in the UK since 1990, and UK legislation was amended in 2000 to implement two European Directives on food irradiation (CEC, 1999a, b). These amendments were consolidated into a single Regulation in each country of the UK in 2009 as part of the FSA programme of regulatory simplification to reduce administrative burden. In 2010, the Regulations

were amended to update the lists of approved food irradiation facilities.

In the UK, one facility in England is licensed to irradiate a range of dried herbs and spices and it is inspected by the FSA. The facility was not inspected during 2015 as no food has been treated for several years. The FSA continues to monitor the situation and the requirement to inspect this facility is kept under review.

## 1.3 Sources of radiation exposure

### 1.3.1 Radioactive waste disposal from nuclear licensed sites


Nuclear licensed sites in the UK discharge radioactive waste as liquid and/or gas as part of their operations. In addition, solid Low Level Waste (LLW) from nuclear licensed sites can be transferred to the Low Level Waste Repository (LLWR) near Drigg for disposal. Solid LLW from Dounreay can be transferred to the new Dounreay LLW Facility which began accepting waste for disposal in April 2015. These discharges and disposals are regulated by the environment agencies under RSA 93 or EPR 10\*.

Figure 1.4 shows the nuclear licensed sites that produce waste containing artificial radionuclides. Nuclear licensed sites are authorised to dispose of radioactive waste (United Kingdom - Parliament, 1993). They are also subject to the Nuclear Installations Act (United Kingdom - Parliament, 1965). The monitoring programmes reported here include studies at each of these sites. Discharges of radioactive waste from other sites such as hospitals, industrial sites and research establishments are also regulated under RSA 93 or EPR 10 but are not subject to the Nuclear Installations Act. Occasionally, these monitoring programmes detect radioactivity in the environment as a result of these discharges. For example, iodine-131 from hospitals is occasionally detected in some river and marine samples. Small amounts of very low level solid radioactive waste are disposed of from some non-nuclear sites. There is also a significant radiological impact due to the legacy of past discharges of radionuclides from non-nuclear industrial activity that also occur naturally in the environment. This includes radionuclides discharged from the former phosphate processing plant at Whitehaven, and so monitoring is carried out near this site. Discharges from other non-nuclear sites are generally considered insignificant in England and Wales and so monitoring to protect public health is not usually carried out by the environment agencies, although some routine monitoring programmes are undertaken. In Scotland, SEPA undertake

---

\* *In England and Wales, the term 'authorisation' has been replaced by 'permit' with EPR 10 taking effect from 6th April 2010. In this report 'permit' has been used to apply to all sites in England and Wales irrespective of whether the period considered includes activities prior to 6th April 2010. 'Authorisation' remains the relevant term for Scotland and Northern Ireland.*

**Figure 1.4.** Principal nuclear site sources of radioactive waste disposal in the UK, 2015 (Showing main initial operation. Some operations are undergoing decommissioning)



routine sampling in the Firth of Clyde and at landfill sites to assess the impact of the non-nuclear industry on the environment. Additionally, to ensure the doses from combined discharges to a sewer network are assessed properly, SEPA periodically undertakes intensive sampling at major sewage treatment plants to monitor the combined discharges from the non-nuclear industry.

Appendix 2 gives a summary of the discharges of liquid and gaseous radioactive waste and disposals of solid radioactive waste from nuclear licensed establishments in the UK during 2015. The tables also list the main discharge and disposal limits that are specified or, in the case of the MoD, administratively agreed. In 2015, discharges and disposals were all below the limits. Section 7 gives information on discharges from non-nuclear sites.

The discharge limits are set through an assessment process, which either the operator or the relevant environment agency can initiate. In support of the process, prospective assessments of doses to the public are made assuming discharges at the specified limits. Discharge limits are set so that doses to the public will be below the source and site dose constraints of 0.3 and 0.5 mSv per year respectively if discharges occurred at the limits (Environment Agency, SEPA, NIEA, HPA and FSA, 2012). The determination of discharge limits also takes into account the dose due to consumption of food. During the determination of the limits, the effect of the planned discharges on the environment and wildlife is also considered. In addition, the regulations require Best Available Techniques or Best Available Technology (BAT), under EPR 10, to be used to further minimise discharges.

The principles of Best Practicable Means (BPM) are applied in Scotland (SEPA, 2012a).

The discharges and disposals made by sites are generally regular throughout the year. However, from time to time there may be unplanned events that cause unintended leakages, spillages or other emissions that are different to the normal or expected pattern of discharges. These events must be reported to the environment agencies and may lead to follow up action, including reactive monitoring by the site, the environment agencies or FSA. In cases where there has been a breach of limits, or if appropriate actions have not been undertaken to ensure discharges are as low as possible, regulatory action may be taken. Where monitoring took place because of these events, the results are presented and discussed in the relevant site text later in this report. Appendix Table A2.4 summarises the types of events that took place in 2015.

### 1.3.2 International agreements, the UK Discharge Strategy and new nuclear power stations

This section gives information on the context of UK radioactive discharges as they relate to international agreements and the future building of new nuclear power stations. The UK has ratified the Convention for the Protection of the Marine Environment of the North-East Atlantic (the 'OSPAR Convention'). This provides a framework for preventing and eliminating pollution in the north-east Atlantic, including the seas around the UK (OSPAR, 2000a). The OSPAR Convention replaced the separate Oslo and Paris Conventions.

In July 1998, the Ministers of the UK Government agreed a long-term Radioactive Substances Strategy (RSS) and signed the Sintra Statement which included the following commitment (OSPAR, 1998):

“We shall ensure that discharges, emissions and losses of radioactive substances are reduced by the year 2020 to levels where the additional concentrations in the marine environment above historic levels, resulting from such discharges, emissions, losses, are close to zero.”

In July 2002, a UK Strategy for Radioactive Discharges was published (Defra, 2002). This described how the UK would implement the agreements reached at the 1998 and subsequent meetings of OSPAR. The aims of the Strategy related to liquid wastes from the major sources, primarily the nuclear industry, and not to gaseous or solid wastes.

Results of a public consultation to update this Strategy were published in 2009 (DECC, 2009). BEIS and the Devolved Administrations have issued a revised Strategy (DECC, Department of the Environment Northern Ireland, the Scottish Government and Welsh Assembly Government, 2009).

The new Strategy was built on the 2002 publication, and expanded its scope to include aerial, as well as liquid discharges, from decommissioning as well as operational activities, and from the non-nuclear as well as the nuclear industry sectors. It also includes considerations of uncertainties associated with discharges from new nuclear power stations, the possible extension of the operational lives of some of the existing nuclear power reactors, and discharges arising from decommissioning activities. The objectives of this revised Strategy are:

- To implement the UK's obligations, rigorously and transparently, in respect of the OSPAR RSS intermediate objective for 2020
- To provide a clear statement of Government policy and a strategic framework for discharge reductions, sector by sector, to inform decision making by industry and regulators

The expected outcomes of the UK Strategy are:

- Progressive and substantial reductions in radioactive discharges, to the extent needed to achieve the sectoral outcomes, while taking into account the uncertainties
- Progressive reductions in concentrations of radionuclides in the marine environment resulting from radioactive discharges, such that by 2020 they add close to zero to historic levels
- Progressive reductions in human exposures to ionising radiation resulting from radioactive discharges, as a result of planned reductions in discharges

To support implementation of UK Government policy, the Scottish Government has issued Statutory Guidance to SEPA (Scottish Government, 2008). Similarly, BEIS and the Welsh Government issued guidance to the Environment Agency (DECC and Welsh Assembly Government, 2009). The Environment Agency has developed Radioactive Substances Regulation (RSR) Environmental Principles (RSR Environmental Principles, or REPs) to form a consistent and standardised framework for the technical assessments that will be made when regulating radioactive substances (Environment Agency, 2008a). It has also issued guidance for assessment of BAT (Environment Agency, 2008b).

Information on work in progress within the OSPAR Convention can be found on OSPAR's website [www.ospar.org](http://www.ospar.org). The basis for OSPAR's approach is the RSS whose primary objective is to prevent marine pollution (OSPAR, 2003), as amended in 2010 (OSPAR, 2010a). A recent report from the OSPAR Radioactive Substances Committee records work completed and planned relating to reporting of discharges, environmental measurements, standards and quality assurance (OSPAR, 2016a). An agreement has been reached on the basis for monitoring of relevance to OSPAR by Contracting Parties (OSPAR, 2006). The programme includes sampling in fifteen divisions of the OSPAR maritime area and is supported by procedures for ensuring quality control. Inputs in the North-East Atlantic have been summarised

for both nuclear and non-nuclear sectors (OSPAR, 2011a; b). The UK submission concerning the implementation of the principle of using BAT has also been published (OSPAR, 2013). Progress by Contracting Parties towards meeting the objectives in RSS has been reviewed (OSPAR, 2009), as has the quality status of the Convention area (OSPAR, 2010b). The Quality Status Report considers radioactivity in food and the environment and refers to results of the monitoring programmes published in earlier issues of this report. The overall conclusions of the review were that there is evidence of:

- A reduction in total beta discharges from the nuclear sector, including technetium-99 discharges
- Reductions in marine concentrations of radioactive substances in most cases
- Estimated doses to humans were well within international and EU limits and
- An indication that the calculated dose rate to marine biota from selected radionuclides from the nuclear sector are low and are below the lowest levels at which any effects are likely to occur

The EC has considered various options for a new policy instrument concerning the protection and conservation of the marine environment and has now issued a Marine Strategy Framework Directive (CEC, 2008). The Directive has been transposed into UK law (United Kingdom - Parliament, 2010b) and is supported by measures to improve management of the marine environment covering the UK, and latterly Scotland and Northern Ireland (United Kingdom - Parliament, 2009; Scotland - Parliament, 2010; Department of the Environment Northern Ireland, 2010). It requires Member States to achieve Good Environmental Status in waters under their jurisdiction by 2020. The UK has submitted an initial assessment to the Commission (EC, 2012).

The importance of an integrated approach to stewardship of the marine environment has been recognised in the UK, and a strategy to achieve this has been published (Defra, Scottish Executive and Welsh Assembly Government, 2002). The report *"Safeguarding Our Seas"* considers conservation and sustainable development of the marine environment and sets out how the UK is addressing those issues in relation to radioactive and other substances and effects. The UK completed a fully integrated assessment of the marine environment in 2005 (Defra, 2005a, b; Defra, Department of the Environment Northern Ireland, Scottish Executive, Welsh Assembly Government, 2005) and has completed a new assessment *"Charting Progress 2"* in 2010 (Defra, 2010). The Department of the Environment Northern Ireland and the Scottish Government have also published individual assessments of the state of the seas around their coasts (Baxter *et al.*, 2011; Department of the Environment Northern Ireland, 2011).

In the 2008 White Paper *"Meeting the Energy Challenge"*, (Department for Business, Enterprise and Regulatory Reform, 2008), the UK Government set out its view that

new nuclear power stations should have a role to play in this country's future energy mix alongside other low-carbon sources; that it would be in the public interest to allow energy companies the option of investing in new nuclear power stations and that the Government should take active steps to facilitate this. More information concerning subsequent national policy statements, consultations and decisions, together with details of the approach for assessing the design of potential new nuclear power stations and approvals for their proposed developments, is available in earlier RIFE reports (e.g. Environment Agency, FSA, NIEA, NRW and SEPA, 2014). The eight nuclear sites, assessed as being potentially suitable for the development of new nuclear power stations, are shown in Figure 1.5.

The assessment process, for the design of potential new nuclear power stations, called "Generic Design Assessment" (GDA) process is continuing in 2016 with consideration of the Hitachi-GE UK ABWR reactor design, as well as progress towards closure of the AP1000 GDA (Environment Agency and ONR, 2016). ONR continues to be engaged in conducting safety and security assessment for Hinkley Point C (ONR, 2016) and is working with companies seeking to apply for nuclear site licenses at:

- Wylfa, Anglesey (Horizon Nuclear Power Limited, ABWR design)
- Moorside, Cumbria (NuGen Limited, AP1000 design)
- Sizewell C, Suffolk (NNB GenCo Limited, EPR design)

### 1.3.3 Managing radioactive liabilities in the UK

The UK Government has ratified the Joint Convention on the Safety of Spent Fuel Management and on the Safety of Radioactive Waste Management (IAEA, 1997). This agreement has an objective to ensure that individuals, society and the environment are protected from the harmful effects of ionising radiation as a result of the management of spent nuclear fuel and radioactive waste. The UK is required, on a triennial basis, to submit National Reports for International peer review, to comply with their obligations with the Joint Convention (for example, DECC, 2010; 2011, 2014a).

The current arrangements for managing civil sector nuclear clean-up are founded in the Energy Act 2004, which led to the establishment of the Nuclear Decommissioning Authority (NDA) in April 2005. NDA is responsible for nuclear sites formerly owned by British Nuclear Fuels Limited (BNFL), including ownership of its assets and liabilities, and United Kingdom Atomic Energy Authority (UKAEA). It is responsible for developing and implementing an overall strategy for cleaning up the civil public sector nuclear legacy safely, securely, and in ways that protect the environment. The current strategy was published in 2016 (NDA, 2016a) and the plan for 2016/19 is available (NDA, 2016b). The health and socio-economic impacts of the strategy have been considered (NDA,

**Figure 1.5.** Potential sites for new nuclear power stations



2016c). NDA published an up-to-date inventory and forecast of radioactive wastes in the UK jointly with BEIS in 2014 (NDA and DECC, 2014).

In 2007, the UK Government and Devolved Administrations issued a UK-wide policy for managing low level waste (Defra, 2007), which includes:

- Maintaining a focus on safety whilst allowing greater flexibility in managing LLW
- An emphasis on community involvement
- NDA creating a UK-wide strategy for managing LLW from the nuclear industry, including considering whether a replacement(s) of the national disposal facility near Drigg in Cumbria might be needed
- Initiating a UK-wide strategy for managing LLW from non-nuclear industries
- Minimising waste

In 2010, The NDA developed and published the “UK Strategy for the Management of Solid Low-Level Radioactive Waste from the Nuclear Industry”, known as the “Nuclear LLW Strategy”. A review of this took place in 2014, culminating in a public consultation on a revised version in January 2015 (DECC, Scottish Government, Welsh Government and Northern Ireland Department of Environment, 2015). The new strategy has now been published (DECC, Scottish Government, Welsh Government and Northern Ireland Department of Environment, 2016). The overall direction of the strategy remains unchanged. However, there have been significant changes including:

- The diversion of significant volumes of LLW from the Low Level Waste Repository (LLWR)
- The development and use of alternate treatment and disposal routes

- The application of the waste hierarchy by waste producers when making waste management decisions
- The identification of opportunities for improvement and the sharing of good practices for LLW management
- The engagement of a broad range of stakeholders within the process
- These changes are reflected in the new strategy.

More generally, consideration of the development of strategy for LLW from the non-nuclear industry resulted in the development and publication of the strategy in two parts:

Part 1 – Arisings of wastes mainly containing anthropogenic radionuclides (DECC, Scottish Government, Welsh Government and the Northern Ireland Department of the Environment, 2012), and most recently,

Part 2 – Arisings of wastes containing Naturally Occurring Radioactive Materials (NORM) in 2014 (DECC, Scottish Government, Welsh Government and the Northern Ireland Department of the Environment, 2014)

UK Government policy is that geological disposal is the best available means of managing higher activity radioactive waste in the long term.

The 2008 White Paper '*Managing Radioactive Waste Safely (MRWS): A Framework for Implementing Geological Disposal*' set out a framework for implementing geological disposal, including a voluntaristic process for identifying a Geological Disposal Facility (GDF) site that was based on local communities' willingness to participate in the process (Defra, Department for Business, Enterprise and Regulatory Reform, Welsh Assembly Government and Northern Ireland Assembly, 2008).

In July 2014, UK Government published a White Paper 'Implementing Geological Disposal' that sets out the policy framework for managing higher activity radioactive waste in the long term through geological disposal (DECC, 2014b). It set out a number of initial actions to implement this policy framework; actions which will be led by BEIS and the developer for a GDF (Radioactive Waste Management Limited, a wholly owned subsidiary company of NDA). Three initial actions, National Geological Screening (NGS); Working with Communities (WWC) and Land-use Planning, underpin the work to deliver outputs prior to formal engagement in 2017 with potential host communities. Consultation on the NGS exercise to bring together information about UK geology that is relevant to the long-term safety of a GDF is underway (Radioactive Waste Management, 2015). No specific sites have been selected or are currently under consideration (DECC, 2016a). A scoping report has considered the proposed content of an Appraisal of Sustainability for a GDF policy statement that is being prepared by BEIS (DECC, 2016b).

Radioactive waste management is a devolved policy issue. Therefore, the Scottish Government, Welsh Government

and Northern Ireland Executive each have responsibility for this issue in respect of their areas.

The Scottish Government is not a sponsor of the programme for implementing geological disposal, but does remain committed to dealing responsibly with radioactive waste arising in Scotland. Scottish Government policy is that the long-term management of higher activity radioactive waste should be in near-surface facilities. Facilities should be located as near to the site as possible (Scottish Government, 2011).

The Welsh Government is committed to securing the long-term safety of radioactive wastes and to the implementation of a framework appropriate to the needs of Wales and continues to play an active part in the MRWS programme to promote the interests of the people of Wales. In 2015, the Welsh Government adopted a policy for geological disposal for the long term management of higher activity radioactive waste (Welsh Government, 2015). The Welsh Government supports the policy of voluntary engagement where potential host communities are able to seek discussions, without prior commitment, about potentially hosting a GDF. Following a further consultation in 2015, the Welsh Government issued a second policy statement setting out its intentions for engaging with potential volunteer host communities (Welsh Government, 2015):

<http://gov.wales/docs/desh/policy/150519-policy-on-the-management-and-disposal-of-higher-activity-radioactive-waste-en.pdf>

The Northern Ireland Executive continues to support the implementation of geological disposal for the UK's higher activity radioactive waste, recognising that it is in the best interests of Northern Ireland that these wastes are managed in the safest and most secure manner.

Independent scrutiny of the Government's long-term management, storage and disposal of radioactive waste is continuing by the Committee on Radioactive Waste Management (CoRWM) who have published their proposed work programme for 2014-2017 (CoRWM, 2014).

Some low level radioactive waste, mostly from non-nuclear sites, and some very low level radioactive waste is currently disposed of in landfill by controlled burial (Section 7). There is still a large amount of solid low level radioactive wastes that will require disposal. Some will be sent to the LLWR near Drigg, the low level radioactive waste from Dounreay can be disposed of at the new Dounreay LLW Facility close to the site, and further alternative disposal options are also being considered. Guidance on requirements for authorisation for geological and near-surface disposal facilities has now been issued (Environment Agency and NIEA (2009), Environment Agency, NIEA and SEPA (2009) and Environment Agency (2013a)). In addition, SEPA has issued a policy statement

which specifies how it will regulate the disposal of LLW from nuclear licensed sites. The position identified has several practical implications including simplification of the process such that individual disposal sites need no longer be named in authorisations (SEPA, 2012b).

Decommissioning of many nuclear sites in Great Britain is underway. The environment agencies have begun a consultation process on draft guidance on the principles, requirements and regulatory process that will apply to nuclear sites in all stages of decommissioning and clean-up (NRW, SEPA and the Environment Agency, 2016). The aim is to ensure that radioactive waste and contamination is managed in a way that is safe so that nuclear sites may eventually be released from regulation under radioactive substances legislation.

NORM is contained in some wastes and is subject to existing regulatory systems which are designed to protect human health and the environment. However, there are improvements that can be achieved and, following a broad ranging consultation, BEIS, the Scottish and Welsh Governments and the Department of the Environment Northern Ireland published the UK NORM Waste Strategy in July 2014 (DECC, the Scottish Government, the Welsh Government and the Department of the Environment Northern Ireland, 2014). The Strategy in respect of the NORM sector is based on stimulating investment in the waste management supply chain. It will achieve this principally through (i) reforming the regulatory framework to ensure it is clear, coherent and effective, (ii) removing policy barriers to the development of a robust and efficient market for NORM waste management and (iii) supporting efforts by waste producers and the waste management supply chain to generate better data and information about current and future NORM waste arisings.

### 1.3.4 Solid radioactive waste disposal at sea

In the past, packaged solid waste of low specific activity was disposed of deep in the North Atlantic Ocean. The last disposal of this type was in 1982. The UK Government announced at the OSPAR Ministerial meeting in 1998 that it was stopping disposal of this material at sea. At that meeting, Contracting Parties agreed that there would no longer be any exception to prohibiting the dumping of radioactive substances, including waste (OSPAR, 1998). The environmental impact of the deep ocean disposals was predicted by detailed mathematical modelling and has been shown to be negligible (OECD, Nuclear Energy Agency, 1985). Disposals of small amounts of waste also took place from 1950 to 1963 in a part of the English Channel known as the Hurd Deep. The results of environmental monitoring of this area are presented in Section 8 and confirm that the radiological impact of these disposals was insignificant.

In England, MMO administers a range of statutory controls that apply to marine works on behalf of the Secretary of State for Environment, Food and Rural Affairs; this includes issuing licences under the MCAA 2009 (United Kingdom - Parliament, 2009) for the disposal of dredged material at sea. Licences for disposals made in Scottish waters and around the coast of Northern Ireland are the responsibility of the Scottish Government (Marine Scotland) and the Department of Agriculture Environment and Rural Affairs, respectively. As of 1 April 2010, licences for Welsh waters are the responsibility of the Welsh Government.

The protection of the marine environment is considered before a licence is issued. Since dredge materials will contain varying concentrations of radioactivity from natural and artificial sources, assessments are carried out, when appropriate, to provide reassurance that there is no significant risk to the food chain or other risk from the disposal. Guidance on exemption criteria for radioactivity in relation to sea disposal is available from IAEA (1999). IAEA has published a system of assessment that can be applied to dredge spoil disposal (IAEA, 2003) and which has been adapted to reflect operational practices in England and Wales (McCubbin and Vivian, 2006). In 2015, one request was received by MMO to apply for a licence for the disposal of dredged material at sea from Maryport Harbour. The results of the assessment are given in Appendix 5.

### 1.3.5 Other sources of radioactivity

There are several other man-made sources of radioactivity that may affect the food chain and the environment. These could include disposals of material from offshore installations, transport incidents, satellite re-entry, releases from overseas nuclear installations and the operation of nuclear powered submarines. PHE has assessed incidents involving the transport of radioactive materials in the UK (Jones and Harvey, 2014). PHE have also considered the effects of discharges from the oil and gas industry into the marine environment (Harvey *et al.*, 2010). Using modelling, the highest individual (per head of population) annual doses for discharges from 2005-2008 were estimated to be less than 0.001 mSv. Submarine berths in the UK are monitored by the MoD (DSTL Radiological Protection Services, 2013). General monitoring of the British Isles is carried out as part of the programmes described in this report, to detect any gross effects from the sources above. No such effects were found in 2015. Low concentrations of radionuclides were detected in the marine environment around the Channel Islands (Section 8) and these may be partly due to discharges from the nuclear fuel reprocessing plant at La Hague in France.

The exploration for, and extraction of, gas from shale rock is being actively investigated in the UK with support from BEIS. This process, along with others for unconventional sources of gas such as coal bed methane, represents a potential source of exposure of the public and workers

to naturally occurring radioactivity. The form of the radioactivity could be gaseous, liquid or solid. Examples of routes of exposure are inhalation of radon gas emissions, and ingestion of water and food where the process has enhanced levels of NORM.

Each of the environment agencies is working to ensure that appropriate regulatory regimes are in operation to control exposures of the public from unconventional gas exploration and extraction. Reports have been published to support engagement with industry, the public and other stakeholders (Environment Agency, 2013b; NIEA, 2013; SEPA, 2013). The Environment Agency has now granted permits for Cuadrilla to carry out shale gas exploration at their site in Roseacre Wood near Elswick in Lancashire and Preston New Road, near Blackpool (Environment Agency and DECC, 2015). A review of potential public health impacts of exposures to radioactivity as a result of shale gas extraction has been issued by PHE (Kibble *et al.*, 2014). Monitoring of radioactivity from the exploration and extraction of shale gas in the environment and food is not undertaken by the environment agencies, FSA or FSS at present. However, the agencies will continue to review the position as specific proposals for development are taken forward and any results of monitoring will be reported in future issues of the RIFE report.

The Environmental Protection Act 1990 provides the basis, through the Environment Act 1995, for a regulatory regime for identifying and remediating contaminated land. The regime was extended in 2006 to provide a system for identifying and remediating land, where contamination is causing people to be exposed to lasting exposure to radiation resulting from the after-effects of a radiological emergency, past practice or post work activity; and where intervention is liable to be justified. A further modification was made in 2007, which extends the regime to cover land contaminated with radioactivity originating from nuclear installations, although to date no sites meeting these criteria have been found. A profile of industries which may have caused land contamination has been published (Defra, 2006). Dose criteria for the designation of contaminated land have been determined for England and Wales (Smith *et al.*, 2006). A report giving an overview of the progress made by local authorities and the Environment Agency in identifying and remediating contaminated land was published in 2009 (Environment Agency, 2009c). BEIS issued revised guidance for radioactive contaminated land to local authorities and the Environment Agency in 2012 (DECC, 2012). The Environment Agency has issued a series of Briefing Notes that provide information on land contaminated with radioactivity in England and Wales (Environment Agency, 2012). To date, no site has been legally designated as 'contaminated land' due to radioactivity in England and Wales.

Equivalent legislation for identifying and remediating contaminated land comprising The Radioactive

Contaminated Land Regulations (Northern Ireland) 2006 and subsequent amending legislation, issued in 2007 and 2010, exists as Statutory Instruments in Northern Ireland (Statutory Instruments, 2007; 2010).

In October 2007, the Radioactive Contaminated Land (Scotland) Regulations came into force by amending Part II A of the Environmental Protection Act 1990. SEPA has powers to inspect land that may be contaminated with radioactivity, to decide if land should be identified as radioactive contaminated land and require remediation if considered necessary. Revised Statutory Guidance was issued to SEPA in 2009. This guidance is broadly similar to that issued to the Environment Agency, apart from the fact that for the designation of radioactive contaminated land, clear dose criteria are set for homogeneous and heterogeneous contamination, and whether or not the probability of receiving the dose should be taken into account. To date, no site has been designated as 'contaminated land' due to radioactivity in Scotland.

The contribution of aerial radioactive discharges from UK installations to concentrations of radionuclides in the marine environment has been studied (Defra, 2004). Tritium and carbon-14 were predicted to be at concentrations that were particularly high in relation to actual measured values in the Irish Sea. However, the study suggested that this was due to unrealistic assumptions being made in the assessment. The main conclusion was that aerial discharges do not make a significant contribution to levels in the marine environment. On occasion, the effects of aerial discharges are detected in the aquatic environment, and conversely the effects of aquatic discharges are detected on land. Where this is found, appropriate comments are made in this report.

All sources of ionising radiation exposure to the UK population are reviewed, the most recent report being published in 2005 (Watson *et al.*, 2005). Sources of naturally occurring radiation and man-made radiation produced for medical use predominate. The average annual dose from naturally occurring radiation was found to be 2.2 mSv and about half of this was from radon exposure indoors. The average annual dose from artificial radiation was 0.42 mSv, mainly derived from medical procedures, such as x-rays. The overall average annual dose was 2.7 mSv. Exposures from non-medical man-made sources were very low and discharges of radioactive wastes contributed less than 0.1 per cent of the total. These figures represent the exposure of the average person.

The RIFE report is directed at establishing the exposure of people who might receive the highest possible doses due to radioactive waste discharges as a result of their age, diet, location or habits. It is the exposure of these people which forms the basis for comparisons with dose limits in EU and UK law.

**Table 1.1. Individual doses – direct radiation pathway, 2015**

Site	Exposure, mSv
<b>Nuclear fuel production and reprocessing</b>	
Capenhurst	0.13
Sellafield	0.005
Springfields	0.020
<b>Research establishments</b>	
Dounreay	0.003
Harwell	0.017
Winfrith	0.014
<b>Nuclear power stations</b>	
Berkeley	Bgd <sup>a</sup>
Bradwell	0.017
Chapelcross	Bgd <sup>a</sup>
Dungeness	0.013 <sup>b</sup>
Hartlepool	0.010
Heysham	0.010
Hinkley Point	0.010 <sup>c</sup>
Hunterston	0.020 <sup>d</sup>
Oldbury	0.001
Sizewell	0.020 <sup>e</sup>
Torness	0.020
Trawsfynydd	0.006
Wylfa	0.012
<b>Defence establishments</b>	
Aldermaston	Bgd <sup>a</sup>
Barrow	Bgd <sup>a</sup>
Burghfield	Bgd <sup>a</sup>
Derby	Bgd <sup>a</sup>
Devonport	Bgd <sup>a</sup>
Faslane	Bgd <sup>a</sup>
Rosyth	Bgd <sup>a</sup>
<b>Radiochemical production</b>	
Amersham	0.14
Cardiff	Bgd <sup>a</sup>
<b>Industrial and landfill sites</b>	
LLWR near Drigg	0.039

<sup>a</sup> Doses not significantly different from natural background

<sup>b</sup> Datum for Dungeness A. Dungeness B (0.010) not used

<sup>c</sup> Datum for Hinkley B. Hinkley A (Bgd<sup>a</sup>) not used

<sup>d</sup> Datum for Hunterston B. Hunterston A (0.003) not used

<sup>e</sup> Datum for Sizewell B. Sizewell A (Bgd<sup>a</sup>) not used

**Table 1.2. Total doses integrated across pathways, 2015**

Site	Representative person <sup>a</sup>	Exposure, mSv	
		Total	Dominant contributions <sup>b</sup>
<b>A Gaseous releases and direct radiation from the site</b>			
Aldermaston and Burghfield	Infant milk consumers	<0.005 <sup>g</sup>	Milk, <sup>3</sup> H <sup>d</sup> , <sup>137</sup> Cs <sup>c</sup> , <sup>234</sup> U, <sup>238</sup> U
Amersham	Local adult inhabitants (0 - 0.25km)	0.14 <sup>g</sup>	Direct radiation
Barrow	Adult potato consumers	<0.005	Gamma dose rate over sediment, potatoes, <sup>137</sup> Cs <sup>c</sup>
Berkeley and Oldbury	Infant milk consumers	<0.005	Milk, <sup>14</sup> C, <sup>35</sup> S <sup>c</sup>
Bradwell	Prenatal children of local inhabitants (0 - 0.25km)	0.017	Direct radiation
Capenhurst	Local adult inhabitants (0 - 0.25km)	0.13 <sup>g</sup>	Direct radiation
Cardiff	Infant milk consumers	<0.005	Milk, <sup>14</sup> C, <sup>32</sup> P <sup>c</sup> , <sup>35</sup> S
Chapelcross	Infant milk consumers	0.022	Milk, <sup>90</sup> Sr, <sup>241</sup> Am <sup>c</sup>
Derby	Adult potato consumers	<0.005 <sup>g</sup>	Potatoes, <sup>137</sup> Cs, <sup>234</sup> U, <sup>238</sup> U
Devonport	Prenatal children of root vegetable consumers	<0.005	Fish, root vegetables, <sup>3</sup> H <sup>c</sup> , <sup>14</sup> C, <sup>131</sup> I <sup>c</sup>
Dounreay	Adult green vegetable consumers	0.010	<sup>90</sup> Sr, <sup>129</sup> I <sup>d</sup> , <sup>238</sup> Pu <sup>d</sup> , <sup>239/240</sup> Pu, <sup>241</sup> Am <sup>c</sup>
Dungeness	Local adult inhabitants (0.5 - 1km)	0.014	Direct radiation
Faslane	Adult consumers of cattle meat	<0.005	Cattle meat, <sup>137</sup> Cs, <sup>241</sup> Am <sup>c</sup>
Hartlepool	Local adult inhabitants (0 - 0.25km)	0.022	Direct radiation, gamma dose rate over sediment
Harwell	Local adult inhabitants (0 - 0.25km)	0.017	Direct radiation
Heysham	Local adult inhabitants (0 - 0.25km)	0.011	Direct radiation, external and inhalation near site
Hinkley Point	Local adult inhabitants (0.5 - 1km)	0.011	Direct radiation
Hunterston	Infant milk consumers	0.025	Milk, <sup>90</sup> Sr, <sup>241</sup> Am <sup>c</sup>
LLWR near Drigg	Local child inhabitants (0.5 - 1km)	0.041	Direct radiation
Rosyth <sup>d</sup>	-	-	-
Sellafield	Local child inhabitants (0 - 0.25km)	0.008 <sup>g</sup>	Direct radiation, potatoes, <sup>106</sup> Ru <sup>c</sup>
Sizewell	Local adult inhabitants (0 - 0.25km)	0.021	Direct radiation
Springfields	Local infant inhabitants (0.5 - 1km)	0.020 <sup>g</sup>	Direct radiation
Torness	Local adult inhabitants (0.5 - 1km)	0.020	Direct radiation
Trawsfynydd	Local infant inhabitants (0.25 - 0.5km)	0.014	Direct radiation, milk, <sup>241</sup> Am
Winfrith	Local adult inhabitants (0.25-0.5km)	0.014	Direct radiation
Wylfa	Local infant inhabitants (0.25 - 0.5km)	0.013	Direct radiation
<b>B Liquid releases from the site</b>			
Aldermaston and Burghfield	Adult occupants over riverbank	<0.005	Exposure over riverbank
Amersham	Adult occupants over riverbank	0.006	Gamma dose rate over riverbank
Barrow	Adult occupants on houseboats	0.051	Gamma dose rate over sediment
Berkeley and Oldbury	Adult occupants over sediment	<0.005	Gamma dose rate over sediment
Bradwell	Adult occupants on houseboats	<0.005	Gamma dose rate over sediment
Capenhurst	Occupants over riverbank aged 10y	0.007	Gamma dose rate over sediment
Cardiff	Prenatal children of occupants over sediment	<0.005	Gamma dose rate over sediment
Chapelcross	Adult wildfowl consumers	0.006	Gamma dose rate over sediment, fish, wildfowl, <sup>239/240</sup> Pu, <sup>241</sup> Am
Derby	Adult consumers of locally sourced water	<0.005	Water, <sup>60</sup> Co <sup>c</sup>
Devonport	Adult fish consumers	<0.005	Fish, <sup>14</sup> C, <sup>131</sup> I <sup>c</sup> , <sup>241</sup> Am <sup>c</sup>
Dounreay	Adult occupants over sediment	0.010	Gamma dose rate over sediment
Dungeness	Prenatal children of occupants over sediment	<0.005	Direct radiation, gamma dose rate over sediment
Faslane	Adult occupants over sediment	<0.005	Gamma dose rate over sediment
Hartlepool	Adult occupants over sediment	0.022	Direct radiation, gamma dose rate over sediment
Harwell	Prenatal children of occupants over sediment	<0.005	Gamma dose rate over riverbank
Heysham	Adult mollusc consumers	0.023	Fish, gamma dose rate over sediment, molluscs, <sup>239/240</sup> Pu, <sup>241</sup> Am
Hinkley Point	Adult occupants over sediment	0.016	Gamma dose rate over sediment
Hunterston	Adult occupants over sediment	<0.005	Gamma dose rate over sediment
LLWR near Drigg <sup>e</sup>	Adult mollusc consumers	0.42 <sup>f</sup>	Crustaceans, molluscs, <sup>210</sup> Po
Rosyth	Adult occupants over sediment	0.006	Gamma dose rate over sediment
Sellafield <sup>e</sup>	Adult mollusc consumers	0.42 <sup>f</sup>	Crustaceans, molluscs, <sup>210</sup> Po
Sizewell	Adult occupants on houseboats	<0.005	Gamma dose rate over sediment
Springfields	Adult occupants on houseboats	0.050	Gamma dose rate over sediment
Torness	Adult fish consumers	0.005	Direct radiation, fish, <sup>241</sup> Am
Trawsfynydd	Adult fish consumers	0.008	Exposure over sediment, fish, <sup>90</sup> Sr, <sup>137</sup> Cs, <sup>241</sup> Am
Whitehaven <sup>e</sup>	Adult mollusc consumers	0.42 <sup>f</sup>	Crustaceans, molluscs, <sup>210</sup> Po
Winfrith	Adult occupants over sediment	0.006	Gamma dose rate over sediment
Wylfa	Adult occupants over sediment	0.005	Gamma dose rate over sediment

**Table 1.2. continued**

Site	Representative person <sup>a</sup>	Exposure, mSv	
		Total	Dominant contributions <sup>b</sup>
<b>C All sources</b>			
Aldermaston and Burghfield	Infant milk consumers	<0.005 <sup>g</sup>	Milk, <sup>3</sup> H <sup>d</sup> , <sup>137</sup> Cs <sup>c</sup> , <sup>234</sup> U, <sup>238</sup> U
Amersham	Local adult inhabitants (0 - 0.25km)	0.14 <sup>g</sup>	Direct radiation
Barrow	Adult occupants on houseboats	0.051	Gamma dose rate over sediment
Berkeley and Oldbury	Infant milk consumers	<0.005	Milk, <sup>14</sup> C, <sup>35</sup> S <sup>c</sup>
Bradwell	Prenatal children of local inhabitants (0 - 0.25km)	0.017	Direct radiation
Capenhurst	Local adult inhabitants (0 - 0.25km)	0.13 <sup>g</sup>	Direct radiation
Cardiff	Prenatal children of occupants over sediment	<0.005	Gamma dose rate over sediment
Chapelcross	Infant milk consumers	0.022	Milk, <sup>90</sup> Sr, <sup>241</sup> Am <sup>c</sup>
Derby	Adult consumers of locally sourced water	<0.005	Water, <sup>60</sup> Co <sup>c</sup>
Devonport	Adult fish consumers	<0.005	Fish, <sup>14</sup> C, <sup>131</sup> I <sup>c</sup> , <sup>241</sup> Am <sup>c</sup>
Dounreay	Adult occupants over sediment	0.010	Gamma dose rate over sediment
Dungeness	Local adult inhabitants (0.5 - 1km)	0.014	Direct radiation
Faslane	Adult consumers of cattle meat	<0.005	Cattle meat, <sup>137</sup> Cs, <sup>241</sup> Am <sup>c</sup>
Hartlepool	Local adult inhabitants (0 - 0.25km)	0.022	Direct radiation, gamma dose rate over sediment
Harwell	Local adult inhabitants (0 - 0.25km)	0.017	Direct radiation
Heysham	Adult mollusc consumers	0.023	Fish, gamma dose rate over sediment, molluscs, <sup>239/240</sup> Pu, <sup>241</sup> Am
Hinkley Point	Adult occupants over sediment	0.016	Gamma dose rate over sediment
Hunterston	Infant milk consumers	0.025	Milk, <sup>90</sup> Sr, <sup>241</sup> Am <sup>c</sup>
LLWR near Drigg <sup>e</sup>	Adult mollusc consumers	0.42 <sup>f</sup>	Crustaceans, molluscs, <sup>210</sup> Po
Rosyth	Adult occupants over sediment	0.006	Gamma dose rate over sediment
Sellafield <sup>e</sup>	Adult mollusc consumers	0.42 <sup>f</sup>	Crustaceans, molluscs, <sup>210</sup> Po
Sizewell	Local adult inhabitants (0 - 0.25km)	0.021	Direct radiation
Springfields	Adult occupants on houseboats	0.050	Gamma dose rate over sediment
Torness	Local adult inhabitants (0.5 - 1km)	0.020	Direct radiation
Trawsfynydd	Local infant inhabitants (0.25 - 0.5km)	0.014	Direct radiation, milk, <sup>241</sup> Am
Whitehaven <sup>e</sup>	Adult mollusc consumers	0.42 <sup>f</sup>	Crustaceans, molluscs, <sup>210</sup> Po
Winfrith	Prenatal children of local inhabitants (0.5 - 1km)	0.014	Direct radiation
Wylfa	Local infant inhabitants (0.25 - 0.5km)	0.013	Direct radiation

<sup>a</sup> Selected on the basis of providing the highest dose from the pathways associated with the sources as defined in A, B or C

<sup>b</sup> Pathways and radionuclides that contribute more than 10% of the total dose. Some radionuclides are reported as being at the limits of detection and based on these measurements, an upper estimate of dose is calculated

<sup>c</sup> The assessed contribution is based on data at limits of detection

<sup>d</sup> The effects of gaseous discharges and direct radiation are not assessed as there are no sources for this site

<sup>e</sup> The effects of liquid discharges from Sellafield, Whitehaven and LLWR near Drigg are considered together when assessing exposures at these sites because their effects are manifested in a common area of the Cumbrian coast

<sup>f</sup> The doses from man-made and naturally occurring radionuclides were 0.078 and 0.35 mSv respectively. The source of naturally occurring radionuclides was a phosphate processing works near Sellafield at Whitehaven. Minor discharges of radionuclides were also made from the LLWR near Drigg into the same area

<sup>g</sup> Includes a component due to natural sources of radionuclides

**Table 1.3. Trends in total doses (mSv) from all sources<sup>a</sup>**

Site	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Aldermaston and Burghfield	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005
Amersham		0.24	0.24	0.22	0.23	0.22	0.22	0.22	0.22	0.22	0.22	0.14	0.14
Barrow										0.057	0.076	0.055	0.051
Berkeley and Oldbury		<i>0.12</i>	<i>0.090</i>	<i>0.042</i>	0.061	0.041	0.058	0.011	0.006	0.014	0.010	<0.005	<0.005
Bradwell		<i>0.090</i>	<i>0.067</i>	<i>0.075</i>	0.070	0.070	0.098	0.13	0.048	<0.005	<0.005	<0.005	0.017
Capenhurst		<i>0.080</i>	<i>0.080</i>	<i>0.085</i>	0.12	0.17	0.19	0.26	0.095	0.085	0.080	0.17	0.13
Cardiff	0.038	0.023	0.023	0.011	0.008	0.007	0.006	0.006	0.006	0.005	0.010	<0.005	<0.005
Chapelcross		<i>0.022</i>	<i>0.023</i>	<i>0.024</i>	<i>0.019</i>	<i>0.021</i>	<i>0.017</i>	<i>0.029</i>	<i>0.037</i>	<i>0.011</i>	<i>0.024</i>	<i>0.014</i>	<i>0.022</i>
Derby							<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005
Devonport		<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005
Dounreay	0.012	0.011	0.043	0.029	0.059	0.078	0.063	0.047	0.018	0.017	0.012	0.012	0.010
Dungeness		<i>0.48</i>	<i>0.55</i>	<i>0.63</i>	<i>0.28</i>	<i>0.40</i>	<i>0.32</i>	<i>0.022</i>	<i>0.021</i>	<i>0.015</i>	<i>0.021</i>	<i>0.021</i>	<i>0.014</i>
Faslane		<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005
Hartlepool	0.021	0.020	0.021	0.021	0.021	0.026	0.027	0.025	0.025	0.015	0.024	0.027	0.022
Harwell		<i>0.017</i>	<i>0.022</i>	<i>0.026</i>	<i>0.022</i>	<i>0.020</i>	<i>0.023</i>	<i>0.018</i>	<i>0.017</i>	<i>0.018</i>	<i>0.010</i>	<i>0.016</i>	<i>0.017</i>
Heysham		<i>0.036</i>	<i>0.028</i>	<i>0.037</i>	<i>0.038</i>	<i>0.046</i>	<i>0.049</i>	<i>0.057</i>	<i>0.025</i>	<i>0.025</i>	<i>0.028</i>	<i>0.023</i>	<i>0.023</i>
Hinkley Point		<i>0.026</i>	<i>0.027</i>	<i>0.048</i>	<i>0.035</i>	<i>0.045</i>	<i>0.055</i>	<i>0.014</i>	<i>0.014</i>	<i>0.013</i>	<i>0.022</i>	<i>0.022</i>	<i>0.016</i>
Hunterston		<i>0.10</i>	<i>0.090</i>	<i>0.074</i>	<i>0.090</i>	<i>0.077</i>	<i>0.067</i>	<i>0.067</i>	<i>0.050</i>	<i>0.032</i>	<i>0.021</i>	<i>0.021</i>	<i>0.025</i>
LLWR near Drigg <sup>b</sup>	0.66	0.58	0.40	0.43	0.37	0.47	0.28	0.18	0.18	0.30	0.061	0.22	0.42
Rosyth		<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	0.006
Sellafield <sup>b</sup>	0.66	0.58	0.40	0.43	0.37	0.47	0.28	0.18	0.18	0.30	0.076 <sup>c</sup>	0.22	0.42
Sizewell		<i>0.045</i>	<i>0.086</i>	<i>0.090</i>	<0.005	0.031	0.026	0.020	0.021	0.021	0.021	0.020	0.021
Springfields		<i>0.17</i>	<i>0.15</i>	0.13	0.11	0.16	0.15	0.17	0.13	0.068	0.060	0.050	0.050
Torness		<i>0.024</i>	<i>0.025</i>	0.024	0.022	0.022	0.022	0.025	0.020	0.020	0.020	0.020	0.020
Trawsfynydd		<i>0.032</i>	<i>0.021</i>	<i>0.028</i>	<i>0.018</i>	<i>0.031</i>	<i>0.018</i>	<i>0.028</i>	<i>0.012</i>	<i>0.025</i>	<i>0.017</i>	<i>0.013</i>	<i>0.014</i>
Whitehaven <sup>b</sup>	0.66	0.58	0.40	0.43	0.37	0.47	0.28	0.18	0.18	0.30	0.061	0.22	0.42
Winfrith	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	<0.005	0.014
Wylfa		0.011	0.010	0.011	0.011	0.011	0.011	0.007	0.008	0.006	<0.005	0.007	0.013

<sup>a</sup> Where no data is given, no assessment was undertaken due to a lack of suitable habits data at the time. Data in italics signify assessments performed to show trends in total dose over the five-year period from 2004–2008, using subsequently obtained habits data

<sup>b</sup> The effects of liquid discharges from Sellafield, Whitehaven and LLWR near Drigg are considered together when assessing exposures at these sites

<sup>c</sup> The highest exposure due to operations at Sellafield was to people living in houseboats near Barrow

**Table 1.4. Source specific doses due to discharges of radioactive waste in the United Kingdom, 2015\***

Establishment	Radiation exposure pathways	Gaseous or liquid source <sup>e</sup>	Exposure, mSv <sup>b</sup> per year	Contributors <sup>c</sup>
<b>Nuclear fuel production and processing</b>				
Capenhurst	Inadvertent ingestion of water and sediment and external <sup>h</sup>	L	0.010	Ext
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005 <sup>i</sup>	<sup>3</sup> H <sup>d</sup> , <sup>99</sup> Tc <sup>d</sup> , <sup>234</sup> U, <sup>238</sup> U
Springfields	Fish and shellfish consumption and external intertidal areas	L	0.020	Ext
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005 <sup>i</sup>	<sup>129</sup> I <sup>d</sup> , <sup>234</sup> Th <sup>d</sup> , <sup>234</sup> U, <sup>238</sup> U
	External in intertidal areas (children playing) <sup>a,h</sup>	L	<0.005	Ext
	Occupancy of houseboats	L	0.054	Ext
	External in intertidal areas (farmers)	L	0.034	Ext
	Wildfowl consumers	L	0.006	Ext
Sellafield <sup>f</sup>	Fish and shellfish consumption and external in intertidal areas (2011-2015 surveys) (excluding naturally occurring radionuclides) <sup>l</sup>	L	0.084	Ext, <sup>239/240</sup> Pu, <sup>241</sup> Am
	Fish and shellfish consumption and external in intertidal areas (2011-2015 surveys) (including naturally occurring radionuclides) <sup>m</sup>	L	0.37	<sup>210</sup> Po
	Fish and shellfish consumption and external in intertidal areas (2015 surveys) (excluding naturally occurring radionuclides) <sup>l</sup>	L	0.098	Ext, <sup>239/240</sup> Pu, <sup>241</sup> Am
	Terrestrial foods, external and inhalation near Sellafield <sup>i</sup>	G	0.012	<sup>14</sup> C, <sup>90</sup> Sr, <sup>106</sup> Ru <sup>d</sup>
	Terrestrial foods at Ravenglass <sup>j</sup>	G/L	0.017	<sup>106</sup> Ru <sup>d</sup> , <sup>144</sup> Ce <sup>d</sup>
	External in intertidal areas (Ravenglass) <sup>a</sup>	L	0.013	Ext, <sup>241</sup> Am
	Occupancy of houseboats (Ribble estuary)	L	0.054	Ext
	Occupancy of houseboats (Barrow)	L	0.050	Ext
	External (skin) to bait diggers	L	0.064 <sup>g</sup>	Beta
	Handling of fishing gear	L	0.10 <sup>g</sup>	Beta
<b>Research establishments</b>				
Culham	Water consumption <sup>o</sup>	L	<0.005	<sup>137</sup> Cs <sup>d</sup>
Dounreay	Fish and shellfish consumption and external in intertidal areas	L	0.011	Ext
	Terrestrial foods, external and inhalation near site	G	0.014	<sup>90</sup> Sr, <sup>129</sup> I <sup>d</sup> , <sup>238</sup> Pu <sup>d</sup> , <sup>239/240</sup> Pu, <sup>241</sup> Am <sup>d</sup>
Harwell	Fish consumption and external to anglers	L	<0.005	Ext
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005	<sup>222</sup> Rn, <sup>3</sup> H <sup>d</sup>
Winfrith	Fish and shellfish consumption and external in intertidal areas	L	<0.005	Ext, <sup>241</sup> Am
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005	<sup>14</sup> C, <sup>137</sup> Cs <sup>d</sup>
<b>Nuclear power production</b>				
Berkeley and Oldbury	Fish and shellfish consumption and external in intertidal areas	L	<0.005	Ext
	Occupancy of houseboats	L	0.006	Ext
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	0.005	<sup>14</sup> C
Bradwell	Fish and shellfish consumption and external in intertidal areas	L	<0.005	Ext, <sup>241</sup> Am
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005	<sup>14</sup> C
Chapelcross	Wildfowl, fish and mollusc consumption and external in intertidal areas	L	0.009	Ext, <sup>239/240</sup> Pu, <sup>241</sup> Am
	Crustacean consumption	L	<0.005	<sup>137</sup> Cs, <sup>241</sup> Am
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	0.017	<sup>14</sup> C, <sup>90</sup> Sr, <sup>241</sup> Am <sup>d</sup>
Dungeness	Fish and shellfish consumption and external in intertidal areas	L	0.006	Ext, <sup>241</sup> Am
	Occupancy of houseboats	L	<0.005	Ext
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	0.005	<sup>14</sup> C, <sup>60</sup> Co <sup>d</sup>
Hartlepool	Fish and shellfish consumption and external in intertidal areas	L	0.020	Ext
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005	<sup>14</sup> C, <sup>35</sup> S <sup>d</sup> , <sup>60</sup> Co <sup>d</sup>
Heysham	Fish and shellfish consumption and external in intertidal areas	L	0.031	Ext, <sup>241</sup> Am
	External in intertidal areas (turf cutters)	L	0.016	Ext
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005	<sup>14</sup> C, <sup>35</sup> S, <sup>137</sup> Cs
Hinkley Point	Fish and shellfish consumption and external in intertidal areas	L	0.023	Ext
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005	<sup>14</sup> C, <sup>35</sup> S, <sup>60</sup> Co <sup>d</sup>
Hunterston	Fish and shellfish consumption and external in intertidal areas	L	<0.005	Ext, <sup>241</sup> Am
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	0.019	<sup>14</sup> C, <sup>90</sup> Sr, <sup>241</sup> Am <sup>d</sup>
Sizewell	Fish and shellfish consumption and external in intertidal areas	L	<0.005	Ext, <sup>241</sup> Am
	Occupancy of houseboats	L	<0.005	Ext
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005	<sup>14</sup> C, <sup>35</sup> S <sup>d</sup>
Torness	Fish and shellfish consumption and external in intertidal areas	L	<0.005	<sup>241</sup> Am
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	0.006	<sup>14</sup> C, <sup>35</sup> S <sup>d</sup> , <sup>90</sup> Sr

**Table 1.4. continued**

Establishment	Radiation exposure pathways	Gaseous or liquid source <sup>e</sup>	Exposure, mSv <sup>b</sup> per year	Contributors <sup>c</sup>
Trawsfynydd	Fish consumption and external to anglers	L	0.007	Ext, <sup>137</sup> Cs, <sup>241</sup> Am
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	0.016	<sup>14</sup> C, <sup>241</sup> Am
Wylfa	Fish and shellfish consumption and external in intertidal areas	L	0.007	Ext, <sup>241</sup> Am
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005	<sup>14</sup> C, <sup>35</sup> S
<b>Defence establishments</b>				
Aldermaston	Fish consumption and external to anglers	L	<0.005 <sup>l</sup>	Ext, <sup>131</sup> I <sup>d</sup> , <sup>234</sup> U
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005 <sup>l</sup>	<sup>3</sup> H <sup>d</sup> , <sup>137</sup> Cs <sup>d</sup> , <sup>234</sup> U, <sup>238</sup> U
Barrow	Occupancy of houseboats	L	0.050	Ext
	Terrestrial food consumption	G	<0.005	<sup>137</sup> Cs <sup>d</sup>
Derby	Water consumption, fish consumption and external to anglers <sup>o</sup>	L	<0.005	<sup>60</sup> Co <sup>d</sup>
	Terrestrial foods, external and inhalation near site	G	<0.005	<sup>137</sup> Cs, <sup>234</sup> U, <sup>238</sup> U
Devonport	Fish and shellfish consumption and external in intertidal areas	L	<0.005	Ext, <sup>14</sup> C, <sup>131</sup> I <sup>d</sup> , <sup>241</sup> Am <sup>d</sup>
	Occupancy of houseboats	L	<0.005	Ext
	Terrestrial foods, external and inhalation near site <sup>p</sup>	G	<0.005	<sup>3</sup> H <sup>d</sup> , <sup>14</sup> C
Faslane	Fish and shellfish consumption and external in intertidal areas	L	<0.005	Ext, <sup>241</sup> Am
	Terrestrial food consumption	G	<0.005	<sup>137</sup> Cs, <sup>241</sup> Am <sup>d</sup>
Holy Loch	External in intertidal areas	L	0.012	Ext
Rosyth	Fish and shellfish consumption and external in intertidal areas	L	0.010	Ext, <sup>241</sup> Am
<b>Radiochemical production</b>				
Amersham	Fish consumption and external to anglers	L	0.006	Ext
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	0.010	<sup>222</sup> Rn
Cardiff	Fish and shellfish consumption and external in intertidal areas <sup>p</sup>	L	0.006	Ext
	Terrestrial foods, external and inhalation near site <sup>i</sup>	G	<0.005	<sup>14</sup> C, <sup>32</sup> P <sup>d</sup> , <sup>35</sup> S, <sup>125</sup> I <sup>d</sup>
	Inadvertent ingestion and riverbank occupancy (River Taff)	L	<0.005	Ext, <sup>14</sup> C <sup>d</sup> , <sup>125</sup> I <sup>d</sup> , <sup>137</sup> Cs
<b>Industrial and landfill</b>				
LLWR near Drigg	Terrestrial foods <sup>j</sup>	G	<0.005	<sup>90</sup> Sr <sup>d</sup> , <sup>106</sup> Ru <sup>d</sup> , <sup>137</sup> Cs, <sup>144</sup> Ce <sup>d</sup>
	Fish and shellfish consumption and external in intertidal areas (2011-2015 surveys) (including naturally occurring radionuclides) <sup>f,m</sup>	L	0.37	<sup>210</sup> Po
	Water consumption <sup>o</sup>	L	<0.005	<sup>90</sup> Sr, <sup>134</sup> Cs <sup>d</sup> , <sup>137</sup> Cs <sup>d</sup> , <sup>210</sup> Po <sup>d</sup>
Whitehaven	Fish and shellfish consumption and external in intertidal areas (2011-2015 surveys) (excluding artificial radionuclides) <sup>f,k</sup>	L	0.29	<sup>210</sup> Po
	Fish and shellfish consumption and external in intertidal areas (2011-2015 surveys) (including artificial radionuclides) <sup>f,n</sup>	L	0.37	<sup>210</sup> Po

\* Source specific dose assessments are performed to provide additional information and as a check on the total dose assessment method

<sup>a</sup> Includes a component due to inadvertent ingestion of water or sediment or inhalation of resuspended sediment where appropriate

<sup>b</sup> Unless otherwise stated represents committed effective dose calculated using methodology of ICRP-60 to be compared with the dose limit of 1 mSv (see Appendix 1). Exposures due to marine pathways include the far-field effects of discharges of liquid waste from Sellafield. Unless stated otherwise, the critical group is represented by adults

<sup>c</sup> The contributors that give rise to more than 10% to the dose; either 'ext' to represent the whole body external exposure from beta or gamma radiation, 'beta' for beta radiation of skin or a radionuclide name to represent a contribution from internal exposure. The source of the radiation listed as contributing to the dose may not be discharged from the site specified, but may be from those of an adjacent site or other sources in the environment such as weapons fallout

<sup>d</sup> The assessed contribution is based on data at limits of detection.

<sup>e</sup> Dominant source of exposure. G for gaseous wastes. L for liquid wastes or surface water near solid waste sites. See also footnote 'c'

<sup>f</sup> The estimates for marine pathways include the effects of liquid discharges from LLWR. The contribution due to LLWR is negligible

<sup>g</sup> Exposure to skin including a component due to natural sources of beta radiation, to be compared with the dose limit of 50 mSv (see Appendix 1)

<sup>h</sup> 10 y old

<sup>i</sup> Includes a component due to natural sources of radionuclides

<sup>j</sup> 1 y old

<sup>k</sup> Excluding the effects of artificial radionuclides from Sellafield

<sup>l</sup> Excluding the effects of enhanced concentrations due to the legacy of discharges of naturally occurring radionuclides from a phosphate processing works, Whitehaven

<sup>m</sup> Including the effects of enhanced concentrations due to the legacy of discharges of naturally occurring radionuclides from a phosphate processing works, Whitehaven

<sup>n</sup> Including the effects of artificial radionuclides from Sellafield

<sup>o</sup> Water is from rivers and streams and not tap water

<sup>p</sup> Prenatal children

## 2. Nuclear fuel production and reprocessing

### Key points

- *Total doses* for the representative person were 42 per cent (or less) of the dose limit for all assessed sites. *Total doses* increased in the vicinity of Sellafield, compared to the values in 2014, but remained well below the limit
- Doses, discharges, environmental concentrations and dose rates in 2015 were broadly similar to those in 2014

### Capenhurst, Cheshire

- *Total dose* for the representative person was 13 per cent of the public dose limit and decreased in 2015
- Gaseous discharges of uranium increased in 2015

### Springfields, Lancashire

- *Total dose* for the representative person was 5 per cent of the public dose limit and unchanged in 2015. The highest exposure was represented by occupancy on a houseboat
- Gaseous discharges of uranium decreased and liquid discharges of uranium increased in 2015
- Gamma dose rates in the vicinity of the houseboats were similar in 2015

### Sellafield, Cumbria

- *Total doses* for the representative person were 42 per cent (or less) of the public dose limit in 2015, up from 22 per cent in 2014
- The highest *total doses* were from seafood affected by past phosphate processing at Whitehaven. Historical discharges from Sellafield made a lesser contribution
- The consumption of seafood contributed to approximately 96 per cent of the *total dose*
- Radiation dose from natural radionuclides was significantly higher in 2015, mostly due to an increase in polonium-210 concentrations in locally caught crustaceans. The contribution to *total dose* from Sellafield discharges also increased due to higher less than values for iodine-129 in fish and shellfish
- Gaseous discharges generally decreased, except for antimony-125 which increased, in 2015
- Liquid discharges were generally similar in 2015
- Concentrations of Sellafield derived radionuclides and dose rates were generally similar to those in 2014. Plutonium radionuclides and americium-241 were generally similar in shellfish

This section considers the results of monitoring, by the Environment Agency, FSA, NIEA and SEPA, of three sites in the UK associated with civil nuclear fuel production and reprocessing. These sites are at:

Capenhurst, a site where uranium enrichment is carried out and management of uranic materials and decommissioning activities are undertaken; Springfields, a site where fuel for nuclear power stations is fabricated; Sellafield, a site where irradiated fuel from nuclear power stations is reprocessed.

The Capenhurst site is owned partly by Urenco UK Limited (UUK) and partly by NDA. UUK holds the Site Licence, and their main commercial business is production of enriched uranium for nuclear power stations. NDA's legacy storage and decommissioning activities are managed by an Urenco Group company, Capenhurst Nuclear Services Limited (CNS). Another Urenco Group company, Urenco ChemPlants Limited (UCP), is currently building a new facility (Tails Management Facility, planned to be commissioned in 2017) on a separate part of the site.

Both the Springfields and Sellafield sites are owned by NDA. The Springfields site is leased long-term to

Springfields Fuels Limited, who carry out nuclear fuel manufacture and other commercial activities and also have a contract with NDA to decommission legacy facilities on the site. The main operations on the Sellafield site are fuel reprocessing, decommissioning and clean-up of redundant nuclear facilities, and waste treatment and storage. Since 2008, the operating model at Sellafield was based on the Parent Body Organisation (PBO) concept, with Nuclear Management Partners Limited (NMP) being the PBO for the Sellafield Site Licence Company (SLC), Sellafield Limited. In 2015, the Government approved the NDA's recommendation for a change to the management arrangements at Sellafield. On 1 April 2016, NDA became the owner of Sellafield Limited, the Site Licence Company responsible for managing and operating Sellafield on behalf of the NDA, replacing the previous management model of ownership by the private sector.

The Windscale site, also owned by NDA, is located on the Sellafield site and in 2008 the site licence for Windscale was transferred to Sellafield Limited, integrating the Windscale and Sellafield sites. Windscale is discussed in Section 2.4. Note that the LLWR site near Drigg is separate from Sellafield and is discussed in Section 7.1.

Gaseous and liquid discharges from each of these sites are regulated by the Environment Agency. In 2015, gaseous and liquid discharges were below permit limits for each of the sites (see Appendix 2). The medium-term trends in doses, discharges and environmental concentrations at these sites have been considered in a summary report (Environment Agency, FSA, NIEA and SEPA, 2010b).

## 2.1 Capenhurst, Cheshire



The site, near Ellesmere Port, was previously split into two adjacent nuclear licensed sites at Capenhurst. One nuclear licensed site was owned by NDA, comprising uranic material storage facilities and activities associated

with decommissioning, and the other owned by UUK, operating three plants producing enriched uranium for nuclear power stations. In 2012, NDA completed the transfer of its Capenhurst site with the transition of Sellafield Limited activities to CNS, creating one nuclear licensed site owned and managed by UUK. The major operators at the site are UUK, CNS and UCP. UCP are currently constructing a new facility, to allow safer long-term storage of depleted uranium, on a separate part of the site. This facility, the Tail Management Facility, will de-convert Uranium Hexafluoride ( $UF_6$ ), or “tails” to Uranium Oxide ( $U_3O_8$ ) to allow the uranium to be stored in a more chemically stable oxide form for potential future reuse in the nuclear fuel cycle and will recover hydrofluoric acid for reuse in the chemical industry. It is anticipated that this facility will become commissioned in 2017. The plant is permitted and, when commissioned, will discharge gaseous waste to the environment, aqueous waste to UUK’s effluent disposal system and will dispose of solid waste by off-site transfer.

The most recent collaborative habits survey undertaken on behalf of the Environment Agency, the FSA and ONR (including all exposure pathways from liquid discharges, gaseous discharges and direct radiation) was conducted in 2008 on behalf of the Environment Agency, the FSA and the Health and Safety Executive (HSE) (Tipple *et al.*, 2009). However, in 2013, an independent habits survey was carried out by Cefas on behalf of UUK. The main aim of the UUK survey was to collect occupancy data for people exposed to direct radiation from the site (living or spending time within 1 km of the site). The UUK survey showed an increase in occupancy in comparison to an equivalent direct radiation survey conducted in 2004.


## Doses to the public

The *total dose* from all pathways and sources is assessed to have been 0.13 mSv (Table 2.1) in 2015, or 13 per cent of the dose limit, and down from 0.17 mSv in 2014. This dose was almost entirely due to direct radiation from the Capenhurst site. The dose assessment identifies a local adult living near to the site as the representative person. The decrease in *total dose* (from 2014) was mostly attributed to a lower estimate of direct radiation from the site in 2015. The trend in *total dose* over the period 2004 – 2015 is given in Figures 1.2 and 2.1. Any changes in *total doses* with time are attributable to changes in the estimates of direct radiation from the site.


Source specific assessments give exposures for high-rate consumers of locally grown foods, and for children playing in and around Rivacre Brook, were less than the *total dose* in 2015 (Table 2.1). The dose for 10 year-old children (who play near the brook and may inadvertently ingest water and sediment) was 0.010 mSv in 2015 and similar to those in recent years. The dose is estimated using cautious assumptions for occupancy of the bank of the brook, inadvertent ingestion rates of water and sediment and gamma dose rates.

## Gaseous discharges and terrestrial monitoring

Uranium is the main radioactive constituent of gaseous discharges from Capenhurst, with small amounts of other radionuclides present in discharges by CNS Limited (previously Sellafield Limited). UUK discharges of uranium increased, in comparison to releases in 2014. The increase in annual emissions in 2015 has been investigated by UUK (see Table A2.4 for further information). The discharges remain within permitted limits. The focus for terrestrial sampling was on the content of technetium-99 and uranium in food (including milk), grass and soil. Results for 2015 are given in Table 2.2(a). Concentrations of radionuclides in milk and food samples around the site were very low and similar to previous years. Technetium-99 and uranium concentrations in soils were also low, whilst technetium-99 concentrations in food and grass are reported as less than values, in 2015. Figure 2.2 shows the trend of technetium-99 concentrations in grass from 2004. The overall trend reflects the reductions in discharges of technetium-99 from recycled uranium. The most recently observed variability (from year to year) in the technetium-99 concentrations is largely based on data reported as less than values. In future, the enrichment of reprocessed uranium is anticipated to increase, which may lead to increases in discharges of technetium-99 and neptunium-237, if recycled uranium is processed. However, no increase is expected in the discharge limits.



**Figure 2.1.** Total dose at nuclear fuel production and reprocessing sites, 2004-2015  
 (Exposures at Sellafield/Whitehaven/LLWR receive a significant contribution to the dose from technologically enhanced naturally occurring radionuclides from previous non-nuclear industrial operations)



**Figure 2.2.** Technetium-99 annual discharges from and concentrations in grass at Capenhurst, 2004-2015

## Liquid waste discharges and aquatic monitoring

The UUK permit for the Capenhurst site allows liquid waste discharges to the Rivacre Brook for uranium and uranium daughters, technetium-99 and non-uranium alpha (mainly neptunium-237). In 2015, discharges from Capenhurst were similar to those in recent years.

Monitoring included the collection of samples of fish and shellfish from the local marine environment (for analysis of a range of radionuclides) and of freshwater and sediments for the analysis of tritium, technetium-99, gamma emitting radionuclides, uranium, neptunium-237, and gross alpha and beta. Dose rate measurements were taken on the banks of the Rivacre Brook. Results for 2015 are given in Tables 2.2(a) and (b). Concentrations of radionuclides in foods from the local marine environment and dose rates were very low and generally similar to those in previous years. Thorium-234 in cockles was detected in 2015, reported just above the less than value. Downstream of the Rivacre Brook (at the location where children play), dose rates were generally similar to those in 2014. The low concentrations in fish and shellfish reflect the distant effects of discharges from Sellafield. As in previous years, sediment samples from the Rivacre Brook contained very low but measurable concentrations of uranium (enhanced above natural levels) and technetium-99. Some enhancement of these radionuclides was measured close to the discharge point. Variations in concentrations in sediment from the brook are to be expected due to differences in the size distribution of the sedimentary particles. Concentrations of radionuclides in freshwaters were also very low. As in recent years, measured dose rates were higher, relative to natural background, near to the discharge point.

## 2.2 Springfields, Lancashire



The Springfields site at Salwick, near Preston, is operated by Springfields Fuels Limited (SFL) under the management of Westinghouse Electric UK Limited, on behalf of NDA. The main commercial activity is the manufacture of fuel elements for

nuclear reactors and the production of uranium hexafluoride. Other important activities include recovery of uranium from residues and decommissioning redundant plant, under contract to NDA, who retain responsibility for the historic nuclear liabilities on the site. Research and development, carried out by the National Nuclear Laboratory, produces small amounts of other gaseous

radionuclides that are also discharged under permit (see Appendix A2.1).

Monitoring around the site is carried out to check not only for uranium concentrations, but also for other radionuclides discharged in the past (such as actinide daughter products from past discharges when uranium ore concentrate (UOC) was the main feed material) and for radionuclides discharged from Sellafield. The monitoring locations (excluding farms) used to determine the effects of gaseous and liquid discharges are shown in Figure 2.3.

The most recent habits survey was undertaken in 2012 (Ly *et al.*, 2013). In 2015, based on a five-year rolling average (2011 – 2015), the habits information was revised, resulting in a slightly lower occupancy rate for high-rate houseboat dwellers. Revised figures for consumption rates, together with occupancy and handling rates, are provided in Appendix 1 (Table X2.2).

## Doses to the public

In 2015, the *total dose* from all pathways and sources is assessed to have been 0.050 mSv (Table 2.1), or 5 per cent of the dose limit, and unchanged from 2014. The representative person was an adult houseboat dweller in a boatyard, who was exposed to external radiation from activity in muddy sediments. Gamma dose rate measurements were not taken aboard a houseboat in 2015 (unlike in 2014). Therefore, dose rates were derived by using measurements outside the houseboat (in 2015), and then adjusting these values by the ratio of on-board and outside dose rates from results reported in 2014. This information was directly applicable to the locations where high-rate occupancy was taking place. *Total doses* (together with dose rates) over the period 2004 – 2015 are given in Figure 2.4. Most recently, the estimated *total dose* has decreased following on from direct measurements (beneath houseboats) being available in 2014.

Source specific assessments indicated that exposures were all less than or similar to the *total dose* (Table 2.1) for;

- Consumers of locally grown food and of seafood
- High-occupancy houseboat dwellers in the Ribble Estuary
- Children playing on the banks of the estuary
- Farmers spending time on the banks of the estuary
- Wildfowling consuming game obtained from the estuary area

In 2015, the source specific assessment gave an estimated dose to a high-occupancy houseboat dweller of 0.054 mSv or approximately 5 per cent of the dose limit for members of the public of 1 mSv, and generally similar to that in 2014 (0.056 mSv). The lower value in 2015 was due to the revised habits information (a small reduction in the occupancy rate). This value is higher, by a small amount, than the *total dose* of 0.050 mSv assessed for the




Figure 2.3. Monitoring locations at Springfields, 2015 (not including farms)




Figure 2.4. Total dose from all sources and dose rates at Springfields, 2004-2015

same representative person. The *total dose* assessment is based on more realistic assumptions. The dose to the representative person for high-rate consumers of seafood (including a contribution from external exposure) was 0.020 mSv in 2015. Of this dose, approximately 0.019 mSv was from external exposure and the remainder was from the consumption of fish and shellfish. The dose in 2014 was 0.021 mSv. The most important radionuclides were caesium-137 and americium-241 from past discharges from the Sellafield site.

As in recent years, assessments were undertaken to determine the dose to wildfowlers from external exposure over salt marsh and the consumption of game, the dose

to farmers from external exposure, the dose to high-rate consumers of locally grown food and the dose to children playing on the banks of the estuary, at Springfields. The estimated doses in 2015 were 0.006 mSv, 0.034 mSv, less than 0.005 mSv and less than 0.005 mSv, respectively, for these pathways (Table 2.1).

It has been previously shown that assessed doses to the public from inhaling Ribble Estuarine sediment re-suspended in the air were much less than 0.001 mSv, and negligible in comparison with other exposure routes (Rollo *et al.*, 1994).

## Gaseous discharges and terrestrial monitoring

Uranium is the main radioactive constituent of gaseous discharges, with small amounts of other radionuclides present in discharges from the National Nuclear Laboratory's research and development facilities. Discharges of uranium decreased in 2015, in comparison to releases in 2014.

The focus of the terrestrial sampling was for the content of tritium, carbon-14, strontium-90, iodine-129, and isotopes of uranium, thorium, plutonium and americium in milk, and vegetables. Grass and soil samples were collected and analysed for isotopes of uranium. Data for 2015 are given in Table 2.3(a). As in previous years, elevated concentrations of uranium isotopes were found in soils around the site, but the isotopic ratio showed they are most likely to be from natural abundance. Low concentrations of thorium were found in vegetable and grass samples. The carbon-14 concentration in beetroot was lower (in comparison to that in 2014), and similar to the default value used to represent the background level. Most other concentrations of radionuclides are reported as less than values. Results were broadly similar to those of previous years.

Figure 2.5 shows the trends over time (2004 – 2015) of uranium discharges and total uranium radionuclide concentrations in food (cabbage; 2004 – 2013; beetroot; 2014 – 2015). Over the period, concentrations of uranium were also found in soil around the site, but the isotopic ratio showed that they were naturally occurring. Total uranium was detected in cabbage and beetroot samples during the period (no data in 2006), but the concentrations were very low. The apparent peak of uranium in cabbage in 2007 was also low and significantly less than that found in soil samples.

## Liquid waste discharges and aquatic monitoring


Permitted discharges of liquid waste (including gross alpha and beta, technetium-99, thorium-230, thorium-232, neptunium-237, uranium and other transuranic radionuclides) are made from the Springfields site to the Ribble Estuary by two pipelines. Discharges of uranium increased in 2015, in comparison to releases in 2014. Other discharges were generally similar to those in recent years, including the short half-life beta-emitting radionuclides (mostly thorium-234) that have decreased following the end of the UOC purification process in 2006. Process improvements in the uranium hexafluoride production plants on the Springfields site have reduced the amounts of other uranium compounds needing recycling; these improvements, alongside a reduction in legacy uranic residue processing, have led to a corresponding reduction in discharges of uranium in most recent years. Discharges of technetium-99 depend almost entirely on

which legacy uranic residues are being processed. Since completion of one particular residue processing campaign (around the end of 2012), technetium-99 discharges have also decreased. In 2015, technetium-99 discharges were decreased, by a small amount, in comparison to those in 2014. The Ribble Estuary monitoring programme consisted of dose rate measurements, and the analysis of sediments for uranium and thorium isotopes, and gamma emitting radionuclides.

Locally obtained fish and shellfish were analysed by gamma-ray spectrometry and for thorium and plutonium isotopes. Results for 2015 are shown in Tables 2.3(a) and (b). As in previous years, radionuclides due to discharges from both Springfields and Sellafield were found in the Ribble Estuary sediment and biota. Radionuclides found in the Ribble Estuary originating from Sellafield were technetium-99, caesium-137 and americium-241. Isotopes of uranium and the short half-life radionuclide thorium-234, from Springfields, were also found. Concentrations of the latter were closely linked to recent discharges from the Springfields site. In 2015, thorium-234 concentrations in sediments (over the range of sampling sites) were generally similar compared to those in recent years. Over a much longer timescale (2004 – 2015), these concentrations have declined due to reductions in discharges as shown by the trend of sediment concentrations at the outfall, Lower Penwortham and Beconsall (Figure 2.5). The most significant change in the discharge trends was the step reduction of short half-life beta emitting radionuclides in liquid discharges, mostly thorium-234. The reduction was because the UOC purification process ended in 2006.

Caesium-137, americium-241 and plutonium radionuclides were found in biota and sediments from the Ribble Estuary in 2015. The presence of these radionuclides was due to past liquid discharges from Sellafield, carried from west Cumbria into the Ribble Estuary by sea currents and adsorbed on fine-grained muds. The concentrations observed were generally similar to those in recent years.

Figure 2.5 also provides trend information over time (2004 – 2015) for a number of other permitted radionuclides and activity concentrations in food. Liquid discharges of uranium radionuclides steadily decreased (and other discharges to a lesser extent) over the whole period, whilst technetium-99 discharges generally decreased overall (but peaked in 2012). Caesium-137 concentrations in flounder and salmon showed variations between years and this was most likely due to natural changes in the environment, although there is some evidence that concentrations in flounder decreased overall. Concentrations of technetium-99 in shrimps generally declined over the whole period, consistent with the reduction in technetium-99 discharges from Sellafield (Figure 2.14); the apparent increases in recent years was due to the inclusion of reported less than values.



**Figure 2.5.** Discharges of gaseous and liquid radioactive wastes and monitoring of the environment, Springfields 2004–2015 (Note different scales used for discharges and activity concentrations)

Gamma dose rates in the estuary were generally higher than expected natural background levels (see Appendix 1, Section 3.7), and this is due to Sellafield-derived gamma-emitting radionuclides (caesium-137 and americium-241). In 2015, gamma dose rates in the estuary, excluding rates taken for houseboat assessments, were generally similar to those in 2014, but with some small variations at some sites. In 2015, gamma dose rates measured in the vicinity of houseboat dwellers (at Becconsall and Freckleton), and beta dose rates from sediments, were generally similar to those in 2014 (where comparisons can be made from similar ground types and locations).

## 2.3 Sellafield, Cumbria



This site is operated by Sellafield Limited but is owned by NDA. The main operations on the Sellafield site are: fuel reprocessing at the Magnox Reprocessing Plant and the Thermal Oxide Reprocessing Plant (THORP); decommissioning and clean-up of redundant nuclear facilities; and waste treatment and storage. The site also contains the Calder Hall Magnox nuclear power station, which ceased generating in 2003 and is undergoing decommissioning.

De-fuelling of Calder Hall is expected to be completed between March 2017 and March 2019 according to the NDA Magnox Operating Plan. The Windscale site is located at Sellafield, and is discussed in Section 2.4.

In 2011, Sellafield Limited and NDA published their plans for decommissioning of the Sellafield site ([http://www.sellafieldsites.com/wp-content/uploads/2012/08/Sellafield\\_Plan.pdf](http://www.sellafieldsites.com/wp-content/uploads/2012/08/Sellafield_Plan.pdf)). In December 2014, an improved performance plan for Sellafield was accepted by NDA ([http://www.sellafieldsites.com/wp-content/uploads/2015/03/Performance-Plan\\_single.pdf](http://www.sellafieldsites.com/wp-content/uploads/2015/03/Performance-Plan_single.pdf)), and replaces the 2011 Sellafield Plan. Sellafield Limited commenced the first retrievals of sludge from legacy pond facilities in early 2015 and continues to prepare for retrievals of intermediate level waste from legacy facilities and to reduce environmental risk. Some of these projects have the potential to impact on discharges to the environment. In 2015, a number of decommissioning projects continued including that of the Calder Hall reactors.

During the financial year 2015/16, 460 \*TeU of spent oxide fuel was reprocessed in THORP, compared with an original performance target of 435 TeU, and was the highest total reprocessing throughput since NDA took ownership of the site. The reprocessing of spent Magnox fuel for 2015/16 was a total of 390 TeU, compared with an original performance target of 477 TeU. The reprocessing of the remaining fuel is scheduled to end in 2018 and 2020 for THORP and Magnox reprocessing, respectively.

Every five years, a full habits survey is conducted in the vicinity of the Sellafield site which investigates the exposure pathways relating to liquid and gaseous discharges, and direct radiation. Annual review habits surveys are undertaken between these full habits surveys. These annual surveys investigate the pathways relating to liquid discharges, review high-rate fish and shellfish consumption by local people (known as the Sellafield Fishing Community) and review their intertidal occupancy rates. The most recent full habits survey was conducted in 2013 (Clyne *et al.*, 2014). In 2015, changes were found in the amounts and mixes of species consumed from the full habits survey conducted in 2013 (Garrod *et al.*, 2015a). Further afield, the most recent habits surveys were conducted in 2012, to determine the consumption and occupancy rates by members of the public on the Dumfries and Galloway coast (Garrod *et al.*, 2013a) and around Barrow and the south-west Cumbrian coast (Garrod *et al.*, 2013b). The results of this survey are used to determine the potential exposure pathways relating to permitted liquid discharges from the Sellafield nuclear licensed site in Cumbria. Revised figures for consumption

rates, together with occupancy rates, are provided in Appendix 1 (Table X2.2).

Habits surveys to obtain data on activities undertaken on beaches relating to potential public exposure to radioactive particles in the vicinity of the Sellafield nuclear licensed site were undertaken in 2007 and 2009 (Clyne *et al.*, 2008; Clyne *et al.*, 2010a).

Monitoring of the environment and food around Sellafield reflects the historical and present day site activities. In view of the importance of this monitoring and the assessment of public radiation exposures, the components of the programme are considered here in depth. The discussion is provided in four sub-sections, relating to the assessment of dose, the effects of gaseous discharges, the effects of liquid discharges and unusual pathways of exposure identified around the site.

### 2.3.1 Doses to the public

#### **Total dose from all pathways and sources**

The *total dose* from all pathways and sources is assessed using consumption and occupancy data from the full habits survey of 2013 (Clyne *et al.*, 2014) and the yearly review in 2015 (Clyne and Garrod, 2016). Calculations are performed for four age groups (adult, 10y, 1y and prenatal). The effects on high-rate consumers of fish and shellfish from historical discharges of naturally occurring radionuclides from non-nuclear industrial activity from the former phosphate works at Whitehaven are included to determine their contribution to the *total dose*. These works were demolished in 2004 and the authorisation to discharge radioactive wastes was revoked. The increase in concentrations of naturally occurring radionuclides due to the historical discharges is difficult to determine above a variable background (see Appendix 1).

In 2015, the highest *total dose* in the vicinity of Sellafield was assessed to have been 0.42 mSv, or 42 per cent of the dose limit to members of the public (Table 2.17). Most of this dose was due to radioactivity from sources other than those resulting from Sellafield discharges. The representative person was an adult, consuming molluscan shellfish at high-rates near Sellafield (and unchanged from 2014). The person also consumed significant quantities of other seafood. This represents a significant increase in *total dose* in 2015, from 0.22 mSv in 2014. This increase in *total dose* was mostly attributable to an increase in the concentrations of polonium-210 in locally caught crustaceans (lobsters and crabs) in 2015, in comparison to those in 2014. Polonium-210 is an important radionuclide in that small changes in levels above background significantly influence the dose contribution from these radionuclides and similarly the value of the estimated dose. A much lesser enhancement in *total dose* was due to the revision of habits information (i.e. from an increase in the

---

\* The twenty-foot equivalent unit (TEU) is an inexact unit of cargo capacity often used to describe the capacity of container ships and container terminals. It is based on the volume of a 20-foot-long (6.1 m) intermodal container, a standard-sized metal box which can be easily transferred between different modes of transportation, such as ships, trains and trucks.

consumption rates of crustaceans (compared to those in 2014). Direct radiation from the Sellafield site (0.005 mSv, Table 1.1) was considered in the *total dose* assessments, but this made an insignificant contribution to the highest *total dose*.

In percentage terms, the most significant contributors to the *total dose* in 2015 were from crustacean consumption, mollusc consumption, external exposure over sediments and fish consumption (81, 10, 4 and 4 per cent, respectively), the most important radionuclides were polonium-210, americium-241, and plutonium-239+240 (80, 7, and 3 per cent, respectively).

Artificial radionuclides discharged by Sellafield (including external radiation) and historical discharges of naturally occurring radionuclides from Whitehaven contributed 0.078 mSv and 0.35 mSv, respectively (values are rounded to two significant figures). In 2014, the contributions were 0.068 mSv and 0.15 mSv, respectively. In 2015, the contribution from the external radiation was approximately 0.017 mSv (0.017 mSv in 2014). Data for naturally occurring radionuclides in fish and shellfish, and their variation in recent years, are discussed in Section 7.

The contribution to the *total dose* of 0.078 mSv in 2015 from artificial radionuclides (including external radiation) was higher than in 2014 (0.068 mSv). In 2015, the contributing radionuclides were mostly americium-241, iodine-129, plutonium-239+240 and carbon-14 (33, 15, 14 and 6 per cent, respectively). External exposure was 22 per cent (25 per cent in 2014) of the *total dose* from artificial radionuclides. The increase in the contribution to the *total dose* from 2014 was mostly attributed to the increase in less than values for iodine-129 in fish and shellfish (given in footnotes, Tables 2.5 and 2.6, respectively) in the 2015 assessment.

The contribution to the *total dose* of 0.35 mSv in 2015 from naturally occurring radionuclides was significantly higher than in 2014 (0.15 mSv). In 2015, the most contributing radionuclide was polonium-210 (97 per cent). The increase in the contribution to the *total dose* from 2014 was mostly due to an increase in the concentrations of polonium-210 in locally caught crustaceans (lobsters and crabs) in 2015, in comparison to those in 2014. In 2015, polonium-210 in locally caught lobsters and crabs contributed 0.24 mSv and 0.089 mSv, respectively (values are rounded to two significant figures) to the *total dose*. Polonium-210 concentrations (above expected background) in mollusc samples contributed 0.007 mSv to the *total dose* in 2015, in comparison to 0.022 mSv in 2014; between 2011 and 2013 there was no contribution from molluscs.

Contributions to the highest *total dose* each year, from all pathways and sources by specific radionuclides, are given in Figure 2.6 over the period 2004 – 2015. The trend of generally declining dose broadly reflects a general reduction in concentrations in seafood of both naturally


occurring and artificial radionuclides from the non-nuclear and nuclear industries respectively. Inter-annual variations were more complex and governed by both natural variability in seafood concentrations and real changes in the consumption and occupancy characteristics of the local population.

The larger step changes (from 2004 to 2005, from 2008 to 2009 and from 2012 to 2013) were due to variations in naturally occurring radionuclides (mainly polonium-210 and lead-210). The changes in *total dose* in the intervening years from 2005 to 2007 were mainly a result of changes in seafood consumption rates. The decrease in 2010 was due to both reductions in naturally occurring radionuclides concentrations (polonium-210) and consumption rates, whilst the variation in the radionuclide contributors in 2011 (from previous years) resulted from a change in the representative person (from a consumer of molluscan shellfish to locally harvested marine plants). The largest proportion of the *total dose*, up till 2008 and again from 2011 – 2012 and 2014 – 2015, was mostly due to enhanced naturally occurring radionuclides from the historical discharges at Whitehaven and a smaller contribution from the historical discharges from Sellafield. From 2008 to 2010, the net result of progressive reductions of the naturally occurring radionuclides contribution to the *total dose* has been a relative increase in the proportion from artificial radionuclides. In 2013, the highest *total dose* (relating to the effects of Sellafield) was entirely due to external radiation from sediments. The change was due to both decreases in naturally occurring radionuclides concentrations (polonium-210) and a revision of habits information, resulting in a change in the representative person. In the following year (2014), the change in *total dose* was due to (i) a significant increase in the proportion of lobsters in the diet of high-rate seafood consumers and (ii) a return to a wide range of seafood species being consumed by individuals.

Other age groups received less exposure than the adult *total dose* of 0.42 mSv in 2015 (10y: 0.21; 1y: 0.13; prenatal: 0.072, rounded to two significant figures). *Total doses* estimated for each age group may be compared with an average dose of approximately 2.2 mSv to members of the UK public from all natural sources of radiation (Watson *et al.*, 2005) and to the annual dose limit to members of the public of 1 mSv.

### **Total dose from gaseous discharges and direct radiation**

In 2015, the dose to a representative person receiving the highest *total dose* from the pathways predominantly relating to gaseous discharges and direct radiation was 0.008 mSv (Table 2.17) and generally similar to that in 2014 (0.009 mSv). The most exposed age group in 2015 was a 10 year-old and the dominant contribution to this dose was direct radiation from the site. This represents a change in the representative person from an adult in



**Figure 2.6.** Contributions to *total dose* from all sources at Sellafield, 2004-2015 (The highest *total dose* in 2013 due to Sellafield discharges was to people living on houseboats near Barrow in Cumbria)

2014. The most significant contributors in 2015 to the *total dose* for a 10 year-old were from direct radiation from the site, the consumption of potatoes, and the combined external and inhalation from the site (60, 25, and 6 per cent, respectively), the most important radionuclide was ruthenium-106 (14 per cent). Other ages received less exposure than the 10 year-old *total dose* of 0.008 mSv in 2015 (adult: 0.008; 1y: 0.008; prenatal: 0.007, equivalent values rounded to one significant figure).

Contributions to the highest *total dose* each year, by specific radionuclides, are given in Figure 2.7 over the period 2004 – 2015. Up until 2007, there was a small decline in *total dose* due to a general reduction in concentrations of radionuclides in food and the environment caused, in part, by reductions in discharges in this period and beforehand. The main feature in the changes in *total dose* over the whole period was the increase in 2009. This resulted from an increase of total radiocaesium in game collected near the site. There is no evidence to suggest that this was caused by a change in site operations. Over the period 2010 – 2014, *total doses* were generally similar between years. The lower *total dose* values in most recent years was mostly due to changes in the monitoring programme in 2014 (Environment Agency, FSA, FSS, NIEA, NRW and SEPA, 2015).

### Total dose from liquid discharges

The people receiving the highest *total dose* from the pathways predominantly relating to liquid discharges are given in Table 2.17. Each *total dose* is the same as that giving their maximum *total dose* for all sources and pathways.

### Source specific doses


Important source specific assessments of exposures, as a result of radioactive waste discharges from Sellafield, continued to be due to high-rate consumption of fish and shellfish and to external exposure from gamma rays over long periods. Other pathways were kept under review, particularly high-rate consumption of locally grown food (from atmospheric discharges), to account for the potential for sea to land transfer at the Ravenglass Estuary to the south of the site and exposure from contact with beta emitters during handling of sediments and/or handling of fishing gear.

### Doses from terrestrial food consumption

In 2015, a 1 year-old child, who was a high-rate consumer of milk and was exposed to external and inhalation pathways from gaseous discharges, received the highest dose for all ages. The estimated dose was 0.012 mSv (Table 2.17) or approximately 1 per cent of the dose limit to members of the public, and unchanged from 2014. Other age groups received less exposure than the 1 year-old child dose of 0.012 mSv in 2015 (adult: 0.008; 10y: 0.010; prenatal: 0.007).

### Doses from seafood consumption

Two sets of habits data are used in these dose assessments. One is based on the habits information seen in the area each year (2015 habits survey). The second is based on a five-year rolling average using habits data gathered from 2011 to 2015. Changes were found in the amounts and mixes of species consumed. For molluscs, the consumption rates increased in 2015, but decreased for the 2011 – 2015 data set. For fish and crustacean, consumption



**Figure 2.7.** Contributions to *total dose* from gaseous discharge and direct radiation sources at Sellafield, 2004-2015 (+ based on limits of detection for concentrations in foods)

rates increased for both 2015 and 2011 – 2015 data sets. The occupancy rate over sediments decreased in 2015, but increased for the 2011 – 2015 data set. The revised habits data are given in Appendix 1 (Table X2.2). Aquatic pathway habits are normally the most important in terms of dose near Sellafield and are surveyed every year. This allows generation of a unique yearly set of data and also rolling five-year averages. The rolling averages are intended to smooth the effects of sudden changes in habits and provide an assessment of dose that follows more closely changes in radioactivity concentrations in food and the environment. These are used for the main assessment of doses from liquid discharges, and follow the recommendations of the report of the Consultative Exercise on Dose Assessments (CEDA) (FSA, 2001a).

Table 2.17 summarises source specific doses to seafood consumers in 2015. The doses from artificial radionuclides to people, who consume a large amount of seafood, were 0.098 mSv and 0.084 mSv using the annual and five-year rolling average habits data, respectively. These doses each include a contribution due to external radiation exposure over sediments. Doses were also generally similar in comparison to those in 2014 (0.094 mSv and 0.089 mSv, respectively),

The dose to a local person (high-rate consumer of seafood) due to the enhancement of concentrations of naturally occurring radionuclides from former non-nuclear industrial activity in the Sellafield area (using maximising assumptions for the dose coefficients and the five-year rolling average habits data) is estimated to have been 0.29 mSv in 2015. Most of this was due to polonium-210 (97 per cent). The reason for the large increase in dose in 2015 (from 0.13 mSv in 2014) is the same as that contributing to the *total dose* for all sources, i.e. mostly due to an increase in the concentrations of polonium-210 in locally caught crustaceans (lobsters and crabs), in comparison to those in

2014. For comparison (with the assessment using the five-year rolling average habits data), the dose from the single-year assessment for the Sellafield seafood consumer from naturally occurring radionuclides (based on consumption rates and habits survey data in 2015) was 0.42 mSv (Table 2.17).

Taking artificial and enhanced natural radionuclides together, the source specific doses were 0.52 mSv and 0.37 mSv (values are rounded to two significant figures) for annual and five-year rolling average habits data, respectively. These estimates are larger and smaller, respectively, than the estimate of *total dose* from all sources of 0.42 mSv. The main reason for this is a difference in the approach to selecting consumption rates for seafood for the representative person. The differences in dose are not unexpected, are within the uncertainties in the assessments and confirm *total dose* as a robust measure of exposure.

Exposures representative of the wider communities associated with fisheries in Whitehaven, Dumfries and Galloway, the Morecambe Bay area, Northern Ireland and North Wales have been kept under review in 2015 (Table 2.17). Those for fisheries in the Isle of Man and Fleetwood have been shown to be generally lower and dose data are available in earlier RIFE reports (e.g. Environment Agency, FSA, NIEA, NRW and SEPA, 2014). Where appropriate, the dose from consumption of seafood is summed with a contribution from external exposure over intertidal areas. The doses received in the wider communities were significantly less than for the local Sellafield population because of the lower concentrations and dose rates further afield. There were generally small changes in the doses (and contribution to doses) in each area in 2015 (Table 2.16), in comparison to those in 2014. All doses were well within the dose limit for members of the public of 1 mSv.

The dose to a person, who typically consumes 15 kg of fish per year from landings at Whitehaven is also given in Table 2.17. This consumption rate used represents an average for a typical consumer of seafood from the north-east Irish Sea. The dose was very low, less than 0.005 mSv in 2015.

## Doses from sediments

The main radiation exposure pathway associated with sediments is due to external dose from gamma-emitting radionuclides adsorbed on intertidal sediments in areas frequented by the public. This dose can make a significant contribution to the total exposure of members of the public in coastal communities of the north-east Irish Sea but particularly in Cumbria and Lancashire. Gamma dose rates currently observed in intertidal areas are mainly due to radiocaesium and naturally occurring radionuclides. For some people, the following pathways may also contribute to doses from sediments: exposure due to beta-emitters during handling of sediments or fishing gear; inhalation of re-suspended beach sediments; and inadvertent ingestion of beach sediments. These pathways are considered later: in the main, they give rise to only minor doses compared with those due to external gamma emitters.

Gamma radiation dose rates over areas of the Cumbrian coast and further afield in 2015 are given in Table 2.9. The results of the assessment of external exposure pathways are included in Table 2.17. The highest whole body exposures due to external radiation resulting from Sellafield discharges, past and present, was received by the representative person living in houseboats in the Ribble Estuary in Lancashire. In 2015, the dose was 0.054 mSv or approximately 5 per cent of the dose limit for members of the public. Other people received lower external doses in 2015. The dose received by a local houseboat dweller at Barrow (due to the effects of Sellafield discharges) was 0.051 mSv and similar to that in 2014 (0.055 mSv). The estimated dose to a person who spends a long time over the marsh in the Ravenglass Estuary was 0.013 mSv and similar to that in 2014 (0.012 mSv). Overall, gamma dose rates measurements in 2015 were generally similar to those in 2014 in the Ravenglass Estuary.

The doses to people in 2015 from a number of other activities were also estimated. Assessments were undertaken for a typical resident using local intertidal areas for recreational purposes at 300 hours per year, and for a typical tourist visiting the coast of Cumbria with a beach occupancy of 30 hours per year. The use by residents for two different environments, at a number of locations (at a distance from the Sellafield influence), were assessed: residents that visit and use beaches and residents that visit local muddy areas or salt marsh. Typical occupancy rates (Clyne *et al.*, 2008; 2010a) are assumed and appropriate gamma dose rates have been used from Table 2.9. The activities for the typical tourist include consumption of local seafood and occupancy on beaches. Concentrations

of radioactivity in fish and shellfish have been used from Tables 2.5 – 2.7, and appropriate gamma dose rates used from Table 2.9. The consumption and occupancy rates for activities of a typical resident and tourist are provided in Appendix 1 (Table X2.2).

In 2015, the doses to people from recreational use of beaches varied from 0.006 to 0.011 mSv with the higher doses being closer to the Sellafield source. The doses for recreational use of salt marsh and muddy areas had a greater variation from less than 0.005 to 0.013 mSv but were of a similar order of magnitude. The values for these activities were similar to those in recent years. The dose to a typical tourist visiting the coast of Cumbria, including a contribution from external exposure, was estimated to be less than 0.005 mSv.

## Doses from handling fishing gear and sediment

Exposures can also arise from contact with beta-emitters during handling of sediments, or fishing gear on which fine particulates have become entrained. Habits surveys keep under review the amounts of time spent by fishermen handling their fishing gear, and by bait diggers and shellfish collectors handling sediment. For those most exposed, the rates for handling nets and pots and for handling sediments are provided in Appendix 1 (Table X2.2). In 2015, the skin doses to a fisherman from handling fishing gear (including a component due to naturally occurring radiation), and a bait digger and shellfish collector from handling sediment, were 0.10 mSv and 0.064 mSv, respectively and both were less than 0.5 per cent of the appropriate annual dose limit of 50 mSv specifically for skin. Therefore, both handling of fishing gear and sediments continued to be minor pathways of radiation exposure.

## Doses from atmospheric sea to land transfer

At Ravenglass, an infant was the representative person from consuming terrestrial foods that were potentially affected by radionuclides transported to land by sea spray. In 2015, the dose (including contributions from Chernobyl and weapon test fallout) was estimated to be 0.017 mSv, which was less than 2 per cent of the dose limit for members of the public and unchanged from 2014. The largest contribution to the dose was from ruthenium-106 in milk (as in 2014). As in previous years, sea-to-land transfer was not of radiological importance in the Ravenglass area.

## Doses from seaweed and seawashed pasture

Doses from seaweed and seawashed pasture pathways were not assessed in 2015, due to earlier changes in the monitoring programme in 2014.

In South Wales the food item laverbread, made from the brown seaweed *Porphyra*, is eaten. Only small quantities of samphire, *Porphyra* and *Rhodomenia* (a red seaweed) are generally consumed and no harvesting of *Porphyra* in west Cumbria, for consumption in the form of laverbread, was reported in 2015. Estimated doses for a high-rate consumer of laverbread are available in earlier RIFE reports (e.g. Environment Agency, FSA, NIEA, NRW and SEPA, 2014) establishing that this exposure pathway is of low radiological significance.

Seaweeds are sometimes used as fertilisers and soil conditioners. Estimated doses for a high-rate consumer of vegetables (assuming these foods were obtained from the monitored plots near Sellafield) are available in earlier RIFE reports (e.g. Environment Agency, FSA, NIEA, NRW and SEPA, 2014). Overall doses from this pathway remain similar, and minor variations from year to year are due to different foods being grown and sampled from the monitored plots. Exposures of vegetable consumers using seaweed from further afield in Northern Ireland, Scotland and North Wales are expected to be much lower than near Sellafield.

Animals may graze on seaweeds on beaches in coastal areas. However, there was no evidence of this taking place significantly near Sellafield. FSA undertook an assessment of the potential dose to a high-rate consumer of meat and liver from sheep grazing the seaweed using data relevant to the Shetlands and Orkneys. This showed that doses would have been well within the dose limit of 1 mSv per year for members of the public in 1998 when concentrations of technetium-99 would have been at substantially higher levels than in 2015 (MAFF and SEPA, 1999). A further research study (relevant to the Scottish islands and coastal communities), conducted by PHE on behalf of the FSA and SEPA, investigated the potential transfer of radionuclides from seaweed to meat products and also to crops grown on land where seaweed had been applied as a soil conditioner (Brown *et al.*, 2009). The study concluded that the highest levels of dose to people using seaweed, as a soil conditioner or an animal feed, were in the range of a few microsieverts and the majority of the doses are at least a factor of 100 lower. The report is available on SEPA's website: <http://www.sepa.org.uk/environment/radioactive-substances/environmental-monitoring-and-assessment/reports/>

### 2.3.2 Gaseous discharges

Regulated discharges to atmosphere are made from a wide range of facilities at the site including the fuel storage

ponds, the reprocessing plants and waste treatment plants, as well as from Calder Hall Power Station. Discharges from Calder Hall are now much reduced since the power station ceased generating electricity in 2003. Discharges to atmosphere during 2015 are summarised in Appendix 2 (Table A2.1). The permit limits gaseous discharges for gross alpha and beta activities, and 13 specified radionuclides. In addition to overall site limits, individual limits have been set on discharges from the main contributing plants on site.


Discharges of gaseous wastes from Sellafield were much less than the permit limits in 2015. Discharges were generally decreased, although antimony-125 increased, in comparison to those in 2014.

### Monitoring around the site related to gaseous discharges

Monitoring of terrestrial foods in the vicinity of Sellafield is conducted by the FSA to reflect the scale of discharges from the site. This monitoring is the most extensive of that for the nuclear licensed sites in the UK. A range of foodstuffs was sampled in 2015 including milk, fruit, vegetables, meat and offal, game, and environmental materials (grass and soil). Samples were obtained from different locations around the site to allow for variations due to the influence of meteorological conditions on the dispersal of gaseous discharges. The analyses conducted included gamma-ray spectrometry and specific measurements for tritium, carbon-14, strontium-90, technetium-99, iodine-129, uranium and transuranic radionuclides.

The results of monitoring in 2015 are given in Table 2.4. The concentrations of all radionuclides around the site were low. Concentrations in terrestrial foodstuffs were generally similar to those in recent years. Concentrations of radionuclides in meat and offal from cattle and sheep were low with many reported as less than values with only limited evidence of the effects of Sellafield's atmospheric discharges detected in data for carbon-14 and strontium-90. Plutonium concentrations and americium-241 in wood pigeon, when detectable, were low and much lower than those found in seafood.

A range of fruit and vegetables was sampled in 2015 and the activity concentrations were generally similar to those found in previous years. In common with meat and offal samples, only limited evidence of the atmospheric discharges from Sellafield was found in some of these foods. In 2015, the maximum iodine-129 concentration in milk was reported as a less than value (unlike in recent years). Small enhancements (above expected background) in concentrations of carbon-14 were found in some food samples (including meat and offal), as in recent years. Concentrations of transuranic radionuclides, when detectable in these foods, were very low. Antimony-125 concentrations were below limits of detection in terrestrial



**Figure 2.8.** Discharges of gaseous wastes and monitoring of milk near Sellafield, 2004-2015

samples in 2015, despite relatively enhanced discharges in recent years. Trends in maximum concentrations of radionuclides in milk, and corresponding discharge levels, near Sellafield over the last decade are shown in Figure 2.8. Over the whole period, concentrations of carbon-14 were relatively constant (with some variation between years, generally consistent with changes in discharges), and caesium-137 concentrations (and strontium-90 to a lesser extent) were declining overall.

### 2.3.3 Liquid discharges


Regulated liquid discharges derive from a variety of sources at the site including the fuel storage ponds, the reprocessing plants, from the retrieval and treatment of legacy wastes, the laundry and from general site drainage. Wastes from these sources are treated and then discharged to the Irish Sea via the sea pipelines that terminate 2.1 km beyond low water mark. Liquid wastes are also discharged from the factory sewer to the River Ehen Estuary. In March 2015, the Environment Agency issued a varied permit to Sellafield Limited that will allow some liquid wastes to also be discharged via the Calder Interceptor Sewer. As part of this permit variation, a restructuring of the liquid discharge limits has introduced annual site limits (incorporating sea pipelines; factory sewer and Calder Interceptor Sewer) that are 10 per cent lower than the previous sea pipeline limits for total alpha, beta and tritium. This change reflects the removal of headroom rather than any change in the dose or risk that the discharges represent. Discharges from the Sellafield pipelines during 2015 are summarised in Appendix 2 (Table A2.2). The current permit sets limits on gross alpha and beta, and 16 individual nuclides. In addition to overall site limits, individual limits have been set on discharges from the main contributing plants on site (Segregated Effluent Treatment Plant, Site Ion Exchange Plant (SIXEP), Enhanced Actinide Removal Plant (EARP) and THORP). All of the discharges in 2015 were well below the limits in the permit. Liquid discharges were generally

similar in 2015, in comparison to those in 2014. Overall, the discharges continue to reflect the varying amounts of fuel reprocessed in the THORP and Magnox reprocessing plant, and periods of planned and unplanned reprocessing plant shutdown that occur from year to year.


Discharges of technetium-99 were low and similar in 2015, to those in 2014. The long-term downward trend, from their peak of 192 TBq in 1995, has continued (Figure 2.9). Technetium-99 discharges from Sellafield are now substantially reduced and met the target set for 2006 in the UK National Discharges Strategy (Defra, 2002). The reduction of technetium-99 discharges was due to the diversion, since 2003, of the Medium Active Concentrate (MAC) waste stream from Magnox reprocessing to vitrification and, between 2003 and 2007, use of a chemical precipitant (Tetraphenylphosphonium Bromide) in EARP to remove technetium-99 from the historic stock of MAC.

### Monitoring of the marine environment

Regular monitoring of the marine environment near to Sellafield and further afield was conducted during 2015, by the Environment Agency and FSA (for England and Wales), NIEA (for Northern Ireland) and SEPA (for Scotland). The monitoring locations for seafood, water, environmental materials and dose rates near the Sellafield site are shown in Figures 2.10 and 2.11. The medium-term trends in discharges, environmental concentrations and dose were considered in a RIFE summary report, and overall showed a decrease in concentrations over time reflecting reduced discharges at Sellafield (Environment Agency, FSA, NIEA and SEPA, 2010b). The Environment Agency has proposed changes to its future monitoring programme that are due to take effect in 2016 and will be reflected in future RIFE reports.



**Figure 2.9.** Technetium-99 in UK seaweed (*Fucus vesiculosus*) from Sellafield liquid discharges between 1989-2015 (Note different scales used for Ardglass, Dunreay, Port William and Sellafield)



**Figure 2.10.** Technetium-99 in seaweed (*fucus vesiculosus*) from Sellafield liquid discharges between, 2004-2015

## Monitoring of fish and shellfish

Concentrations of beta/gamma activity in fish from the Irish Sea and from further afield are given in Table 2.5. Data are listed by location of sampling or landing point, north to south in Cumbria, then in approximate order of increasing distance from Sellafield. Results are available for previous specific surveys in the 'Sellafield Coastal Area' (extending 15 km to the north and to the south of Sellafield, from St Bees Head to Selker, and 11 km offshore) and the smaller 'Sellafield Offshore Area' (consisting of a rectangle, 1.8 km wide by 3.6 km long, situated south of the pipelines) in earlier RIFE reports (e.g. Environment Agency, FSA, NIEA, NRW and SEPA, 2014). Concentrations of specific naturally occurring radionuclides in fish and shellfish in the Sellafield area are given in Section 7.


The concentrations of most radionuclides have decreased over the previous decades in response to decreases in discharges. Concentrations generally continue to reflect changes in discharges over time periods, characteristic of radionuclide mobility and organism uptake. Trends in concentrations of radionuclides, and corresponding discharge levels, in seafood near Sellafield (over the last decade) are shown in Figures 2.13 – 2.18. There was variability from year to year, particularly for the more mobile radionuclides. Liquid discharges of technetium-99 in 2015 were similar to those in 2014. Overall, concentrations of technetium-99 in fish and shellfish have shown a continued reduction, from the relatively elevated levels shown at the beginning of the reported period, but were generally similar (with minor variations) over most recent years (Figure 2.15). For the transuranic elements (Figures 2.17 – 2.18), the long-term trends of reductions in concentrations from earlier decades appear to be slowing. Over the last decade, despite generally decreasing discharges, concentrations of americium-241 and plutonium-239+240 in fish and shellfish have shown some variations from year to year. Overall, concentrations of plutonium radionuclides and americium-241 in winkles

were generally similar (with minor variations) in 2015 compared to those in recent years.

Beta/gamma-emitting radionuclides detected in fish included: tritium, carbon-14, strontium-90 and caesium-137 (Table 2.5). Overall, concentrations of caesium-137 in fish species were generally similar in comparison to those in 2014. Over the longer time period, activity concentrations in fish and shellfish appear to be generally declining (with minor variations) at a slow rate (Figure 2.16). Activity concentrations in fish (and shellfish) generally reflected progressive dilution with increasing distance from Sellafield. However, the rate of decline of caesium-137 concentrations with distance was not as marked as was the case when significant reductions in discharges were achieved some years ago. There was therefore a greater contribution from historical sources.

Up until 2013, brown trout was sampled for analysis from the River Calder, which flows through the Sellafield site. Results for previous measured caesium-137 concentrations, and long-term trend information, are available in earlier RIFE reports (e.g. Environment Agency, FSA, NIEA, NRW and SEPA, 2014). The changes in concentrations were likely to be due the combined effects of Sellafield discharges and fallout from Chernobyl, accentuated by the movement of such fish in the Calder river system.

Other artificial beta/gamma-emitting radionuclides detected in fish included carbon-14 and tritium. With an expected carbon-14 concentration from natural sources  $\sim 25$  Bq kg<sup>-1</sup>, the data suggest a continued local enhancement of carbon-14 due to discharges from Sellafield. In 2015, carbon-14 provided the highest activity concentration in marine fish (plaice, 120 Bq kg<sup>-1</sup>), with lower concentrations of tritium (total) and Organically Bound Tritium (OBT). In previous years, virtually all of the total tritium was associated with organic matter. In 2015, the limited tritium results suggest that the total tritium in marine samples was a combination of OBT (associated with organic matter) and tritiated water. Nevertheless, due to



**Figure 2.11.** Monitoring locations in Cumbria, 2015 (not including farms)

the low toxicity of this isotope of hydrogen and the low concentrations observed, the overall dose implication was very small.

For shellfish, a wide range of radionuclides is detectable, owing to generally greater uptake of radioactivity by these organisms from sediments. Generally, molluscs tend to contain higher concentrations than crustaceans and both contain higher concentrations than fish. Concentrations of beta/gamma-emitting radionuclides are shown in Table 2.6 (Table 2.7 for plutonium-241). There can be substantial

variations between species; for example, lobsters tend to concentrate more technetium-99 than crabs (see also Knowles *et al.*, 1998; Swift and Nicholson, 2001). The highest concentrations from Sellafield discharges were of tritium, carbon-14, and technetium-99. Comparing 2015 and 2014 data across a wide range of sampling locations and shellfish species (where comparisons can be made), technetium-99 concentrations were similar (with minor variations), but reduced in comparison to those years prior to 2012 due to the progressive reductions in discharges of this radionuclide. Low concentrations of ruthenium-106







Figure 2.12. Monitoring locations at Sellfield, 2015 (not including farms)




**Figure 2.13.** Carbon-14 liquid discharge from Sellafield and concentrations in plaice, lobsters and winkles near Sellafield, 2004-2015




**Figure 2.14.** Cobalt-60 liquid discharge from Sellafield and concentrations in plaice, lobsters and winkles near Sellafield, 2004-2015




**Figure 2.15.** Technetium-99 liquid discharge from Sellafield and concentrations in plaice, lobsters and winkles near Sellafield, 2004-2015



**Figure 2.16.** Caesium-137 liquid discharge from Sellafield and concentrations in plaice, lobsters and winkles near Sellafield, 2004-2015



**Figure 2.17.** Plutonium-239+240 liquid discharge from Sellafield and concentrations in plaice, lobsters and winkles near Sellafield, 2004-2015



**Figure 2.18.** Americium-241 liquid discharge from Sellafield and concentrations in plaice, lobsters and winkles near Sellafield, 2004-2015

were positively detected in mussels in 2015. Concentrations of other radionuclides (non-transuranic) in 2015 were also broadly similar (where comparisons can be made) to those in 2014.

Transuranic radionuclide data for fish and shellfish samples (chosen on the basis of potential radiological significance) in 2015 are given in Table 2.7. Transuranic elements are less mobile than other radionuclides in seawater and have a high affinity for sediments; this is reflected in higher concentrations of transuranic elements in shellfish compared with fish. Comparing 2015 and 2014 data across a wide range of sampling locations and shellfish species further afield from Sellafield, concentrations in shellfish were generally similar (where comparisons can be made). Those from the north-eastern Irish Sea were the highest transuranic concentrations found in foodstuffs in the UK. In comparison to 2014 data, the concentrations in shellfish were generally similar for plutonium radionuclides and americium-241 at most of the north-eastern Irish Sea locations in 2013, with a small increase in activity concentrations in winkles (from Parton and Nethertown). Variations of these observations in previous years were likely to have resulted from a combination of mechanisms including natural environmental variability and redistribution of sediments due to natural processes.

## Monitoring of sediments

Radionuclides in Sellafield liquid discharges are taken up into sediments along the Cumbrian Coast in particular in muddier (fine grained) areas such as estuaries. Some of these areas are used by the public. Levels of radionuclides are regularly monitored, both because of their relevance to exposure and in order to keep distributions of radioactivity under review. The results for 2015 are shown in Table 2.8. Radionuclides detected include cobalt-60, strontium-90, caesium-137 and transuranic elements. The highest concentrations found are close to the site and in fine particulate materials in estuaries and harbours, rather than the coarser-grained sands on open beaches. The concentrations of long-lived radionuclides, particularly caesium-137 and the transuranic elements, largely reflect past discharges from Sellafield, which were considerably higher than in recent years. Over the last 30 years discharges have fallen significantly as the site provided enhanced treatment to remove radionuclides prior to discharge. Overall, concentrations in sediments were similar in 2015, although small enhancements of plutonium radionuclides and americium-241 were measured from Maryport Harbour, in comparison to those in recent years.

The trends over time (1989 – 2015) for activity concentrations in mud from Ravenglass with discharges from Sellafield are shown in Figures 2.19 – 2.22. The concentrations of most radionuclides have decreased over the past 25 years in response to decreases in discharges, with sustained reductions in discharges of caesium-137


and transuranic elements. Discharges of cobalt-60 have been variable in the earlier years but reduced over the last decade, as reflected in the sediment concentrations at Ravenglass, with some evidence of a lag time between discharge and sediment concentration (Figure 2.21). The cobalt-60 concentration in mud from Ravenglass is the lowest reported value in 2015, following the similar downward trend as recent years. Over the last decade, caesium-137 and transuranic concentrations in sediments have remained relatively constant (Figures 2.19, 2.20 and 2.22). Since the mid-1990s, discharges of caesium-137, plutonium isotopes and americium-241 have remained at low levels, but with some variability. There is as a suggestion of small progressive increases in caesium-137 and transuranic elements activities in sediments (peaking in 2006), and transuranic elements possibly increasing again over the most recent 5 years. The likely explanation is that changes in these concentrations are due to remobilisation and subsequent accretion of fine-grained sediments containing higher activity concentrations. For americium-241, there is also an additional contribution due to radioactive in-growth from the parent plutonium-241 already present in the environment. The effect is less apparent in fish and shellfish (Figures 2.16 – 2.18) and will continue to be monitored.

Concentrations of caesium-137 and americium-241 in sediments from coastal locations of the north-east Irish Sea are also shown in Figure 2.23. Concentrations of both radionuclides diminish with distance from Sellafield. Overall, concentrations in 2015 at a given location were generally similar to those in 2014, and any fluctuations were most likely due to the normal variability expected to be in the environment. Limited evidence suggests that small peaks in activity concentrations have occurred in sediments at some locations at distance from Sellafield in recent years, but these are still below peak values reported over the whole period of time (except at Carslith). The effect appears to be more pronounced for americium-241 and is likely to be due to the spreading of activity away from Sellafield combined with the effect of grow-in from plutonium-241 (Hunt *et al.*, 2013).


A research study, commissioned by FSA, determined the depth distributions of technetium-99 concentrations in sea-bed cores to produce an estimate of the total inventory residing in the sub-tidal sediments of the Irish Sea (Jenkinson *et al.*, 2014). The study concluded that the inventory of technetium-99 was estimated to have been of the order of 30 TBq (or approximately 2 per cent of the total cumulative Sellafield discharge), with approximately 8 TBq present in surface material and thereby potentially most susceptible to re-dissolution or re-suspension.

## Monitoring of dose rates


Dose rates are regularly monitored at a large number of locations, both in the Sellafield vicinity and further afield, using environmental radiation dosimeters. Table 2.9 lists




**Figure 2.19.** Caesium-137 liquid discharge from Sellafield and concentration in mud at Ravenglass, 1988-2015



**Figure 2.20.** Plutonium-alpha liquid discharge from Sellafield and plutonium-239+240 concentration in mud at Ravenglass, 1988-2015



**Figure 2.21.** Cobalt-60 liquid discharge from Sellafield and concentration in mud at Ravenglass, 1988-2015



**Figure 2.22.** Americium-241 liquid discharge from Sellafield and concentration in mud at Ravenglass, 1988-2015

the locations monitored by the environment agencies and the gamma dose rates in air at 1 m above ground. Where comparisons can be made from similar ground types and locations, dose rates over intertidal areas throughout the Irish Sea in 2015 were generally similar to those in recent years. Any variations between years are likely to have been due to normal variability expected to be present in the environment. As in previous years, gamma dose rates measured on the banks of the River Calder, which flows through the Sellafield site, continued to show significant excess above natural background downstream of the site (of approximately  $0.04 \mu\text{Gy h}^{-1}$ ). Although the dose rates were locally enhanced, occupancy by the public, mainly anglers, is low in this area (unlikely to be more than a few tens of hours per year). On this basis the resulting doses were much less than those at other intertidal areas as discussed earlier in this section.

Gamma dose rates above mud and salt marshes, from a range of coastal locations in the vicinity of Sellafield, are shown in Figure 2.24. Gamma dose rates at sandy locations are generally lower than those above mud or salt marshes. The general decrease in dose rates with increasing distance from Sellafield, which was apparent under conditions of higher discharges several decades ago, is no longer so prominent in recent years. Spatial variability of dose rates is expected, depending on ground type; generally higher dose rates being recorded over areas with finely divided sediments. For each location, there has been variation over time. Close to Sellafield (at Carleton Marsh and Newbiggin), there was limited evidence to suggest that dose rates were slowly declining over the whole period. Locations that are further afield from Sellafield show dose rate values that only marginally exceeded average UK natural background rates.


Over the last 30 years, levels of radioactivity in the environment around Sellafield have declined as a result of reduced discharges. In more recent years the levels in the Esk estuary have shown a less clear trend, with concentrations of some radionuclides fluctuating from

year to year (for example, see Figure 2.20). This effect could be due to the dynamic nature of the sediment in the estuary, which is eroded and transported by tide and freshwater, periodically exposing older deeper sediment containing radioactivity from historical discharges. Due to the variations seen in recent years and local concerns, the Environment Agency initiated a more detailed study of dose rates in the Esk Estuary in 2007. The objectives of the study were to assess the current level of external gamma radiation exposure in the estuary, and changes in the measured dose rates, relative to a more detailed survey of the estuary undertaken in 1989 (Kelly and Emptage, 1991). A six-week survey of gamma dose rates was undertaken at a total of 576 locations in the Esk Estuary. The University of Liverpool (Institute for Sustainable Water Integrated Management and Ecosystem Research (SWIMMER)) undertook the study.


The mean dose rate across all 576 locations was  $0.14 \mu\text{Gy h}^{-1}$ , with a range of  $0.07 - 0.28 \mu\text{Gy h}^{-1}$ . This indicates a significant decrease compared to the mean dose rate reported in 1989 (at similar locations) of  $0.23 \mu\text{Gy h}^{-1}$  (range  $0.07 - 0.61 \mu\text{Gy h}^{-1}$ ). The highest gamma dose rates measured in both surveys were from comparable locations within the estuary. The reduced dose rates in the 2007 survey were due to the effects of reductions in radionuclide discharges from the Sellafield site and also radioactive decay of the inventory within the Esk Estuary sediments and soils since 1989. The full report on this study has been published by the Environment Agency (Wood *et al.*, 2011).

## Monitoring of fishing gear

During immersion in seawater, fishing gear may entrain particles of sediment on which radioactivity is adsorbed. Fishermen handling this gear may be exposed to external radiation, mainly to skin from beta particles. Fishing gear is regularly monitored using surface contamination monitors. Results for 2015 are given in Table 2.10. Overall, where



**Figure 2.23.** Concentrations of americium-241 and caesium-137 in coastal sediments in North West England and South West Scotland between 2003-2015 (Note different scales used for Newbiggin and Carleton Marsh)



**Figure 2.24.** Gamma dose rates above fine coastal sediments (mud and salt marshes) in North West England and South West Scotland between 2003-2015

comparisons can be made, measured dose rates were generally similar to those in 2014.

## Contact dose-rate monitoring of intertidal areas

Results from measurements of beta dose rates on shoreline sediments (using contamination monitors), to allow estimation of exposure of people who handle sediments regularly, are given in Table 2.11. Overall, positively detected dose rates were generally higher in 2015, in comparison to those in recent years (where comparisons can be made from similar ground types and locations). Beta dose rates were higher in sand at Whitehaven (outer harbour), St Bees, Sellafield pipeline and Tarn Bay, and in salt marsh at Ravenglass, in 2015 (compared to 2014).

More general beta/gamma monitoring for the Environment Agency of contamination on beaches using portable probes continued to establish whether there are any localised 'hot spots' of activity, particularly in strand lines and beach debris. In 2015, no material was found using these probes in excess of the action level equivalent to 0.01 mSv h<sup>-1</sup>.

In 2008, the Environment Agency published a formal programme of work for the assessment of contamination by radioactive particles on and around the west Cumbrian coastline. The assessment was focused on public protection from high activity discrete radioactive particles that have been released to the environment from activities at the Sellafield site (Environment Agency, 2008c). The work so far has included investigating the distribution and behaviour of Sellafield-related particles, particle analysis and identification, risks from particles, and a review of particle dispersion and transport models focused on the Eastern Irish Sea and Solway Firth.

Since vehicle-mounted beach survey work began in November 2006, and up to the end of March 2015, approximately 2,000 hectares of beach area has been surveyed by the Sellafield site operator's contractors (Sellafield Limited, 2015). The survey equipment used currently (since August 2009) is the Groundhog™ Synergy system, which is an improvement on the use of the original Groundhog™ Evolution system. The Groundhog™ Synergy system has a specific capability in relation to the detection of medium/high energy gamma emitting radionuclides and also provides improved detection capability for low energy gamma emissions, increasing the ability to detect particles containing americium-241.

During 2015/16, further beach monitoring was completed in line with the Environment Agency's specification of approximately 166 hectares and exceeded the Environment Agency's specification of 160 hectares (Sellafield Limited, 2016). The number of radioactive finds identified in the period from April 2015 to March 2016 was 349 (compared with 383 in the previous year), of which 83 per cent

were classified as particles (less than 2 mm in size) and the remainder as stones (larger than 2 mm in size). The number of finds were typical of those in recent years. The majority of the finds were concentrated on a 5 km stretch of beach running NW from the Sellafield site. All have been removed from the beaches.

Monitoring along the Cumbrian coast will continue, with the current proposal being a further 150 hectares to be surveyed between April 2016 and March 2017, as part of the operator's routine environmental monitoring programme. A programme of work is in place to address the remaining uncertainties in the origins, fate and effects of the particles with the aim of returning the work to a routine monitoring programme only by about 2018.

In 2012, PHE reported their review of the results and position on risk following the introduction of the improved Synergy™ monitoring system. The report concluded that the increase in particle finds following the introduction of this system was a result of its improved capability and also that advice previously given by PHE to the Environment Agency following a detailed assessment of risks in 2010 remained valid (Brown and Etherington, 2011; Etherington *et al.*, 2012). The report restated the conclusion that based on the currently available information, the overall health risks to beach users are very low and significantly lower than other risks people accept when using the beaches. As such, PHE advice remained that no special precautionary actions were required to limit access to or use of the beaches. A more recent report by PHE describes the assessed health risks from the consumption of seafood (including those to commercial fishermen) from radioactive particles in the vicinity of the Sellafield Site (Oatway and Brown, 2015). Based on currently available information, it is concluded that the overall health risks to both seafood consumers and commercial fishermen are very low.

In June 2014, a single particle was found which was unusual in that its activity was predominantly from beta radiation (strontium-90). PHE has undertaken detailed dose measurements of this particle and established that the skin dose rate from this one particle exceeds the criterion recommended to prompt an urgent review of the risk assessment. PHE has provided advice that "While the retrieval of one particle cannot in itself be regarded as a substantial public health issue, this find should now lead to a reassessment of monitoring capabilities for particles with [a] high content of strontium-90, along with any other measures that may be considered appropriate, to support a reappraisal of the potential health risks."

In August 2015, a beta detection trial was carried out (using hand held detectors, allowing monitoring of areas inaccessible to vehicles) to investigate additional capability for the detection of strontium-90 finds on the beaches at Sellafield. Approximately 1.7 hectares were surveyed and 7 finds were detected (and removed) in the trial. None of the finds were found by the beta detector and none were strontium-90 rich particles. These results, along with

a re-evaluation of the detection limits for the Synergy™ monitoring system, has enabled Sellafield Limited to conclude that there is not a significant population of undetected strontium-90 rich particles on the beaches.

In relation to food safety, and following a previous assessment of the particles frequency and the activity concentrations, FSA's guidance to the Environment Agency supported PHE's advice. The Environment Agency will also continue to work with relevant authorities to keep the situation under review.

In 2007, SEPA published a strategy document for the assessment of the potential impact of Sellafield radioactive particles on members of the public in south-west Scotland (SEPA, 2007) and the beach monitoring programme was temporarily extended to include two locations on the north Solway coastline (Kirkcudbright Bay and Southernness). This was based on some limited modelling work on the movement of particles undertaken for the Environment Agency following a request by SEPA. No particles were detected at these locations. SEPA is maintaining a watching brief on the situation in as much as it may affect Scotland.

Between 2010 and 2013, the Environment Agency provided updates on further progress of the enhanced beach monitoring (Environment Agency, 2010; 2011; 2013c) with work prior to 2010 described elsewhere (Environment Agency, FSA, NIEA and SEPA, 2010a).

Further detail on enhanced beach monitoring data compiled so far can be obtained from Sellafield Limited website: <http://sustainability.sellafieldsites.com/environment/environment-page/particles-in-the-environment/>

## Monitoring of seaweed

In addition to occasional use in foods and as fertilisers, seaweeds are useful environmental indicator materials (as radionuclides are concentrated by seaweeds), facilitating assessments and assisting the tracing of these radionuclides in the environment. Table 2.12 gives the results of measurements in 2015 of seaweeds from shorelines of the Cumbrian coast and further afield.

*Fucus* seaweeds are particularly useful indicators of most fission product radionuclides: samples of *Fucus vesiculosus* are collected both in the Sellafield vicinity and further afield to show the extent of Sellafield contamination in north European waters. The effects of technetium-99 discharges from Sellafield on concentrations in seaweed, between 1989 and 2015, are shown in Figure 2.9. In the north-east Irish Sea, technetium-99 concentrations have been reasonably constant over the present decade, consistent with the relatively low discharges; the highest concentrations which were found near Sellafield were much less than those in the mid 1990's and the decade thereafter (in response to the progressive reduction in

discharges). In general, there was also a large reduction in concentrations of technetium-99 in *Fucus vesiculosus* with distance from Sellafield, as the effect of the discharges becomes diluted in moving further afield.

Technetium-99 concentrations in seaweed (Table 2.12) collected from sites in Cumbria were generally higher (by small amounts) in comparison to those in 2014. At one specific location (Auchencairn, Scotland), known to have had fluctuating concentrations in previous years, activity concentrations in seaweed (*Fucus*) were also higher in 2015 compared with those in 2014. Variations in concentrations in the past were most likely the result of complex hydrographic transport patterns in the Irish Sea, with technetium-99 being dispersed to a variable degree before arriving at distant locations (Leonard *et al.*, 2004). It may also be noted that as the effects of the high technetium discharges of the 1990s continue to disperse, there is the potential for areas distant from Sellafield to exhibit concentrations greater than those in closer proximity, such as Auchencairn, and as observed in seawater in Liverpool Bay for 1998 (McCubbin *et al.*, 2002).

Seaweeds are sometimes used as fertilisers and soil conditioners and this potential pathway for the transfer of radionuclides into the food chain has been previously monitored to determine the activity concentrations in a range of vegetables, and in grass and soil. Results are available in earlier RIFE reports (e.g. Environment Agency, FSA, NIEA, NRW and SEPA, 2014).

No harvesting of *Porphyra* in west Cumbria, for consumption in the form of laverbread, was reported in 2015; this pathway has therefore remained dormant. Samples of *Porphyra* were regularly collected from selected locations along UK shorelines of the Irish Sea in previous years. Results from surveys providing activity concentrations in *Porphyra*, and in samples of the major manufacturers' laverbread (that were regularly collected from markets in South Wales), are available in earlier RIFE reports (e.g. Environment Agency, FSA, NIEA, NRW and SEPA, 2014).

## Monitoring of seawashed pasture

The potential transfer of technetium-99 to milk, meat and offal from animals grazing tide-washed pasture was considered using a modelling approach in the report for 1997 (MAFF and SEPA, 1998). The maximum potential dose was calculated to be 0.009 mSv at that time. Follow-up sampling of tide-washed pastures at Newton Arlosh, Cumbria and Hutton Marsh, Lancashire in 2006 suggested that this dose estimate remains valid (Environment Agency, Environment and Heritage Service, FSA and SEPA, 2007).

## Monitoring of sea to land transfer

Terrestrial foodstuffs are monitored near Ravenglass to check on the extent of transfer of radionuclides from sea to land in this area. In 2015, samples of milk and livestock were collected and analysed, for radionuclides which were released in liquid effluent discharges from Sellafield. Results from surveys for activity concentrations in crops, fruit and environmental indicators are available in earlier RIFE reports (e.g. Environment Agency, FSA, NIEA, NRW and SEPA, 2014).

The results of measurements in 2015 are given in Table 2.13. Generally, the activity concentrations, where positively detected, showed lower concentrations than were found in the immediate vicinity of Sellafield (Table 2.4). As in previous years, the evidence for sea to land transfer was very limited in 2015. Technetium-99 and plutonium-238 concentrations are reported as less than values (or close to the less than value). Small concentrations of artificial nuclides were detected in some samples but the concentrations were very low. In recent years, where detectable, observed isotopic ratios of  $^{238}\text{Pu}:^{239+240}\text{Pu}$  concentrations were somewhat higher than 0.025, a value which might be expected if the source was only (or entirely) due to fallout. This may suggest a Sellafield influence.

## Monitoring of fishmeal

Low concentrations of man-made radioactivity were found in fishmeal, which is fed to farmed fish, poultry, pigs, cows and sheep. A theoretical study has established that any indirect onward transmission of radioactivity into human diet as a result of this pathway is unlikely to be of radiological significance (Smith and Jeffs, 1999). A detailed survey was undertaken in 2003 to confirm these findings. Samples were obtained from 14 fish farms in Scotland and three in Northern Ireland. They demonstrated that concentrations of radionuclides are indeed very low, most being less than the limits of detection, and the few that were positively determined were all less than  $1 \text{ Bq kg}^{-1}$  (FSA, 2003). Previous reported results (published in previous RIFE reports, Tables 2.5 and 2.7) for activity concentrations in farmed salmon from the west of Scotland confirm the findings of the 2003 study.

## Monitoring of waters

Evidence of the effects of liquid discharges from Sellafield on concentrations of radionuclides in seawater is determined by sampling from research vessels and the shore. The results of the seawater programme are given in Section 8.

Sampling of fresh water from rivers and lakes in west Cumbria is conducted as part of the regular environmental monitoring programme around Sellafield; however, other

environmental materials are likely to be more indicative of direct site-related effects. Some of the sources monitored provide public drinking water. The results for 2015 are included in Table 2.14. Tritium, gross alpha and gross beta concentrations in public supplies were below the investigation levels for drinking water in the European Directive 2013/51.

Small amounts of activity are discharged from Sellafield under permit via the factory sewer outfall to the River Ehen Estuary, immediately prior to the confluence with the River Calder. As in recent years, there was no evidence of tritium 100m downstream of the outfall in 2015 (Table 2.14). These waters are not potable and any low concentrations observed previously are of no radiological significance. Table 2.14 also includes the results of monitoring from the Ehen Spit (Figure 2.11) near Sellafield where water issues from the ground at low tide. This release is not due to regulated discharges of liquid wastes but to ground water migration from the Sellafield site. The water is brackish so it will not be used as a drinking water source and therefore the only consumption would be inadvertent. Enhanced gross beta and tritium concentrations were observed in 2015 with concentrations similar to those in recent years. The dose from inadvertent consumption of water from Ehen Spit has been shown to be insignificant (Environment Agency, 2002a).

## 2.3.4 Monitoring of unusual pathways

In 1998, high concentrations of caesium-137 (of up to  $110,000 \text{ Bq kg}^{-1}$ ) were found in feral pigeons sampled in Seascale by MAFF. Consumption of the breast meat of only 20 birds contaminated at the highest concentration would have given a dose of 1 mSv to high-rate consumers. Advice issued by MAFF in 1998 was that people should not handle, slaughter or consume pigeons within a radius of 10 miles of the site. A full review of the incident was published in 1999 (Copeland Borough Council *et al.*, 1999). It was found that pigeons had access to the roof spaces in buildings on the Sellafield site and had become contaminated with radionuclides including caesium-137. The pigeons were also congregating in large numbers at a bird sanctuary in Seascale village and the environment around had become contaminated. Since then, the site operator has undertaken remedial measures, including a substantial cull of feral pigeons in the area and preventing access to the loft spaces in buildings on the Sellafield site. Results of the analysis of a wood pigeon sample collected in 2015 are included in Table 2.4. The maximum caesium-137 concentration in the muscle of wood pigeon ( $0.09 \text{ Bq kg}^{-1}$ ) in 2015 was detected just above the less than value, and generally similar to the maximum value reported in 2014 ( $0.27 \text{ Bq kg}^{-1}$ ). These radiocaesium concentrations have had fluctuating levels in recent years prior to 2011. Concentrations of artificial radionuclides were low and would add little to the exposure of local consumers. The FSA will continue to monitor this pathway.

Following the review of the pigeon incident, the Environment Agency began to sample and analyse sediments from road drains (gully pots) in Seascale and Whitehaven in 1999. Gully pots in road drains collect sediments washed off road surfaces and provide good indicators of contamination of urban environments. The results of analyses in 2015 are shown in Table 2.15, and were generally similar to those in recent years. In 2010, elevated concentrations (of strontium-90, caesium-137, americium-241 and plutonium radionuclides) in sediments were reported for one of the five Seascale road drains (Seascale SS 233). Investigations, including monitoring of additional Seascale road drains, were conducted in 2011 to confirm that the elevation had ceased or to inform appropriate action. The results indicate that the elevated levels in 2010 were not sustained during the period 2011 to 2013, and that these results were mostly consistent with other road drains sampled. The enhancements may have arisen from unusual weather conditions in that year, releasing radioactivity trapped within the drainage path. Generally, over a longer period, activity concentrations in road drains have fallen significantly since remedial measures were taken to reduce contamination.

## 2.4 Windscale, Cumbria



Windscale was historically a separate licensed site located at Sellafield. NDA has ownership of the site. In 2008, the Windscale permit was transferred from UKAEA to Sellafield Limited, and combined with

the Sellafield site permit. At Windscale there are three nuclear reactors, two of which were shut down in 1957 and the third in 1981. Most of the radioactive wastes derive from decontamination and decommissioning operations, some of which are of the early Windscale reactor buildings. Decommissioning activities began in the mid-1980s and these activities are continuing. The reactor decommissioning of the Windscale Advanced Gas Cooled Reactor (AGR) was completed in 2011. The current plan is to achieve complete decommissioning of Windscale by 2050 (NDA, 2016a). Gaseous wastes are regulated from specific stacks on the Windscale site; liquid radioactive wastes are disposed of, after appropriate treatment, to the Irish Sea via the Sellafield site pipelines. Both gaseous and liquid discharges are included as part of the regulated Sellafield discharges (Appendix 2). Discharges of both gaseous and liquid radioactive wastes are minor compared to those from the Sellafield nuclear licensed site.

Regular monitoring of the environment by the Environment Agency and FSA in relation to any releases from the Windscale site is conducted as part of the overall programme for the Sellafield site. The results of this monitoring and the implications in terms of dose to people in Cumbria are described in Section 2.3.

**Table 2.1. Individual doses – Capenhurst and Springfields, 2015**

Site	Representative person <sup>a</sup>	Exposure, mSv per year						
		All pathways	Seafood	Other local food	External radiation from intertidal areas, river banks or fishing gear	Intakes of sediment and water	Gaseous plume related pathways	Direct radiation from site
Capenhurst								
<b>Total dose – all sources</b>	<b>Local adult inhabitants (0 - 0.25km)</b>	<b>0.13<sup>c</sup></b>	-	<b>&lt;0.005</b>	-	-	<b>&lt;0.005</b>	<b>0.13</b>
Source	Infant inhabitants and consumers of locally grown food	<0.005 <sup>c</sup>	-	<0.005	-	-	<0.005	-
specific doses	Children playing at Rivacre Brook	0.010 <sup>c</sup>	-	-	0.009	<0.005	-	-
Springfields								
<b>Total dose – all sources</b>	<b>Adult occupants on houseboats</b>	<b>0.050</b>	-	-	<b>0.050</b>	-	-	-
Source	Seafood consumers	0.020 <sup>c</sup>	<0.005	-	0.019	-	-	-
specific doses	Houseboat occupants	0.054	-	-	0.054	-	-	-
	Children playing at Lower Penwortham <sup>c</sup>	<0.005	-	-	<0.005	<0.005	-	-
	External in intertidal areas (farmers)	0.034	-	-	0.034	-	-	-
	Wildfowl consumer	0.006 <sup>c</sup>	-	<0.005	0.005	-	-	-
	Inhabitants and consumers of locally grown food	<0.005 <sup>c</sup>	-	<0.005	-	-	<0.005	-

<sup>a</sup> The total dose is the dose which accounts for all sources including gaseous and liquid discharges and direct radiation. The total dose for the representative person with the highest dose presented. Other dose values are presented for specific sources, either liquid discharges or gaseous discharges, and their associated pathways. They serve as a check on the validity of the total dose assessment. The representative person is an adult unless otherwise stated

<sup>b</sup> Exposure to skin for comparison with the 50 mSv dose limit

<sup>c</sup> Includes a component due to natural sources of radionuclides

**Table 2.2(a). Concentrations of radionuclides in food and the environment near Capenhurst, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>							
			<sup>3</sup> H	<sup>99</sup> Tc	<sup>137</sup> Cs	<sup>234</sup> Th	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U	<sup>237</sup> Np
<b>Marine samples</b>										
Flounder	Liverpool Bay	1	<25		1.5					
Shrimps	Wirral	1	<25	1.1	0.78					
Mussels	Liverpool Bay	1	<25		0.95					
Cockles	Dee Estuary	1	<25	1.5	0.90	5.6				
Sediment	Rivacre Brook	2 <sup>E</sup>		92	<1.2	44	73	2.9	41	<1.0
Sediment	Rivacre Brook (1.5 km downstream)	2 <sup>E</sup>		30	1.4	20	31	<1.5	21	<2.0
Sediment	Rossmore (3.1 km downstream)	2 <sup>E</sup>		22	<1.0	<13	23	<1.4	17	<1.0
Sediment	Rivacre Brook (4.3 km downstream)	2 <sup>E</sup>		13	0.62	<7.0	11	<0.91	8.5	<1.0
Freshwater	Rivacre Brook	2 <sup>E</sup>	<3.9	<0.30			0.034	<0.0015	0.016	<0.037
Freshwater	Rivacre Brook (1.5 km downstream)	2 <sup>E</sup>	<3.0	<0.30			0.025	<0.0017	0.014	<0.021
Freshwater	Rossmore (3.1 km downstream)	2 <sup>E</sup>	<3.1	<0.20			0.022	<0.0014	0.014	<0.021
Freshwater	Rivacre Brook (4.3 km downstream)	2 <sup>E</sup>	<3.2	<0.30			0.023	<0.0019	0.013	<0.021
<hr/>										
Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>							
			<sup>238</sup> Pu	<sup>239</sup> Pu+ <sup>240</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm+ <sup>244</sup> Cm	Gross alpha	Gross beta	
<b>Marine samples</b>										
Flounder	Liverpool Bay	1			<0.06					
Shrimps	Wirral	1	0.0020	0.013	0.015	*	*			
Mussels	Liverpool Bay	1			1.2					
Cockles	Dee Estuary	1	0.067	0.45	1.3	*	*			
Sediment	Rivacre Brook	2 <sup>E</sup>						300	970	
Sediment	Rivacre Brook (1.5 km downstream)	2 <sup>E</sup>						<110	450	
Sediment	Rossmore (3.1 km downstream)	2 <sup>E</sup>						<110	530	
Sediment	Rivacre Brook (4.3 km downstream)	2 <sup>E</sup>						<120	420	
Freshwater	Rivacre Brook	2 <sup>E</sup>						<0.088	0.51	
Freshwater	Rivacre Brook (1.5 km downstream)	2 <sup>E</sup>						<0.036	0.37	
Freshwater	Rossmore (3.1 km downstream)	2 <sup>E</sup>						<0.035	0.29	
Freshwater	Rivacre Brook (4.3 km downstream)	2 <sup>E</sup>						<0.039	0.16	

Table 2.2(a). continued

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>d</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			<sup>3</sup> H <sup>c</sup>	<sup>99</sup> Tc	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U
<b>Terrestrial samples</b>							
Milk		2	<3.0	<0.038	<0.0022	<0.00079	<0.0010
Milk	max			<0.041		<0.00094	<0.0013
Beetroot		1		<0.22	0.011	<0.00043	0.010
Grass		1		<0.24	0.018	0.00080	0.018
Grass/herbage	North of Ledsham	1 <sup>E</sup>		<0.72	<0.48	<0.18	0.46
Grass/herbage	South of Capenhurst	1 <sup>E</sup>		<0.96	<0.19	<0.095	<0.20
Grass/herbage	Dunkirk Lane (0.9 km South of Site)	1 <sup>E</sup>		<1.6	0.29	<0.065	<0.22
Grass/herbage	East of station	1 <sup>E</sup>		<1.4	0.17	<0.048	0.22
Soil	North of Ledsham	1 <sup>E</sup>		7.8	21	<1.6	21
Soil	South of Capenhurst	1 <sup>E</sup>		7.2	21	<2.0	21
Soil	Dunkirk Lane (0.9 km South of Site)	1 <sup>E</sup>		<2.1	20	<1.2	20
Soil	East of station	1 <sup>E</sup>		<2.8	19	1.2	20

\* Not detected by the method used

<sup>a</sup> Except for milk and water where units are Bq l<sup>-1</sup>, and for soil and sediment where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'Max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> In distillate fraction of sample

<sup>d</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

Table 2.2(b). Monitoring of radiation dose rates near Capenhurst, 2015

Location	Ground type	No. of sampling observations	µGy h <sup>-1</sup>
<b>Mean gamma dose rates at 1m over substrate</b>			
Rivacre Brook Plant outlet	Grass and mud	1	0.090
Rivacre Brook Plant outlet	Grass	1	0.094
Rivacre Brook 1.5 km downstream	Grass	2	0.082
Rossmore Road West 3.1 km downstream	Grass	2	0.083
Rivacre Brook 4.3 km downstream	Mud	1	0.081
Rivacre Brook 4.3 km downstream	Grass	1	0.082

**Table 2.3(a). Concentrations of radionuclides in food and the environment near Springfields, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>									
			<sup>3</sup> H	<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>99</sup> Tc	<sup>129</sup> I	<sup>137</sup> Cs	<sup>228</sup> Th	<sup>230</sup> Th	<sup>232</sup> Th
<b>Marine samples</b>												
Flounder	Ribble Estuary	1			<0.05					2.3		
Bass	Ribble Estuary	1			<0.07					4.0		
Shrimps <sup>d</sup>	Ribble Estuary	1		64	<0.07		<0.88			1.3	0.015	0.0072
Mussels <sup>d</sup>	Ribble Estuary	1			<0.06					0.97	0.21	0.15
Wildfowl	Ribble Estuary	1	<2.2	38	<0.05	<0.046		<0.90		0.85		0.011
Samphire	Marshside Sands	1			<0.05		<0.34			0.38		
Sediment	River Ribble outfall	4 <sup>E</sup>			<0.88					120	29	51
Sediment	Savick Brook	2 <sup>E</sup>			<0.92					160	33	74
Sediment	Lea Gate	2 <sup>E</sup>			<1.1					160	40	73
Sediment	Lower Penwortham Park	4 <sup>E</sup>			<1.0					140	32	78
Sediment	Penwortham rail bridge	4 <sup>E</sup>			<0.97					110	31	56
Sediment	Penwortham rail bridge - West bank	2 <sup>E</sup>			<0.50					75	29	40
Sediment	Penwortham position 1	4 <sup>E</sup>			<1.0					120	27	53
Sediment	Penwortham position 2	1 <sup>E</sup>			<0.32					9.1	19	21
Sediment	Lytham Yacht Club	1 <sup>E</sup>			<1.1					210	43	76
Sediment	Beconsall	4 <sup>E</sup>			<0.88					83	22	37
Sediment	Freckleton	1 <sup>E</sup>			<1.1					210	41	74
Sediment	Hutton Marsh	1 <sup>E</sup>			<1.7					360	42	170
Sediment	Longton Marsh	1 <sup>E</sup>			<0.95					480	53	260
Grass (washed)	Hutton Marsh	1 <sup>E</sup>					<0.55					
Grass (unwashed)	Hutton Marsh	1 <sup>E</sup>					<0.58					
Soil	Hutton Marsh	1 <sup>E</sup>					32					

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>									
			<sup>234</sup> Th	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U	<sup>237</sup> Np	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am	Gross alpha	Gross beta
<b>Marine samples</b>												
Flounder	Ribble Estuary	1									<0.05	
Bass	Ribble Estuary	1									<0.18	
Shrimps <sup>d</sup>	Ribble Estuary	1					0.000058	0.0015	0.0090	0.016		
Mussels <sup>d</sup>	Ribble Estuary	1						0.077	0.46	0.79		
Wildfowl	Ribble Estuary	1						0.0011	0.0070	0.012		
Samphire	Marshside Sands	1								0.18		
Sediment	River Ribble outfall	4 <sup>E</sup>	170	21	<1.3	23				110	470	1200
Sediment	Savick Brook	2 <sup>E</sup>	2100	25	<1.5	25				150	470	3900
Sediment	Lea Gate	2 <sup>E</sup>	1800	30	1.7	31				150	540	2100
Sediment	Lower Penwortham Park	4 <sup>E</sup>	430	28	<1.5	29				120	470	1300
Sediment	Penwortham rail bridge	4 <sup>E</sup>	<390	24	<1.8	25				87	540	1100
Sediment	Penwortham rail bridge - West bank	2 <sup>E</sup>	360	22	<1.4	25				75	470	930
Sediment	Penwortham position 1	4 <sup>E</sup>	280	23	<1.5	23				94	430	940
Sediment	Penwortham position 2	1 <sup>E</sup>	33	14	<1.1	15				6.4	130	630
Sediment	Lytham Yacht Club	1 <sup>E</sup>	220	28	1.5	28				170	600	1400
Sediment	Beconsall	4 <sup>E</sup>	<100	20	<1.3	20				78	420	840
Sediment	Freckleton	1 <sup>E</sup>	420	35	2.3	36				180	590	1300
Sediment	Hutton Marsh	1 <sup>E</sup>	53	33	<1.6	31				270	820	1400
Sediment	Longton Marsh	1 <sup>E</sup>	69	33	<2.2	34				310	1100	1500

**Table 2.3(a). continued**

Material	Location or selection <sup>a</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>					
			<sup>3</sup> H	<sup>14</sup> C	<sup>90</sup> Sr	<sup>129</sup> I	<sup>137</sup> Cs	Total Cs
<b>Terrestrial samples</b>								
Beetroot		1	<2.5	14	<0.045	<0.023	<0.05	0.019
Sediment	Deepdale Brook	2 <sup>E</sup>					<0.90	
Grass		1	<5.5	13	0.14	<0.027	0.16	0.36
Freshwater <sup>e</sup>	Ulnes Walton	1 <sup>E</sup>	<3.1				<0.23	

Material	Location or selection <sup>a</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>					
			<sup>230</sup> Th	<sup>232</sup> Th	<sup>234</sup> Th	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U
<b>Terrestrial samples</b>								
Milk		2				0.0027	<0.0010	<0.0026
Milk	max					0.0037	<0.0011	0.0041
Beetroot		1	0.013	0.0077		0.0074	0.00036	0.0093
Sediment	Deepdale Brook	2 <sup>E</sup>			62	34	1.6	34
Grass		1	0.031	0.035		0.065	0.0027	0.065
Grass	Site fence	1 <sup>E</sup>				<0.17	<0.093	<0.12
Grass	Opposite site entrance	1 <sup>E</sup>				0.25	<0.086	0.23
Grass	Opposite windmill	1 <sup>E</sup>				<0.69	<0.29	<0.66
Grass	Deepdale Brook	1 <sup>E</sup>				<0.29	<0.11	<0.22
Grass	Lea Town	1 <sup>E</sup>				<0.14	<0.066	<0.14
Grass	N of Lea Town	1 <sup>E</sup>				<0.38	<0.19	0.32
Soil	Site fence	1 <sup>E</sup>				20	<1.5	20
Soil	Opposite site entrance	1 <sup>E</sup>				94	3.7	82
Soil	Opposite windmill	1 <sup>E</sup>				110	5.7	100
Soil	Deepdale Brook	1 <sup>E</sup>				110	4.5	100
Soil	Lea Town	1 <sup>E</sup>				32	1.6	32
Soil	N of Lea Town	1 <sup>E</sup>				55	1.6	57
Freshwater	Deepdale Brook	4 <sup>E</sup>				0.33	0.015	0.33
Freshwater <sup>e</sup>	Ulnes Walton	1 <sup>E</sup>	<0.0011	<0.00070		0.070	<0.0035	0.064

Material	Location or selection <sup>a</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>					
			<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Pu	<sup>241</sup> Am	Gross alpha	Gross beta
<b>Terrestrial samples</b>								
Beetroot		1	0.000036	0.000094	<0.19	0.000080		
Sediment	Deepdale Brook	2 <sup>E</sup>					400	820
Grass		1	0.00014	0.0013	<0.30	0.0023		
Freshwater	Deepdale Brook	4 <sup>E</sup>					0.43	0.81
Freshwater <sup>e</sup>	Ulnes Walton	1 <sup>E</sup>					0.12	0.12

<sup>a</sup> Data are arithmetic means unless stated as 'max'. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>b</sup> Except for milk and freshwater where units are Bq l<sup>-1</sup> and for sediment and soil where dry concentrations apply

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>d</sup> <sup>242</sup>Cm and <sup>243+244</sup>Cm were not detected by the method used

<sup>e</sup> The concentration of <sup>228</sup>Th was <0.0040 Bq kg<sup>-1</sup>

<sup>E</sup> Measurements are made on behalf of the Food Standards Agency unless labelled "E". In that case they are made on behalf of the Environment Agency

**Table 2.3(b). Monitoring of radiation dose rates near Springfields, 2015**

Location	Material or ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Lytham Yacht Club	Salt marsh	1	0.097
Warton Mud Marsh	Salt marsh	2	0.11
Warton Mud Marsh	Salt marsh <sup>a</sup>	2	0.12
Warton Salt Marsh	Salt marsh	2	0.10
Freckleton	Salt marsh	1	0.097
Naze Point	Salt marsh	2	0.10
Banks Marsh	Salt marsh	1	0.11
Banks Marsh	Salt marsh <sup>a</sup>	1	0.12
Hesketh Bank	Salt marsh	2	0.10
Becconsall Boatyard	Mud and salt marsh	1	0.090
Becconsall Boatyard	Salt marsh	2	0.087
Becconsall Boatyard	Grass and mud	1	0.085
Longton Marsh	Salt marsh	1	0.11
Hutton Marsh	Salt marsh	1	0.12
River Ribble outfall	Mud	3	0.088
River Ribble outfall	Mud and sand	1	0.092
Savick Brook, confluence with Ribble	Salt marsh	1	0.092
Savick Brook, confluence with Ribble	Grass	1	0.094
Savick Brook, tidal limit	Salt marsh	1	0.096
Savick Brook, tidal limit	Grass and mud	1	0.091
Savick Brook, Lea Gate	Grass and mud	1	0.097
Savick Brook, Lea Gate	Grass	1	0.097
South bank opposite outfall	Salt marsh	1	0.11
Penwortham Bridge cadet hut	Mud	2	0.086
Lower Penwortham Park	Grass	4	0.078
Lower Penwortham Railway Bridge	Mud	1	0.081
Lower Penwortham Railway Bridge	Mud and sand	1	0.083
Lower Penwortham Railway Bridge	Pebbles and sand	1	0.081
Lower Penwortham Railway Bridge	Grass	1	0.080
River Darwen	Grass	4	0.085
Riverbank Angler location 1	Mud	1	0.082
Riverbank Angler location 1	Grass and sand	1	0.080
Riverbank Angler location 1	Grass	1	0.077
Riverbank Angler location 2	Sand	1	0.079
Ulnes Walton, BNFL area survey	Grass	3	0.082
<b>Mean beta dose rates</b>			
Lytham – Granny's Bay	Mud	1	0.19
Banks Marsh	Salt marsh	1	0.094
Warton Mud Marsh	Salt marsh	1	0.10
Warton Salt Marsh	Salt marsh	1	0.066

<sup>a</sup> 15cm above substrate

**Table 2.4. Concentrations of radionuclides in terrestrial food and the environment near Sellafield, 2015**

Material	Selection <sup>a</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>									
			Organic <sup>3</sup> H		<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>99</sup> Tc	<sup>106</sup> Ru	<sup>125</sup> Sb	<sup>129</sup> I	<sup>131</sup> I
Milk		9	<3.2	<3.2	17	<0.05	<0.038	<0.038	<0.44	<0.13	<0.0046	<0.0040
Milk	max		<4.1	<4.1	19	<0.06	0.054	<0.039	<0.51	<0.16	<0.0088	<0.0066
Beef kidney		1	<29	<29	30	<0.02	<0.051	<0.24	<0.14	<0.10		
Beef liver		1	<4.1	<4.1	31	<0.05	0.023	<0.23	<0.26	<0.15	<0.024	
Beef muscle		1	<2.4	<2.4	38	<0.06	0.057	<0.21	<0.48	<0.15	<0.024	
Blackberries		1	<9.4	<9.4	23	<0.05	0.79	<0.23	<0.57	<0.15	<0.020	
Cabbage		1	<2.9	<2.9	5.7	<0.07	0.086		<0.44	<0.13	<0.012	
Carrots		1	<2.0	<2.0	10	<0.07	<0.035	<0.24	<0.59	<0.12	<0.010	
Eggs		1	<2.5	<2.5	23	<0.04	<0.042		<0.62	<0.12	<0.033	
Mushrooms		1	<3.2	<3.2	14	<0.08	<0.045		<0.64	<0.21	<0.038	
Pheasant		1	<10	<10	45	<0.05	<0.043	<0.24	<0.43	<0.14	<0.018	
Potatoes		1	<4.0	<4.0	15	<0.10	<0.043		<0.71	<0.13	<0.021	
Rabbit		1	<4.7	<4.7	28	<0.09	<0.049	<0.22	<0.72	<0.21	<0.026	
Sheep muscle		2	<4.5	<4.5	32	<0.06	<0.042	<0.25	<0.69	<0.15	<0.021	
Sheep muscle	max		<6.2	<6.2	35	<0.07	<0.044	<0.26	<0.81	<0.19	<0.027	
Sheep offal		2	<6.7	<6.7	31	<0.06	<0.045	<0.41	<0.49	<0.16	<0.015	
Sheep offal	max		<9.1	<9.1	32		0.046	<0.46	<0.50		<0.016	
Swede		1	<2.5	<2.5	11	<0.03	0.10		<0.59	<0.21	<0.019	
Wood pigeon muscle		2	<7.0	<7.0	35	<0.04	<0.047		<0.28	<0.11	<0.017	
Wood pigeon muscle	max		<8.0	<8.0		<0.05	<0.048		<0.41		<0.018	
Grass		1	<2.3	<2.3	<4.9	<0.07	1.6		<0.43	<0.18	<0.025	
Soil <sup>d</sup>		1	<4.2	<4.2	11	<0.26	2.1	<4.0	<2.5	<0.83	<0.022	

Material	Selection <sup>a</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>									
			<sup>137</sup> Cs	Total Cs	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Pu	<sup>241</sup> Am	
Milk		9	<0.10	<0.088					<0.000033	<0.000032	<0.12	<0.000025
Milk	max		<0.17	0.14				<0.000047	<0.000042		<0.000038	
Beef kidney		1	0.42	0.51	0.0032	<0.00048	0.0025	<0.000046	0.00023	<0.21	0.0011	
Beef liver		1	0.23	0.48				0.00028	0.0024	<0.22	0.0022	
Beef muscle		1	0.15	0.25				<0.000048	<0.000084	<0.22	0.000028	
Blackberries		1	0.22	0.16				0.000081	0.00061	<0.21	0.0012	
Cabbage		1	<0.06	0.056				<0.000026	0.000055	0.13	0.000044	
Carrots		1	<0.08	0.047							<0.10	
Eggs		1	<0.06	0.081				<0.000037	0.000029	<0.23	0.000035	
Mushrooms		1	0.30	0.22				0.0024	0.016	0.27	0.041	
Pheasant		1	0.06	0.12				<0.000038	0.000081	<0.24	0.00013	
Potatoes		1	0.14	0.087	0.011	<0.00098	0.016	<0.000060	0.00066	<0.30	0.00067	
Rabbit		1	7.5	7.5				<0.000043	0.000065	<0.21	0.00019	
Sheep muscle		2	0.52	0.52				<0.000047	0.000079	<0.23	<0.00014	
Sheep muscle	max			0.54				<0.000053	0.00013	<0.24	0.00019	
Sheep offal		2	0.23	0.32	<0.0013	<0.00033	0.0015	0.00037	0.0032	<0.20	0.0023	
Sheep offal	max		0.27	0.34	0.0019	<0.00038	0.0017	0.00040	0.0036	<0.22	0.0025	
Swede		1	0.09	0.045	<0.00082	<0.00058	<0.00058				<0.10	
Wood pigeon muscle		2	<0.07	0.14				<0.000089	<0.00012	<0.35	0.000089	
Wood pigeon muscle	max		0.09	0.24				<0.000090	0.00017	<0.47	0.00013	
Grass		1	0.30	0.31				0.0010	0.011	<0.22	0.013	
Soil <sup>d</sup>		1	18	23				0.22	4.0	17	1.5	

<sup>a</sup> Data are arithmetic means unless stated as 'max'. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>b</sup> Except for milk where units are Bq l<sup>-1</sup>

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>d</sup> The concentration of <sup>226</sup>Ra was 14 Bq kg<sup>-1</sup>

**Table 2.5. Beta/gamma radioactivity in fish from the Irish Sea vicinity and further afield, 2015**

Location	Material	No. of sampling observations	Mean radioactivity concentration (fresh), Bq kg <sup>-1</sup>							
			Organic <sup>3</sup> H	<sup>3</sup> H	<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>95</sup> Nb	<sup>95</sup> Zr	<sup>99</sup> Tc
<b>Cumbria</b>										
Parton	Cod <sup>c</sup>	2			55	<0.09	<0.047	<0.10	<0.15	<0.75
Whitehaven	Cod <sup>c</sup>	2			70	<0.07	<0.023	<0.10	<0.13	<0.75
Whitehaven	Plaice <sup>a,c</sup>	2	36	72	120	<0.07	<0.073	<0.11	<0.14	1.3
Ravenglass	Plaice <sup>b,c</sup>	2	33	64	110	<0.08	<0.048	<0.30	<0.25	1.4
<b>Lancashire and Merseyside</b>										
Morecambe Bay (Morecambe)	Flounder	2	<28	<35	55	<0.10	<0.025	<0.16	<0.19	<0.48
Ribble Estuary	Flounder	1				<0.05		<0.13	<0.14	
Ribble Estuary	Bass	1				<0.07		<0.20	<0.20	
Liverpool Bay	Plaice	1		<25		<0.05		<0.16	<0.16	
<b>Scotland</b>										
Shetland	Fish meal (salmon)	1 <sup>S</sup>				<0.15		<0.41	<0.43	
Shetland	Fish meal (herring)	1 <sup>S</sup>				<0.13		<0.37	<0.40	
Shetland	Fish oil (salmon)	1 <sup>S</sup>				<0.10		<0.28	<0.30	
Shetland	Fish oil (herring)	1 <sup>S</sup>				<0.10		<0.28	<0.30	
West of Scotland	Salmon	1 <sup>S</sup>				<0.10		<0.28	<0.26	
West of Scotland	Mackerel	1 <sup>S</sup>				<0.10		<0.33	<0.29	
Kircudbright	Plaice	2 <sup>S</sup>			23	<0.10		<0.55	<0.39	<0.14
Inner Solway	Flounder	2 <sup>S</sup>			26	<0.10	<0.10	<0.68	<0.46	<0.25
Inner Solway	Salmon	1 <sup>S</sup>		<5.0		<0.10		<0.44	<0.36	
Inner Solway	Sea trout	1 <sup>S</sup>		<5.0		<0.10		<0.94	<0.58	
<b>Wales</b>										
North Anglesey	Plaice	1	<25	<25	36	<0.03		<0.09	<0.09	
<b>Northern Ireland</b>										
North coast	Lesser spotted dogfish	4 <sup>N</sup>				<0.15		<0.48	<0.44	
Ardglass	Herring	2 <sup>N</sup>				<0.09		<0.41	<0.33	
Kilkeel	Cod	4 <sup>N</sup>			30	<0.05		<0.10	<0.12	
Kilkeel	Plaice	4 <sup>N</sup>				<0.05		<0.14	<0.14	
Kilkeel	Skates / rays	4 <sup>N</sup>				<0.10		<0.25	<0.25	
Kilkeel	Haddock	4 <sup>N</sup>				<0.06		<0.09	<0.11	
<b>Further afield</b>										
Norwegian Sea	Cod	2				<0.06		<0.40	<0.26	

Table 2.5. continued

Location	Material	No. of sampling observations	Mean radioactivity concentration (fresh), Bq kg <sup>-1</sup>						Gross beta
			<sup>106</sup> Ru	<sup>125</sup> Sb	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>144</sup> Ce	<sup>155</sup> Eu	
<b>Cumbria</b>									
Parton	Cod <sup>c</sup>	2	<0.62	<0.19	<0.08	2.3	<0.33	<0.18	150
Whitehaven	Cod <sup>c</sup>	2	<0.53	<0.16	<0.07	2.5	<0.29	<0.15	120
Whitehaven	Plaice <sup>a,c</sup>	2	<0.54	<0.16	<0.07	2.1	<0.32	<0.15	130
Ravenglass	Plaice <sup>b,c</sup>	2	<0.63	<0.17	<0.08	2.1	<0.32	<0.15	140
<b>Lancashire and Merseyside</b>									
Morecambe Bay									
(Morecambe)	Flounder	2	<0.72	<0.21	<0.09	4.3	<0.32	<0.15	
Ribble Estuary	Flounder	1	<0.39	<0.11	<0.05	2.3	<0.18	<0.08	
Ribble Estuary	Bass	1	<0.63	<0.20	<0.08	4.0	<0.42	<0.19	
Liverpool Bay	Plaice	1	<0.43	<0.12	<0.06	1.5	<0.22	<0.09	
<b>Scotland</b>									
Shetland	Fish meal (salmon)	1 <sup>S</sup>	<1.4	<0.39	<0.15	0.27	<0.92	<0.36	
Shetland	Fish meal (herring)	1 <sup>S</sup>	<1.3	<0.34	<0.13	0.47	<0.80	<0.35	
Shetland	Fish oil (salmon)	1 <sup>S</sup>	<0.88	<0.26	<0.10	<0.10	<0.53	<0.22	
Shetland	Fish oil (herring)	1 <sup>S</sup>	<0.95	<0.27	<0.10	<0.10	<0.52	<0.22	
West of Scotland	Salmon	1 <sup>S</sup>	<0.75	<0.21	<0.10	<0.10	<0.44	<0.19	
West of Scotland	Mackerel	1 <sup>S</sup>	<0.71	<0.20	<0.10	0.44	<0.49	<0.20	
Kircudbright	Plaice	2 <sup>S</sup>	<0.87	<0.24	<0.10	<0.11	<0.54	<0.23	
Inner Solway	Flounder	2 <sup>S</sup>	<0.98	<0.28	<0.11	<4.6	<0.58	<0.24	
Inner Solway	Salmon	1 <sup>S</sup>	<0.91	<0.25	<0.10	0.15	<0.53	<0.21	
Inner Solway	Sea trout	1 <sup>S</sup>	<1.1	<0.27	<0.10	1.9	<0.71	<0.24	
<b>Wales</b>									
North Anglesey	Plaice	1	<0.29	<0.08	<0.03	0.41	<0.17	<0.08	
<b>Northern Ireland</b>									
North coast	Lesser spotted dogfish	4 <sup>N</sup>	<1.2	<0.32	<0.14	0.60	<0.57	<0.23	
Ardglass	Herring	2 <sup>N</sup>	<0.79	<0.20	<0.10	<0.20	<0.35	<0.15	
Kilkeel	Cod	4 <sup>N</sup>	<0.42	<0.12	<0.05	0.74	<0.23	<0.11	
Kilkeel	Plaice	4 <sup>N</sup>	<0.43	<0.12	<0.05	0.50	<0.24	<0.11	
Kilkeel	Skates / rays	4 <sup>N</sup>	<0.74	<0.21	<0.10	0.87	<0.37	<0.17	
Kilkeel	Haddock	4 <sup>N</sup>	<0.43	<0.12	<0.06	0.30	<0.21	<0.10	
<b>Further afield</b>									
Norwegian Sea	Cod	2	<0.48	<0.12	<0.05	0.14	<0.24	<0.09	

<sup>a</sup> The concentrations of <sup>129</sup>I and <sup>147</sup>Pm were <1.1 and <0.021 Bq kg<sup>-1</sup> respectively

<sup>b</sup> The concentrations of <sup>129</sup>I and <sup>147</sup>Pm were <1.1 and 0.054 Bq kg<sup>-1</sup> respectively

<sup>c</sup> Data for natural radionuclides for some of these samples may be available in Table 7.7

<sup>N</sup> Measurements labelled "N" are made on behalf of the Northern Ireland Environment Agency

<sup>S</sup> Measurements labelled "S" are made on behalf of the Scottish Environment Protection Agency

**Table 2.6. Beta/gamma radioactivity in shellfish from the Irish Sea vicinity and further afield, 2015**

Location	Material	No. of sampling observations	Mean radioactivity concentration (fresh), Bq kg <sup>-1</sup>								
			Organic		<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>95</sup> Nb	<sup>95</sup> Zr	<sup>99</sup> Tc	<sup>106</sup> Ru
			<sup>3</sup> H	<sup>3</sup> H							
<b>Cumbria</b>											
Parton	Crabs <sup>d</sup>	2			110	<0.11	0.092	<0.18	<0.23	3.8	<0.80
Parton	Lobsters <sup>d</sup>	2			100	<0.06	<0.11	<0.09	<0.12	37	<0.45
Parton	Winkles <sup>d</sup>	2			100	<0.53	0.77	<0.18	<0.22	34	<1.4
Whitehaven	<i>Nephrops</i> <sup>a,d</sup>	2	<25	<35	75	<0.08	0.076	<0.29	<0.24	25	<0.64
Whitehaven outer harbour	Mussels <sup>d</sup>	2			93	<0.25	0.25	<0.07	<0.10	26	1.3
Nethertown	Winkles <sup>b,d</sup>	4	<26	<26	120	0.94	1.5	<0.15	<0.19	26	<3.2
Sellafield coastal area	Crabs <sup>c,d</sup>	2	100	90	150	0.49	0.24	<0.20	<0.22	8.2	<0.87
Sellafield coastal area	Lobsters <sup>d</sup>	2	150	160	160	<0.16	0.054	<0.21	<0.24	180	<0.81
Ravenglass	Mussels <sup>d</sup>	2			160	0.46	0.22	<0.07	<0.09	62	2.2
Seascale Area	Common prawns <sup>d</sup>	2	<25	28	100	<0.07	<0.059	<0.15	<0.16	<0.61	<0.52
<b>Lancashire and Merseyside</b>											
Morecambe Bay (Morecambe)	Shrimps	2	35	42	85	<0.06	<0.063	<0.23	<0.20	0.52	<0.50
Morecambe Bay (Morecambe)	Mussels	2	66	64	85	<0.08	0.21	<0.10	<0.14	41	<0.57
Morecambe Bay (Middleton Sands)	Winkles	2	39	<25	61	<0.09	0.12	<0.13	<0.18	6.8	<0.68
Ribble Estuary	Shrimps	1			64	<0.07		<0.12	<0.15	<0.88	<0.52
Ribble Estuary	Mussels	1				<0.06		<0.16	<0.16		<0.47
Liverpool Bay	Mussels	1		<25		<0.13		<0.16	<0.22		<0.94
Dee Estuary	Cockles	1		<25		<0.04		<0.06	<0.08	1.5	<0.32
Wirral	Shrimps	1		<25		<0.04		<0.09	<0.10	1.1	<0.33
<b>Scotland</b>											
Kinlochbervie	Crabs	2 <sup>s</sup>				<0.10		<0.14	<0.14	<10	<0.40
Lewis	Mussels	1 <sup>s</sup>				<0.10		<1.2	<0.56		<0.82
Skye	Lobsters	1 <sup>s</sup>				<0.10		<0.74	<0.41	3.9	<0.67
Skye	Mussels	1 <sup>s</sup>				<0.10		<0.63	<0.35		<0.58
Islay	Crabs	1 <sup>s</sup>				<0.10		<0.45	<0.24		<0.35
Islay	Scallops	1 <sup>s</sup>				<0.10		<0.45	<0.31		<0.61
Kirkcudbright	Scallops	2 <sup>s</sup>				<0.10		<0.36	<0.24	<0.19	<0.68
Kirkcudbright	Queens	2 <sup>s</sup>				<0.10		<0.47	<0.32	<0.20	<0.55
Kirkcudbright	Crabs <sup>d</sup>	1 <sup>s</sup>			94	<0.10	0.14	<0.35	<0.29	3.3	<0.67
Kirkcudbright	Lobsters <sup>d</sup>	2 <sup>s</sup>			65	<0.10	<0.10	<0.35	<0.33	43	<0.90
Kirkcudbright	Limpets <sup>d</sup>	1 <sup>s</sup>				<0.10		<0.56	<0.35		<0.63
Kirkcudbright	Winkles <sup>d</sup>	2 <sup>s</sup>				<0.18	<0.10	<0.10	<0.12	20	<0.46
Cutters Pool	Winkles	1 <sup>s</sup>				0.12		<0.12	<0.14		<0.48
Southernness	Winkles	2 <sup>s</sup>		<5.0		<0.10	0.16	<0.29	<0.24	25	<0.67
North Solway coast	Cockles	1 <sup>s</sup>		<5.0		<0.10		<0.46	<0.35		<0.75
North Solway coast	Mussels	2 <sup>s</sup>		<5.0	77	<0.14	0.37	<0.31	<0.29	23	<0.82
Inner Solway	Shrimps	2 <sup>s</sup>		<5.0		<0.10	<0.10	<0.55	<0.36	<0.16	<0.82
<b>Wales</b>											
North Anglesey	Crabs	1	40	<25	56	<0.06		<0.11	<0.15		<0.56
North Anglesey	Lobsters	1	<25	<25	58	<0.08		<0.09	<0.13	16	<0.62
<b>Northern Ireland</b>											
Ballycastle	Lobsters	2 <sup>N</sup>				<0.11		<0.45	<0.38	2.6	<0.95
County Down	Scallops	2 <sup>N</sup>				<0.07		<0.80	<0.41		<0.63
Kilkeel	Crabs	4 <sup>N</sup>				<0.06		<0.16	<0.17		<0.54
Kilkeel	Lobsters	4 <sup>N</sup>				<0.05		<0.13	<0.13	8.1	<0.44
Kilkeel	<i>Nephrops</i>	4 <sup>N</sup>				<0.12		<0.33	<0.33	6.0	<0.93
Minerstown	Winkles	4 <sup>N</sup>				<0.05		<0.09	<0.11		<0.42
Carlingford Lough	Mussels	2 <sup>N</sup>				<0.11		<0.24	<0.28	1.8	<0.96
<b>Further afield</b>											
Cromer	Crabs	2				<0.07		<0.15	<0.17		<0.59
Southern North Sea	Cockles	2				<0.03		<0.04	<0.06		<0.22

Table 2.6. continued

Location	Material	No. of sampling observations	Mean radioactivity concentration (fresh), Bq kg <sup>-1</sup>							Gross beta
			<sup>110m</sup> Ag	<sup>125</sup> Sb	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>144</sup> Ce	<sup>147</sup> Pm	<sup>155</sup> Eu	
<b>Cumbria</b>										
Parton	Crabs <sup>d</sup>	2	<0.13	<0.23	<0.10	0.74	<0.45		<0.22	120
Parton	Lobsters <sup>d</sup>	2	<0.07	<0.14	<0.05	1.1	<0.28		<0.14	170
Parton	Winkles <sup>d</sup>	2	<0.12	<0.24	<0.09	3.7	<0.40		<0.20	200
Whitehaven	<i>Nephrops</i> <sup>a,d</sup>	2	<0.11	<0.18	<0.07	2.2	<0.31	0.094	<0.14	170
Whitehaven outer harbour	Mussels <sup>d</sup>	2	<0.06	<0.16	<0.05	1.3	<0.19		<0.09	80
Nethertown	Winkles <sup>b,d</sup>	4	<0.13	<0.31	<0.09	4.9	<0.39	0.82	<0.21	190
Sellafield coastal area	Crabs <sup>c,d</sup>	2	<0.14	<0.22	<0.09	1.0	<0.40	0.17	<0.19	110
Sellafield coastal area	Lobsters <sup>d</sup>	2	<0.15	<0.23	<0.10	1.3	<0.43		<0.21	230
Ravenglass	Mussels <sup>d</sup>	2	<0.06	<0.22	<0.05	1.0	<0.21		<0.10	120
Seascale Area	Common prawns <sup>d</sup>	2	<0.09	<0.14	<0.06	1.2	<0.29		<0.14	98
<b>Lancashire and Merseyside</b>										
Morecambe Bay (Morecambe)	Shrimps	2	<0.08	<0.15	<0.06	2.5	<0.31		<0.14	
Morecambe Bay (Morecambe)	Mussels	2	<0.08	<0.16	<0.07	2.2	<0.27		<0.13	65
Morecambe Bay (Middleton Sands)	Winkles	2	<0.11	<0.21	<0.09	2.6	<0.36		<0.18	88
Ribble Estuary	Shrimps	1	<0.09	<0.14	<0.07	1.3	<0.25		<0.11	
Ribble Estuary	Mussels	1	<0.08	<0.14	<0.06	1.0	<0.27		<0.13	
Liverpool Bay	Mussels	1	<0.14	<0.27	<0.11	0.95	<0.42		<0.17	
Dee Estuary	Cockles	1	<0.05	<0.09	<0.04	0.90	<0.16		<0.08	
Wirral	Shrimps	1	<0.05	<0.10	<0.04	0.78	<0.23		<0.10	
<b>Scotland</b>										
Kinlochbervie	Crabs	2 <sup>S</sup>	<0.10	<0.12	<0.10	<0.10	<0.24		<0.11	
Lewis	Mussels	1 <sup>S</sup>	<0.10	<0.21	<0.10	<0.10	<0.56		<0.21	
Skye	Lobsters	1 <sup>S</sup>	<0.10	<0.17	<0.10	<0.10	<0.41		<0.15	
Skye	Mussels	1 <sup>S</sup>	<0.10	<0.15	<0.10	<0.10	<0.37		<0.13	
Islay	Crabs	1 <sup>S</sup>	<0.10	<0.10	<0.10	<0.10	<0.23		<0.10	
Islay	Scallops	1 <sup>S</sup>	<0.10	<0.16	<0.10	<0.10	<0.35		<0.14	
Kirkcudbright	Scallops	2 <sup>S</sup>	<0.10	<0.16	<0.10	<0.10	<0.35		<0.14	
Kirkcudbright	Queens	2 <sup>S</sup>	<0.11	<0.19	<0.10	<0.15	<0.41		<0.17	
Kirkcudbright	Crabs <sup>d</sup>	1 <sup>S</sup>	<0.10	<0.19	<0.10	0.69	<0.45		<0.19	
Kirkcudbright	Lobsters <sup>d</sup>	2 <sup>S</sup>	<0.14	<0.26	<0.10	0.79	<0.52		<0.23	
Kirkcudbright	Limpets <sup>d</sup>	1 <sup>S</sup>	<0.10	<0.16	<0.10	0.79	<0.35		<0.13	
Kirkcudbright	Winkles <sup>d</sup>	2 <sup>S</sup>	<0.10	<0.14	<0.10	0.73	<0.29		<0.13	
Cutters Pool	Winkles	1 <sup>S</sup>	<0.10	<0.15	<0.10	1.7	<0.29		<0.13	
Southernness	Winkles	2 <sup>S</sup>	<0.13	<0.20	<0.11	0.78	<0.39		<0.17	
North Solway coast	Cockles	1 <sup>S</sup>	<0.10	<0.21	<0.10	0.36	<0.48		<0.19	
North Solway coast	Mussels	2 <sup>S</sup>	<0.15	<0.26	<0.10	1.7	<0.54		<0.22	
Inner Solway	Shrimps	2 <sup>S</sup>	<0.14	<0.22	<0.10	3.0	<0.49		<0.19	
<b>Wales</b>										
North Anglesey	Crabs	1	<0.09	<0.16	<0.07	0.37	<0.31		<0.15	
North Anglesey	Lobsters	1	<0.10	<0.18	<0.07	0.50	<0.32		<0.16	120
<b>Northern Ireland</b>										
Ballycastle	Lobsters	2 <sup>N</sup>	<0.16	<0.25	<0.11	0.16	<0.43		<0.16	
County Down	Scallops	2 <sup>N</sup>	<0.12	<0.16	<0.07	0.11	<0.36		<0.15	
Kilkeel	Crabs	4 <sup>N</sup>	<0.09	<0.15	<0.07	<0.10	<0.28		<0.13	
Kilkeel	Lobsters	4 <sup>N</sup>	<0.07	<0.12	<0.06	0.15	<0.22		<0.10	
Kilkeel	<i>Nephrops</i>	4 <sup>N</sup>	<0.16	<0.25	<0.12	0.66	<0.43		<0.18	
Minerstown	Winkles	4 <sup>N</sup>	<0.07	<0.12	<0.05	<0.15	<0.23		<0.11	
Carlingford Lough	Mussels	2 <sup>N</sup>	<0.16	<0.28	<0.12	<0.34	<0.52		<0.23	
<b>Further afield</b>										
Cromer	Crabs	2	<0.09	<0.17	<0.07	<0.07	<0.32		<0.15	
Southern North Sea	Cockles	2	<0.03	<0.07	<0.03	0.07	<0.12		<0.06	

<sup>a</sup> The concentration of <sup>129</sup>I was <1.0 Bq kg<sup>-1</sup>

<sup>b</sup> The concentration of <sup>129</sup>I was <1.5 Bq kg<sup>-1</sup>

<sup>c</sup> The concentration of <sup>129</sup>I was <1.1 Bq kg<sup>-1</sup>

<sup>d</sup> Data for natural radionuclides for some of these samples may be available in Table 7.7

<sup>N</sup> Measurements labelled "N" are made on behalf of the Northern Ireland Environment Agency

<sup>S</sup> Measurements labelled "S" are made on behalf of the Scottish Environment Protection Agency

**Table 2.7. Concentrations of transuranic radionuclides in fish and shellfish from the Irish Sea vicinity and further afield, 2015**

Location	Material	No. of sampling observations	Mean radioactivity concentration (fresh), Bq kg <sup>-1</sup>						
			<sup>237</sup> Np	<sup>238</sup> Pu	<sup>239</sup> Pu+ <sup>240</sup> Pu	<sup>241</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm+ <sup>244</sup> Cm
<b>Cumbria</b>									
Parton	Cod	2		0.0011	0.0064	<0.12	0.012	*	0.000016
Parton	Crabs	2		0.028	0.15	0.96	0.67	*	*
Parton	Lobsters	2		0.027	0.13	0.82	1.5	*	0.0038
Parton	Winkles	2		0.96	5.5	25	11	*	*
Whitehaven	Cod	2		0.0049	0.026	<0.25	0.038	*	*
Whitehaven	Plaice	2	0.00010	0.00083	0.0047	<0.17	0.011	*	0.000029
Whitehaven	<i>Nephrops</i>	2	0.0013	0.053	0.31	1.6	1.5	*	0.0017
Whitehaven outer harbour	Mussels	2		0.56	2.9	15	7.3	*	*
Nethertown	Winkles	4	0.015	1.5	8.5	40	19	*	*
Sellafield coastal area	Crabs	2	0.0021	0.060	0.32	2.2	1.6	*	*
Sellafield coastal area	Lobsters	2		0.047	0.22	1.9	3.0	*	*
Ravenglass	Plaice	2	0.000052	0.00097	0.0045	<0.15	0.011	*	*
Ravenglass	Mussels	2		0.81	4.2	24	9.6	*	*
Seascale	Prawns	2		0.0034	0.020	<0.30	0.044	0.00014	0.000098
<b>Lancashire and Merseyside</b>									
Morecambe Bay (Morecambe)	Flounder	2		0.00034	0.0022		0.0046	*	0.000039
Morecambe Bay (Morecambe)	Shrimps	2		0.0039	0.024		0.041	*	*
Morecambe Bay (Morecambe)	Mussels	2		0.31	1.8	9.5	3.5	*	*
Morecambe Bay (Middleton Sands)	Winkles	2		0.21	1.3	6.5	2.5	*	0.0041
Ribble Estuary	Flounder	1					<0.05		
Ribble Estuary	Bass	1					<0.18		
Ribble Estuary	Shrimps	1	0.000058	0.0015	0.0090		0.016	*	*
Ribble Estuary	Mussels	1		0.077	0.46		0.79	*	*
Liverpool Bay	Mussels	1					<0.06		
Liverpool Bay	Plaice	1					1.2		
Dee Estuary	Cockles	1		0.067	0.45		1.3	*	*
Wirral	Shrimps	1		0.0020	0.013		0.015	*	*
<b>Scotland</b>									
Shetland	Fish meal (salmon)	1 <sup>s</sup>		<0.0034	0.0085		0.020		
Shetland	Fish meal (herring)	1 <sup>s</sup>		<0.0036	0.023		0.0083		
Shetland	Fish oil (salmon)	1 <sup>s</sup>		<0.00080	<0.00060		<0.021		
Shetland	Fish oil (herring)	1 <sup>s</sup>		<0.00060	<0.00070		0.0053		
West of Scotland	Salmon	1 <sup>s</sup>		0.011	0.047		0.044		
West of Scotland	Mackerel	1 <sup>s</sup>		<0.0038	<0.0038		0.014		
Kinlochbervie	Crabs	2 <sup>s</sup>		<0.035	<0.0035		0.16		
Lewis	Mussels	1 <sup>s</sup>					<0.11		
Skye	Lobsters	1 <sup>s</sup>					<0.10		
Skye	Mussels	1 <sup>s</sup>					<0.10		
Islay	Crabs	1 <sup>s</sup>					<0.10		
Islay	Scallops	1 <sup>s</sup>					<0.10		
Kirkcudbright	Plaice	2 <sup>s</sup>		0.0018	0.0070		0.016		
Kirkcudbright	Scallops	2 <sup>s</sup>		0.016	0.087		0.024		
Kirkcudbright	Queens	2 <sup>s</sup>		<0.0016	0.017		0.013		
Kirkcudbright	Crabs	1 <sup>s</sup>		<0.059	0.20	<3.0	1.2		
Kirkcudbright	Lobsters	2 <sup>s</sup>		0.014	0.080	0.28	0.55		
Kirkcudbright	Winkles	2 <sup>s</sup>		0.21	1.3		2.6		
Kirkcudbright	Limpets	1 <sup>s</sup>					2.4		
Cutters Pool	Winkles	1 <sup>s</sup>					6.6		
Southernness	Winkles	2 <sup>s</sup>		0.099	0.61	<2.4	1.3		
North Solway coast	Cockles	1 <sup>s</sup>		0.070	0.42		0.60		
North Solway coast	Mussels	2 <sup>s</sup>		0.46	2.7	8.7	5.1		
Inner Solway	Flounder	1 <sup>s</sup>		<0.0081	<0.0072		0.010		
Inner Solway	Salmon	1 <sup>s</sup>					<0.12		
Inner Solway	Sea trout	1 <sup>s</sup>					<0.14		
Inner Solway	Shrimps	1 <sup>s</sup>		<0.0019	0.012		0.034		

Table 2.7. continued

Location	Material	No. of sampling observations	Mean radioactivity concentration (fresh), Bq kg <sup>-1</sup>						
			<sup>237</sup> Np	<sup>238</sup> Pu	<sup>239</sup> Pu+ <sup>240</sup> Pu	<sup>241</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm+ <sup>244</sup> Cm
<b>Wales</b>									
North Anglesey	Plaice	1					<0.11		
North Anglesey	Crabs	1					<0.19		
North Anglesey	Lobsters	1		0.0053	0.032	0.17	0.13	*	*
Northern Ireland									
North coast	Lesser spotted dogfish	4 <sup>N</sup>					<0.15		
Ballycastle	Lobsters	2 <sup>N</sup>					0.20		
County Down	Scallops	2 <sup>N</sup>					<0.18		
Ardglass	Herring	2 <sup>N</sup>					<0.09		
Kilkeel	Cod	4 <sup>N</sup>					<0.08		
Kilkeel	Plaice	4 <sup>N</sup>					<0.13		
Kilkeel	Skates / rays	4 <sup>N</sup>					<0.13		
Kilkeel	Haddock	4 <sup>N</sup>					<0.09		
Kilkeel	Crabs	4 <sup>N</sup>					<0.15		
Kilkeel	Lobsters	4 <sup>N</sup>					<0.09		
Kilkeel	<i>Nephrops</i>	1 <sup>N</sup>		0.0083	0.051		0.23	0.00023	0.00015
Minerstown	Winkles	1 <sup>N</sup>		0.022	0.15		0.11	*	*
Carlingford Lough	Mussels	2 <sup>N</sup>					<0.25		
<b>Further afield</b>									
Norwegian Sea	Cod	2					<0.06		
Cromer	Crabs	2					<0.15		
Southern North Sea	Cockles	1		0.0017	0.012		0.0087	*	*

\* Not detected by the method used

<sup>N</sup> Measurements labelled "N" are made on behalf of the Northern Ireland Environment Agency

<sup>S</sup> Measurements labelled "S" are made on behalf of the Scottish Environment Protection Agency

**Table 2.8. Concentrations of radionuclides in sediment from the Cumbrian coast and further afield, 2015**

Location	Material	No. of sampling observations	Mean radioactivity concentration (dry), Bq kg <sup>-1</sup>								
			<sup>54</sup> Mn	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>95</sup> Nb	<sup>95</sup> Zr	<sup>106</sup> Ru	<sup>125</sup> Sb	<sup>134</sup> Cs	<sup>137</sup> Cs
<b>Cumbria</b>											
Newton Arlosh	Sediment	4	<0.59			<0.41	<1.4	<4.4	<2.3	<0.56	170
Maryport Outer Harbour	Sediment	2	<0.58	6.7		<0.28	<0.80	<2.4	<1.3	<0.32	61
Workington Harbour	Sediment	2	<0.54			<0.52	<1.4	<3.8	<1.9	<0.52	48
Harrington Harbour	Sediment	2	<0.65			<0.39	<1.1	<3.2	<1.6	<0.41	120
Whitehaven Outer Harbour	Sediment	4	<0.43	<3.0		<0.28	<0.65	<2.2	<1.2	<0.30	56
St Bees beach	Sediment	4	0.77			<0.23	<0.71	<2.2	<1.2	<0.29	53
Sellafield beach, S of former pipeline	Sediment	2	<0.51			<0.23	<0.85	<2.3	<1.1	<0.30	42
River Calder - downstream	Sediment	2	<0.39			<0.27	<0.67	<2.3	<1.3	<0.33	57
River Calder - upstream	Sediment	2	<0.59			<0.47	<1.2	<3.3	<1.8	<0.52	40
Seascale beach	Sediment	4	<0.46			<0.23	<0.57	<1.9	<1.0	<0.28	28
Ravenglass - Carleton Marsh	Sediment	4	2.6			<0.40	<0.92	8.7	<1.8	<0.44	210
River Mite Estuary (erosional)	Sediment	4	2.6	48		<0.65	<1.7	<9.5	<3.4	<0.73	360
Ravenglass - Raven Villa	Sediment	4	1.4			<0.37	<0.96	<5.2	<1.7	<0.41	110
Newbiggin (Eskmeals)	Sediment	4	3.5	94		<0.44	<1.2	<5.6	<2.1	<0.48	300
Haverigg	Sediment	2	0.77			<0.31	<0.87	<2.5	<1.3	<0.35	47
Millom	Sediment	2	<0.72			<0.39	<1.3	<3.5	<1.8	<0.46	65
Low Shaw	Sediment	2	<0.48			<0.30	<1.1	<3.0	<1.6	<0.41	60
Walney Channel - N of discharge point	Sediment	2	<0.41			<0.34	<0.92	<2.7	<1.4	<0.37	70
Walney Channel - S of discharge point	Sediment	2	<0.71			<0.33	<0.86	<2.9	<1.4	<0.37	88
Sand Gate Marsh	Sediment	4	<0.48			<0.34	<1.1	<3.2	<1.7	<0.42	77
Kents Bank	Sediment	4	<0.51			<0.38	<1.3	<4.1	<2.1	<0.48	220
<b>Lancashire</b>											
Morecambe	Sediment	2	<0.28								3.0
Half Moon Bay	Sediment	2	<0.51								33
Red Nab Point	Sediment	2	<0.50								33
Potts Corner	Sediment	2	<0.43								11
Sunderland Point	Sediment	4	<0.50			<0.45	<1.1	<3.4	<1.8	<0.49	63
Conder Green	Sediment	4	<0.58			<0.51	<1.4	<3.9	<2.0	<0.56	63
Hambleton	Sediment	4	<0.56			<0.46	<1.2	<4.3	<2.2	<0.51	240
Skippool Creek	Sediment	4	<0.91			<0.67	<2.0	<6.7	<3.5	<0.83	190
Fleetwood	Sediment	4	<0.35			<0.23	<0.76	<2.1	<1.1	<0.30	7.5
Blackpool	Sediment	4	<0.31			<0.20	<0.56	<1.8	<1.0	<0.28	2.2
Crossens Marsh	Sediment	4	<1.4			<1.0	<3.2	<10	<5.0	<1.3	150
Ainsdale	Sediment	4	<0.31			<0.20	<0.70	<1.9	<1.0	<0.28	3.1
Rock Ferry	Sediment	4	<0.50			<0.36	<1.1	<3.1	<1.6	<0.43	56
New Brighton	Sediment	4	<0.32			<0.22	<0.68	<1.9	<1.0	<0.29	2.7
<b>Scotland</b>											
Campbeltown	Sediment	1 <sup>S</sup>	<0.10	<0.10		<0.37	<0.22	<0.38	<0.11	<0.10	7.2
Garlieston	Sediment	1 <sup>S</sup>	<0.10	0.11		<0.53	<0.15	<0.63	<0.21	<0.10	24
Innerwell	Sediment	2 <sup>S</sup>	<0.10	0.17		<0.41	<0.15	<0.75	<0.26	<0.11	37
Carlsruith	Sediment	1 <sup>S</sup>	<0.10	0.17		<0.60	<0.38	<0.72	<0.24	<0.10	19
Skyreburn	Sediment	2 <sup>S</sup>	<0.10	<0.10		<0.39	<0.30	<0.76	<0.24	<0.10	21
Kirkcudbright	Sediment <sup>a</sup>	2 <sup>S</sup>	<0.10	<0.24		<0.27	<0.20	<0.79	<0.31	<0.11	74
Balcary Bay	Sediment <sup>a</sup>	1 <sup>S</sup>	<0.10	0.29		<0.63	<0.27	<0.81	<0.29	<0.10	59
Palnackie Harbour	Sediment	2 <sup>S</sup>	<0.10	0.34		<0.38	<0.21	<0.70	<0.36	<0.11	78
Gardenburn	Sediment	2 <sup>S</sup>	<0.10	0.36		<0.74	<0.38	<0.94	<0.36	<0.12	84
Kippford Slipway	Sediment	2 <sup>S</sup>	<0.10	0.82		<0.33	<0.23	3.0	0.66	<0.13	160
Kippford Merse	Sediment	1 <sup>S</sup>	<0.10	0.33		<0.51	<0.26	1.1	<0.31	<0.10	310
Kirkconnell Merse	Sediment	1 <sup>S</sup>	<0.10	0.39		<0.34	<0.19	<0.75	0.61	<0.10	400
Southernness	Sediment	1 <sup>S</sup>	<0.10	<0.10		<0.20	<0.20	<0.65	<0.22	<0.10	19

Table 2.8. continued

Location	Material	No. of sampling observations	Mean radioactivity concentration (dry), Bq kg <sup>-1</sup>								Gross alpha	Gross beta
			<sup>144</sup> Ce	<sup>154</sup> Eu	<sup>155</sup> Eu	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Pu	<sup>241</sup> Am			
<b>Cumbria</b>												
Newton Arlosh	Sediment	4	<2.3	<1.4	<2.0					240	630	840
Maryport Outer Harbour	Sediment	2	<1.7	<0.80	<0.69	21	120	680		140	640	730
Workington Harbour	Sediment	2	<2.0	<1.3	<0.86					33	370	780
Harrington Harbour	Sediment	2	<2.1	<1.1	<0.86					150	340	880
Whitehaven Outer Harbour	Sediment	4	<1.5	<0.85	<0.66	15	89	440		140	290	550
St Bees beach	Sediment	4	<1.3	<0.80	<0.58					150	190	470
Sellafield beach, S of former pipeline	Sediment	2	<1.4	<0.83	<0.58					130	350	430
River Calder - downstream	Sediment	2	<1.6	<0.90	<0.68					70	150	550
River Calder - upstream	Sediment	2	<2.0	<1.5	<1.7					<0.73	330	1300
Seascale beach	Sediment	4	<1.2	<0.75	<0.56					130	250	500
Ravenglass - Carleton Marsh	Sediment	4	<2.4	2.2	<1.1					740	1300	1100
River Mite Estuary (erosional)	Sediment	4	<3.2	3.2	<1.5	120	680	3500		1300	2300	1200
Ravenglass - Raven Villa	Sediment	4	<1.9	1.2	<0.84					460	940	790
Newbiggin (Eskmeals)	Sediment	4	<2.7	3.4	<1.2	110	620	3500		1200	1900	1100
Haverigg	Sediment	2	<1.8	<0.84	<0.73					160	430	660
Millom	Sediment	2	<1.9	<1.2	<0.82					180	350	630
Low Shaw	Sediment	2	<1.6	<1.2	<0.68					110	300	560
Walney Channel - N of discharge point	Sediment	2	<1.8	<0.93	<0.79					160	680	660
Walney Channel - S of discharge point	Sediment	2	<1.8	<0.88	<0.80					210	470	780
Sand Gate Marsh	Sediment	4	<1.9	<1.2	<0.79					70	240	650
Kents Bank	Sediment	4	<2.4	<1.2	<1.4					130	430	880
<b>Lancashire</b>												
Morecambe	Sediment	2								1.2		
Half Moon Bay	Sediment	2				5.0	29			59		
Red Nab Point	Sediment	2								48		
Potts Corner	Sediment	2								9.8		
Sunderland Point	Sediment	4	<2.0	<1.2	<0.87					96	380	780
Conder Green	Sediment	4	<2.1	<1.4	<0.92					76	370	870
Hambleton	Sediment	4	<2.5	<1.4	<1.1					240	590	1200
Skippool Creek	Sediment	4	<3.0	<2.3	<1.3					210	640	1100
Fleetwood	Sediment	4	<1.2	<0.85	<0.50					11	<120	350
Blackpool	Sediment	4	<0.99	<0.75	<0.44					3.6	<120	200
Crossens Marsh	Sediment	4	<4.0	<3.3	<1.8					140	470	1100
Ainsdale	Sediment	4	<1.0	<0.77	<0.44					2.8	<110	260
Rock Ferry	Sediment	4	<1.7	<1.2	<0.75					45	340	820
New Brighton	Sediment	4	<1.1	<0.80	<0.47					2.8	<110	280
<b>Scotland</b>												
Campbeltown	Sediment	1 <sup>S</sup>	<0.47	<0.10	0.78					1.5		
Garlieston	Sediment	1 <sup>S</sup>	<0.78	<0.19	0.44	5.6	40			66		
Innerwell	Sediment	2 <sup>S</sup>	<0.73	<0.16	0.80	8.3	49			83		
Carsluith	Sediment	1 <sup>S</sup>	<0.76	0.30	0.66	11	68			120	180	810
Skyreburn	Sediment	2 <sup>S</sup>	<0.75	<0.18	0.72	2.7	16			22		
Kirkcudbright	Sediment <sup>a</sup>	2 <sup>S</sup>	<0.76	<0.37	<0.71					100		
Balcary Bay	Sediment <sup>a</sup>	1 <sup>S</sup>	<1.1	0.39	0.84					120		
Palnackie Harbour	Sediment	2 <sup>S</sup>	<0.77	<0.31	1.1	12	74			150		
Gardenburn	Sediment	2 <sup>S</sup>	<1.2	0.41	0.88	53	47			160		
Kippford Slipway	Sediment	2 <sup>S</sup>	<0.93	0.81	<0.67	85	110			350		
Kippford Merse	Sediment	1 <sup>S</sup>	<0.74	1.1	0.90	34	280			440	160	1200
Kirkconnell Merse	Sediment	1 <sup>S</sup>	<0.87	0.80	1.2	150	27			270		
Southernness	Sediment	1 <sup>S</sup>	<0.63	<0.15	<0.31	2.9	17			28		

**Table 2.8. continued**

Location	Material	No. of sampling observations	Mean radioactivity concentration (dry), Bq kg <sup>-1</sup>								
			<sup>54</sup> Mn	<sup>60</sup> Co	<sup>95</sup> Nb	<sup>95</sup> Zr	<sup>106</sup> Ru	<sup>125</sup> Sb	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>144</sup> Ce
<b>Wales</b>											
Rhyl	Sediment	2	<0.69	<0.52	<1.5	<4.4	<2.3	<0.61	51	<2.1	
Llandudno	Sediment	2	<0.28	<0.23	<0.55	<1.8	<0.96	<0.26	1.7	<1.1	
Caerhun	Sediment	2	<0.64	<0.47	<1.2	<3.9	<2.1	<0.57	47	<2.0	
Llanfairfechan	Sediment	2	<0.48	<0.31	<0.98	<2.8	<1.5	<0.42	20	<1.6	
<b>Northern Ireland</b>											
Carrichue	Mud and shell	1 <sup>N</sup>	<0.26	<0.23	<1.3	<1.0	<2.3	<0.65	<0.29	1.4	<1.9
Carrichue	Sand and shell	1 <sup>N</sup>	<0.29	<0.20	<1.7	<1.2	<2.2	<0.67	<0.30	1.3	<1.9
Portrush	Sand	2 <sup>N</sup>	<0.24	<0.19	<1.7	<1.1	<2.0	<0.56	<0.26	0.48	<1.5
Oldmill Bay	Mud and sand	1 <sup>N</sup>	<0.29	<0.23	<3.8	<1.7	<2.5	<0.66	<0.32	5.9	<1.7
Oldmill Bay	Sand	1 <sup>N</sup>	<0.25	<0.18	<1.9	<1.2	<2.1	<0.65	<0.26	4.6	<1.8
Ballymacormick	Mud and sand	1 <sup>N</sup>	<0.51	<0.39	<6.2	<2.9	<3.9	<0.99	<0.55	7.0	<2.3
Ballymacormick	Sand	1 <sup>N</sup>	<0.28	<0.21	<1.7	<1.2	<2.2	<0.69	<0.32	4.8	<2.0
<b>Strangford Lough</b>											
- Nicky's Point	Mud	2 <sup>N</sup>	<0.44	<0.34	<2.9	<1.9	<3.6	<1.0	<0.47	14	<2.6
Dundrum Bay	Mud	1 <sup>N</sup>	<0.43	<0.38	<2.7	<1.8	<3.2	<0.85	<0.49	4.1	<2.0
Dundrum Bay	Mud and sand	1 <sup>N</sup>	<0.55	<0.42	<4.9	<2.7	<4.4	<1.1	<0.58	3.5	<2.6
Carlingford Lough	Mud	2 <sup>N</sup>	<0.74	<0.62	<4.3	<2.9	<5.9	<1.7	<0.81	43	<3.4

Location	Material	No. of sampling observations	Mean radioactivity concentration (dry), Bq kg <sup>-1</sup>									
			<sup>154</sup> Eu	<sup>155</sup> Eu	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm + <sup>244</sup> Cm	Gross alpha	Gross beta	
<b>Wales</b>												
Rhyl	Sediment	2	<1.7	<2.1			39				290	970
Llandudno	Sediment	2	<0.68	<0.47			1.3				<120	240
Caerhun	Sediment	2	<1.6	<0.88			30				400	760
Llanfairfechan	Sediment	2	<1.2	<0.66			15				220	480
<b>Northern Ireland</b>												
Carrichue	Mud and shell	1 <sup>N</sup>	<0.68	<0.82	0.080	0.57	1.0	*	*			
Carrichue	Sand and shell	1 <sup>N</sup>	<0.64	<0.83			<1.0					
Portrush	Sand	2 <sup>N</sup>	<0.63	<0.65			<0.66					
Oldmill Bay	Mud and sand	1 <sup>N</sup>	<0.82	<0.65			13					
Oldmill Bay	Sand	1 <sup>N</sup>	<0.56	<0.75			2.8					
Ballymacormick	Mud and sand	1 <sup>N</sup>	<1.4	<0.93			8.1					
Ballymacormick	Sand	1 <sup>N</sup>	<0.73	<0.88			1.2					
<b>Strangford Lough</b>												
- Nicky's Point	Mud	2 <sup>N</sup>	<1.1	<1.2			4.0					
Dundrum Bay	Mud	1 <sup>N</sup>	<1.2	<0.84			1.2					
Dundrum Bay	Mud and sand	1 <sup>N</sup>	<1.4	<1.1			2.0					
Carlingford Lough	Mud	2 <sup>N</sup>	<2.0	<1.5	1.8	11	8.7	*	*			

\* Not detected by the method used

<sup>S</sup> Measurements labelled "S" are made on behalf of the Scottish Environment Protection Agency

<sup>N</sup> Measurements labelled "N" are made on behalf of the Northern Ireland Environment Agency  
All other measurements are made on behalf of the Environment Agency

<sup>a</sup> Data for natural radionuclides for some of these samples may be available in Table 7.7

**Table 2.9. Gamma radiation dose rates over areas of the Cumbrian coast and further afield, 2015**

Location	Ground type	No. of sampling observations	Mean gamma dose rate in air at 1m, $\mu\text{Gy h}^{-1}$
<b>Cumbria, Rockcliffe-Harrington</b>			
Rockcliffe Marsh	Salt marsh	2	0.084
Burgh Marsh	Salt marsh	2	0.087
Port Carlisle 1	Mud and sand	3	0.086
Port Carlisle 1	Sand	1	0.085
Port Carlisle 2	Salt marsh	2	0.088
Port Carlisle 2	Grass	2	0.086
Greenend 1	Mud	1	0.092
Greenend 1	Mud and sand	1	0.089
Greenend 1	Sand	2	0.090
Greenend 2	Salt marsh	3	0.088
Greenend 2	Grass	1	0.089
Cardurnock Marsh	Salt marsh	4	0.079
Newton Arlosh	Salt marsh	4	0.093
Silloth harbour	Mud and stones	1	0.099
Silloth harbour	Sand	1	0.10
Silloth harbour	Sand and stones	2	0.10
Silloth silt pond	Grass and sand	1	0.081
Silloth silt pond	Grass	3	0.080
Allonby	Sand	4	0.085
Maryport harbour	Mud and sand	1	0.088
Maryport harbour	Sand	1	0.086
Workington harbour	Sand and shingle	1	0.11
Workington harbour	Pebbles and sand	1	0.11
Harrington harbour	Sand and shingle	1	0.11
Harrington harbour	Pebbles and sand	1	0.11
<b>Cumbria, Whitehaven-Drigg</b>			
Whitehaven - outer harbour	Sand	2	0.092
Whitehaven - outer harbour	Sand and shingle	1	0.096
Whitehaven - outer harbour	Pebbles and sand	1	0.089
St Bees	Sand	3	0.077
St Bees	Sand and shingle	1	0.072
Nethertown beach	Shingle	2	0.12
Braystones	Shingle	2	0.12
Sellafield dunes	Grass	2	0.11
North of former pipeline on foreshore	Sand	2	0.091
South of former pipeline on foreshore	Sand	2	0.090
River Calder downstream of site	Grass	2	0.089
River Calder upstream of site	Stones	1	0.11
River Calder upstream of site	Pebbles and rock	1	0.12
Seascale	Sand	2	0.081
Seascale	Sand and shingle	1	0.10
Seascale	Shingle	1	0.088
Seascale grass / car park	Grass	4	0.085

**Table 2.9. continued**

Location	Ground type	No. of sampling observations	Mean gamma dose rate in air at 1m, $\mu\text{Gy h}^{-1}$
<b>Cumbria, Ravenglass-Askam</b>			
Ravenglass - Carleton Marsh	Salt marsh	3	0.13
Ravenglass - Carleton Marsh	Grass and salt marsh	1	0.14
Ravenglass - River Mite estuary (erosional)	Salt marsh	4	0.14
Ravenglass - Raven Villa	Salt marsh	4	0.13
Ravenglass - boat area	Sand	1	0.11
Ravenglass - boat area	Sand and stones	1	0.11
Ravenglass - boat area	Stones	1	0.11
Ravenglass - boat area	Pebbles and stones	1	0.11
Ravenglass - ford	Mud	1	0.090
Ravenglass - ford	Mud and sand	1	0.10
Ravenglass - ford	Sand	2	0.096
Muncaster Bridge	Salt marsh	3	0.11
Muncaster Bridge	Grass	1	0.12
Ravenglass - salmon garth	Mud and sand	1	0.11
Ravenglass - salmon garth	Sand	1	0.10
Ravenglass - salmon garth	Sand and stones	1	0.11
Ravenglass - salmon garth	Stones	1	0.11
Ravenglass - Eskmeals Nature Reserve	Mud	1	0.12
Ravenglass - Eskmeals Nature Reserve	Salt marsh	3	0.11
Newbiggin/Eskmeals viaduct	Salt marsh	4	0.12
Newbiggin/Eskmeals Bridge	Salt marsh	4	0.12
Tarn Bay	Sand	1	0.075
Tarn Bay	Sand and shingle	1	0.11
Silecroft	Shingle	2	0.12
Haverigg	Mud and sand	1	0.094
Haverigg	Sand	1	0.076
Millom	Sand	2	0.10
Low Shaw	Salt marsh	2	0.082
Askam	Mud	1	0.070
Askam	Sand	1	0.068
<b>Cumbria, Walney-Arnside</b>			
Walney Channel, N of discharge point	Mud	2	0.089
Walney Channel, S of discharge point	Mud	2	0.093
Tummer Hill Marsh	Salt marsh	2	0.11
Roa Island	Mud	1	0.091
Roa Island	Sand	1	0.095
Greenodd Salt Marsh	Salt marsh	1	0.082
Greenodd Salt Marsh	Grass	1	0.078
Sand Gate Marsh	Salt marsh	3	0.088
Sand Gate Marsh	Grass	1	0.083
Kents Bank 2	Salt marsh	3	0.091
Kents Bank 2	Grass	1	0.088
High Foulshaw	Mud	1	0.087
High Foulshaw	Sand and mud	1	0.077
High Foulshaw	Salt marsh	1	0.099
High Foulshaw	Grass and mud	1	0.082
Arnside 1	Mud and sand	2	0.082
Arnside 1	Sand	1	0.094
Arnside 1	Grass and sand	1	0.081
Arnside 2	Salt marsh	4	0.098

Table 2.9. continued

Location	Ground type	No. of sampling observations	Mean gamma dose rate in air at 1m, $\mu\text{Gy h}^{-1}$
<b>Lancashire and Merseyside</b>			
Morecambe Central Pier	Sand	2	0.076
Half Moon Bay	Sand	1	0.079
Half Moon Bay	Sand and stones	1	0.087
Red Nab Point	Sand	2	0.086
Middleton Sands	Sand	2	0.078
Sunderland Point	Mud and sand	4	0.10
Sunderland	Mud and sand	1	0.093
Sunderland	Salt marsh	3	0.091
Colloway Marsh	Salt marsh	4	0.12
Lancaster	Grass	4	0.082
Aldcliffe Marsh	Salt marsh	4	0.092
Conder Green	Mud	1	0.095
Conder Green	Salt marsh	3	0.090
Pilling Marsh	Salt marsh	4	0.098
Knott End	Sand	2	0.078
Heads - River Wyre	Salt marsh	4	0.099
Height o' th' hill - River Wyre	Salt marsh	4	0.11
Hambleton	Salt marsh	4	0.10
Skippool Creek 1	Salt marsh	4	0.11
Skippool Creek 2	Salt marsh	4	0.11
Skippool Creek (board walk)	Wood	4	0.095
Skippool Creek Bivand (board walk)	Salt marsh	1	0.092
Skippool Creek Bivand (board walk)	Wood	3	0.094
Skippool Creek Bivand 1	Mud	3	0.084
Skippool Creek Bivand 1	Mud and salt marsh	1	0.086
Fleetwood Marsh Nature Park	Salt marsh	4	0.11
Fleetwood shore 1	Sand	4	0.078
Fleetwood shore 2	Salt marsh	4	0.12
Blackpool	Sand	2	0.064
Crossens Marsh	Salt marsh	4	0.091
Ainsdale	Sand	4	0.066
Rock Ferry	Mud and sand	4	0.089
New Brighton	Sand	4	0.065
West Kirby	Sand	4	0.071
Little Neston Marsh 1	Salt marsh	2	0.085
Little Neston Marsh 2	Grass	2	0.073
Flint 1	Mud	2	0.090
Flint 2	Salt marsh	2	0.098
<b>Scotland</b>			
Piltanton Burn	Salt marsh	2 <sup>S</sup>	0.070
Garlieston	Mud	2 <sup>S</sup>	0.070
Innerwell	Sediment	2 <sup>S</sup>	0.077
Bladnoch	Mud	2 <sup>S</sup>	0.078
Carsluith	Sand	2 <sup>S</sup>	0.086
Skyreburn Bay (Water of Fleet)	Salt marsh	2 <sup>S</sup>	0.085
Kirkcudbright	Salt marsh	2 <sup>S</sup>	0.067
Cutters Pool	Winkle bed	4 <sup>S</sup>	0.087
Balcary Bay	Sediment	2 <sup>S</sup>	0.089
Gardenburn	Mud	2 <sup>S</sup>	0.081
Palnackie Harbour	Mud	2 <sup>S</sup>	0.076
Kippford - Slipway	Mud	2 <sup>S</sup>	0.11
Kippford - Merse	Mud	2 <sup>S</sup>	0.10
Kirkconnell Marsh	Mud	2 <sup>S</sup>	0.10
Southernness	Sand	2 <sup>S</sup>	0.063
<b>Wales</b>			
Prestatyn	Sand	2	0.068
Rhyl	Salt marsh	2	0.090
Llandudno	Sand and shingle	2	0.096
Caerhun	Salt marsh	2	0.091
Llanfairfechan	Sand and shells	2	0.079

**Table 2.9. continued**

Location	Ground type	No. of sampling observations	Mean gamma dose rate in air at 1m, $\mu\text{Gy h}^{-1}$
<b>Northern Ireland</b>			
Lisahally	Mud	1 <sup>N</sup>	0.071
Donnybrewer	Shingle	1 <sup>N</sup>	0.051
Carrichue	Mud	1 <sup>N</sup>	0.052
Bellerena	Mud	1 <sup>N</sup>	0.055
Benone	Sand	1 <sup>N</sup>	0.058
Castlerock	Sand	1 <sup>N</sup>	0.060
Portstewart	Sand	1 <sup>N</sup>	0.050
Portrush, Blue Pool	Sand	1 <sup>N</sup>	0.058
Portrush, White Rocks	Sand	1 <sup>N</sup>	0.063
Portballintrae	Sand	1 <sup>N</sup>	0.053
Giant's Causeway	Sand	1 <sup>N</sup>	0.056
Ballycastle	Sand	1 <sup>N</sup>	0.056
Cushendun	Sand	1 <sup>N</sup>	0.055
Cushendall	Sand and stones	1 <sup>N</sup>	0.058
Red Bay	Sand	1 <sup>N</sup>	0.062
Carnlough	Sand	1 <sup>N</sup>	0.053
Glenarm	Sand	1 <sup>N</sup>	0.055
Half Way House	Sand	1 <sup>N</sup>	0.053
Ballygally	Sand	1 <sup>N</sup>	0.053
Drains Bay	Sand	1 <sup>N</sup>	0.052
Larne	Sand	1 <sup>N</sup>	0.057
Whitehead	Sand	1 <sup>N</sup>	0.056
Carrickfergus	Sand	1 <sup>N</sup>	0.058
Jordanstown	Sand	1 <sup>N</sup>	0.060
Helen's Bay	Sand	1 <sup>N</sup>	0.059
Groomsport	Sand	1 <sup>N</sup>	0.059
Millisle	Sand	1 <sup>N</sup>	0.072
Ballywalter	Sand	1 <sup>N</sup>	0.061
Ballyhalbert	Sand	1 <sup>N</sup>	0.057
Cloghy	Sand	1 <sup>N</sup>	0.058
Portaferry	Shingle and stones	1 <sup>N</sup>	0.084
Kircubbin	Sand	1 <sup>N</sup>	0.063
Greyabbey	Sand	1 <sup>N</sup>	0.076
Ards Maltings	Mud	1 <sup>N</sup>	0.071
Island Hill	Mud	1 <sup>N</sup>	0.070
Nicky's Point	Mud	1 <sup>N</sup>	0.086
Strangford	Shingle and stones	1 <sup>N</sup>	0.078
Kilclief	Sand	1 <sup>N</sup>	0.080
Ardglass	Mud	1 <sup>N</sup>	0.068
Killough	Mud	1 <sup>N</sup>	0.079
Ringmore Point	Sand	1 <sup>N</sup>	0.078
Tyrella	Sand	1 <sup>N</sup>	0.070
Dundrum	Sand	1 <sup>N</sup>	0.085
Newcastle	Sand	1 <sup>N</sup>	0.078
Annalong	Sand	1 <sup>N</sup>	0.080
Cranfield Bay	Sand	1 <sup>N</sup>	0.070
Mill Bay	Sand	1 <sup>N</sup>	0.090
Greencastle	Sand	1 <sup>N</sup>	0.080
Rostrevor	Sand	1 <sup>N</sup>	0.11
Narrow Water	Mud	1 <sup>N</sup>	0.10

<sup>S</sup> Measurements labelled "S" are made on behalf of the Scottish Environment Protection Agency

<sup>N</sup> Measurements labelled "N" are made on behalf of the Northern Ireland Environment Agency  
All other measurements are made on behalf of the Environment Agency

**Table 2.10. Beta radiation dose rates on contact with fishing gear on vessels operating off Sellafield, 2015**

Vessel or location	Type of gear	No. of sampling observations	Mean beta dose rate in tissue, $\mu\text{Sv h}^{-1}$
101	Nets	2	<0.075
104	Nets	1	<0.076
105	Nets	1	<0.075
106	Nets	2	<0.095
109	Nets	2	<0.092
111	Nets	1	<0.071
112	Nets	1	0.099
113	Nets	1	<0.073
Saltcoats	Pots	1	<0.086
Sellafield area – Batch 1	Pots	1	0.17
Sellafield area – Batch 2	Pots	1	0.10
Drigg/Selker – Batch 1	Pots	1	0.11
Drigg/Selker – Batch 2	Pots	1	<0.088

**Table 2.11. Beta radiation dose rates over intertidal areas of the Cumbrian coast, 2015**

Location	Ground type	No. of sampling observations	Mean beta dose rate in tissue, $\mu\text{Sv h}^{-1}$
Whitehaven – outer harbour	Sand	1	0.21
Whitehaven – outer harbour	Sand and shingle	1	0.10
Whitehaven – outer harbour	Pebbles and sand	1	0.14
St Bees	Sand	2	0.12
St Bees	Sand and shingle	1	0.11
Sellafield pipeline	Sand	1	0.065
Ravenglass – Raven Villa	Salt marsh	4	0.11
Tarn Bay	Sand and shingle	1	0.23

**Table 2.12. Concentrations of radionuclides in aquatic plants from the Cumbrian coast and further afield, 2015**

Location	Material	No. of sampling observations	Mean radioactivity concentration (fresh), Bq kg <sup>-1</sup>							
			<sup>60</sup> Co	<sup>65</sup> Zn	<sup>90</sup> Sr	<sup>95</sup> Zr	<sup>95</sup> Nb	<sup>99</sup> Tc	<sup>106</sup> Ru	<sup>110m</sup> Ag
<b>Cumbria</b>										
Silloth	Seaweed	2	<0.67			<0.44	<0.86	110	<4.2	<0.65
Harrington Harbour	Seaweed	2	<0.61			<0.39	<0.72	240	<3.6	<0.58
St Bees	Seaweed	2	<0.61		0.27	<0.39	<0.72	480	<3.6	<0.56
Sellafield	Seaweed	2	<0.91		<0.42	<0.60	<1.2	1200	<5.0	<0.84
Ravenglass	Samphire	1 <sup>F</sup>	<0.05	<0.11		<0.24	<0.18	0.38	<0.43	<0.07
Ravenglass	Seaweed	2	<0.49			<0.32	<0.59	110	<2.9	<0.48
<b>Lancashire</b>										
Half Moon Bay	Seaweed	2	<0.72			<0.51	<0.97	140	<4.5	<0.80
Marshside Sands	Samphire	1 <sup>F</sup>	<0.05	<0.12		<0.13	<0.14	<0.34	<0.36	<0.07
<b>Scotland</b>										
Aberdeen	<i>Fucus vesiculosus</i>	1 <sup>S</sup>	<0.10	<0.19		<0.22	<0.21	9.2	<0.55	<0.10
Lerwick	<i>Fucus vesiculosus</i>	1 <sup>S</sup>	<0.10	<0.37		<2.4	<0.88	1.7	<0.96	<0.12
Kinlochbervie	<i>Fucus vesiculosus</i>	2 <sup>S</sup>	<0.10	<0.21		<0.70	<0.37	12	<0.62	<0.10
Lewis	<i>Fucus vesiculosus</i>	1 <sup>S</sup>	<0.10	<0.17		<0.42	<0.26	8.0	<0.46	<0.10
Islay	<i>Fucus vesiculosus</i>	1 <sup>S</sup>	<0.10	<0.10		<0.19	<0.14	16	<0.28	<0.10
Campbeltown	<i>Fucus vesiculosus</i>	1 <sup>S</sup>	<0.10	<0.18		<0.42	<0.30	61	<0.58	<0.10
Port William	<i>Fucus vesiculosus</i>	4 <sup>S</sup>	<0.10	<0.18		<0.34	<0.25	53	<0.52	<0.10
Garlieston	<i>Fucus vesiculosus</i>	4 <sup>S</sup>	<0.10	<0.18		<0.35	<0.23	53	<0.52	<0.12
Auchencairn	<i>Fucus vesiculosus</i>	4 <sup>S</sup>	<0.10	<0.20		<0.38	<0.28	290	<0.58	<0.11
<b>Isle of Man<sup>b</sup></b>										
	Seaweed	3	<0.84			<0.59	<1.1	58	<5.0	<0.85
<b>Wales</b>										
Cemaes Bay	Seaweed	2	<0.46			<0.30	<0.57	13	<2.7	<0.45
Porthmadog	Seaweed	2	<0.46			<0.33	<0.62	1.5	<3.0	<0.47
Lavernock Point	Seaweed	2	<0.80			<0.53	<1.0	<0.61	<4.5	<0.85
Fishguard	Seaweed	2	<0.87			<0.56	<1.1	4.1	<5.1	<0.88
<b>Northern Ireland</b>										
Portrush	<i>Fucus</i> spp.	4 <sup>N</sup>	<0.03	<0.09		<0.06	<0.07		<0.27	<0.05
Portaferry <sup>a</sup>	<i>Rhodomenia</i> spp.	4 <sup>N</sup>	<0.07	<0.18		<0.16	<0.18	0.26	<0.57	<0.10
Ardglass	<i>Fucus vesiculosus</i>	1 <sup>N</sup>	<0.04	<0.12		<0.15	<0.14		<0.35	<0.06
Ardglass	<i>Ascophyllum nodosum</i>	3 <sup>N</sup>	<0.06	<0.16		<0.10	<0.13	42	<0.46	<0.08
Carlingford Lough	<i>Fucus vesiculosus</i>	1 <sup>N</sup>	<0.05	<0.14		<0.07	<0.11		<0.39	<0.07
Carlingford Lough	<i>Ascophyllum nodosum</i>	3 <sup>N</sup>	<0.09	<0.23		<0.19	<0.21	77	<0.64	<0.11

Table 2.12. continued

Location	Material	No. of sampling observations	Mean radioactivity concentration (fresh), Bq kg <sup>-1</sup>							
			<sup>125</sup> Sb	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>144</sup> Ce	<sup>155</sup> Eu	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am
<b>Cumbria</b>										
Silloth	Seaweed	2	<2.4	<0.58	3.0	<1.7				2.8
Harrington Harbour	Seaweed	2	<2.1	<0.52	1.9	<1.5				2.2
St Bees	Seaweed	2	<2.2	<0.53	1.6	<1.6		0.95	4.7	2.2
Sellafield	Seaweed	2	<2.9	<0.75	2.4	<1.8		0.99	4.4	1.8
Ravenglass	Samphire	1 <sup>F</sup>	<0.11	<0.05	0.37	<0.19	<0.07			1.0
Ravenglass	Seaweed	2	<1.8	<0.43	3.3	<1.4				10
<b>Lancashire</b>										
Half Moon Bay	Seaweed	2	<2.7	<0.67	3.1	<1.8				<1.6
Marshside Sands	Samphire	1 <sup>F</sup>	<0.10	<0.04	0.38	<0.17	<0.07			0.18
<b>Scotland</b>										
Aberdeen	<i>Fucus vesiculosus</i>	1 <sup>S</sup>	<0.15	<0.10	<0.10	<0.35	<0.16			<0.10
Lerwick	<i>Fucus vesiculosus</i>	1 <sup>S</sup>	<0.24	<0.10	<0.10	<0.64	<0.24			<0.17
Kinlochbervie	<i>Fucus vesiculosus</i>	2 <sup>S</sup>	<0.17	<0.10	0.12	<0.41	<0.16			<0.10
Lewis	<i>Fucus vesiculosus</i>	1 <sup>S</sup>	<0.12	<0.10	0.23	<0.29	<0.12			<0.10
Islay	<i>Fucus vesiculosus</i>	1 <sup>S</sup>	<0.10	<0.10	<0.10	<0.18	<0.10			<0.10
Campbeltown	<i>Fucus vesiculosus</i>	1 <sup>S</sup>	<0.15	<0.10	0.20	<0.33	<0.14			<0.10
Port William	<i>Fucus vesiculosus</i>	4 <sup>S</sup>	<0.14	<0.10	0.43	<0.34	<0.15			<0.36
Garlieston	<i>Fucus vesiculosus</i>	4 <sup>S</sup>	<0.12	<0.10	3.1	<0.34	<0.14			5.6
Auchencairn	<i>Fucus vesiculosus</i>	4 <sup>S</sup>	<0.15	<0.10	2.0	<0.37	<0.15			3.0
<b>Isle of Man<sup>b</sup></b>										
	Seaweed	3	<2.9	<0.78	<0.62	<2.1	<0.97			<0.68
<b>Wales</b>										
Cemaes Bay	Seaweed	2	<1.6	<0.41	<0.36	<1.4				<0.50
Porthmadog	Seaweed	2	<1.7	<0.45	<0.38	<1.4				<0.49
Lavernock Point	Seaweed	2	<2.6	<0.64	<0.56	<1.8	<0.88			<0.59
Fishguard	Seaweed	2	<2.9	<0.76	<0.64	<2.3				<0.70
<b>Northern Ireland</b>										
Portrush	<i>Fucus</i> spp.	4 <sup>N</sup>	<0.08	<0.04	<0.04	<0.15	<0.07			<0.08
Portaferry <sup>a</sup>	<i>Rhodomenia</i> spp.	4 <sup>N</sup>	<0.15	<0.07	0.41	<0.28	<0.12	0.064	0.38	0.70
Ardglass	<i>Fucus vesiculosus</i>	1 <sup>N</sup>	<0.09	<0.05	0.43	<0.17	<0.08			0.24
Ardglass	<i>Ascophyllum nodosum</i>	3 <sup>N</sup>	<0.12	<0.06	0.18	<0.24	<0.12			<0.12
Carlingford Lough	<i>Fucus vesiculosus</i>	1 <sup>N</sup>	<0.11	<0.06	0.28	<0.19	0.31			<0.06
Carlingford Lough	<i>Ascophyllum nodosum</i>	3 <sup>N</sup>	<0.18	<0.09	0.23	<0.37	<0.20			<0.24

<sup>a</sup> <sup>242</sup>Cm and <sup>243+244</sup>Cm were not detected by the method used

<sup>b</sup> Data for natural radionuclides for some of these samples may be available in Table 7.7

<sup>F</sup> Measurements labelled "F" are made on behalf of the Food Standards Agency

<sup>N</sup> Measurements labelled "N" are made on behalf of the Northern Ireland Environment Agency

<sup>S</sup> Measurements labelled "S" are made on behalf of the Scottish Environment Protection Agency

All other measurements are made on behalf of the Environment Agency

**Table 2.13. Concentrations of radionuclides in terrestrial food and the environment near Ravenglass, 2015**

Material and selection <sup>a</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>										
		<sup>3</sup> H	<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>95</sup> Nb	<sup>95</sup> Zr	<sup>99</sup> Tc	<sup>106</sup> Ru	<sup>125</sup> Sb	<sup>129</sup> I	<sup>134</sup> Cs
Milk	3	<3.1	18	<0.06	0.028	<0.06	<0.09	<0.038	<0.44	<0.13	<0.0037	<0.06
Milk max					0.043	<0.07	<0.10		<0.47	<0.15	<0.0040	<0.07
Beef kidney	1	<2.6	12	<0.19	<0.046	<0.30	<0.39	<0.31	<1.6	<0.48		<0.21
Beef liver	1	<3.2	29	<0.07	<0.052	<0.09	<0.15	<0.59	<0.59	<0.11	<0.017	<0.07
Beef muscle	1	<2.6	22	<0.09	<0.043	<0.07	<0.10	<0.23	<0.34	<0.16	<0.024	<0.07
Sheep muscle	2	<5.3	28	<0.04	<0.043	<0.12	<0.12	<0.26	<0.70	<0.15	<0.017	<0.07
Sheep muscle max		<6.3		<0.06	<0.044	<0.14	<0.13	<0.27	<0.74	<0.17	<0.019	<0.08
Sheep offal	2	<6.4	32	<0.04	<0.044	<0.06	<0.09	<0.29	<0.33	<0.12	<0.017	<0.05
Sheep offal max		<7.2	34	<0.05	<0.047	<0.08	<0.11	<0.36	<0.40			

Material and selection <sup>a</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>									
		<sup>137</sup> Cs	Total Cs	<sup>144</sup> Ce	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Pu	<sup>241</sup> Am
Milk	3	<0.09		<0.32				<0.000042	<0.000020	<0.12	<0.000041
Milk max		<0.12		<0.37				<0.000076	<0.000033	<0.14	0.000067
Beef kidney	1	5.8	5.8	<0.92	0.012	0.0015	0.0094	<0.000052	0.00022	<0.23	0.00064
Beef liver	1	3.3	3.2	<0.43				0.00050	0.0030	<0.38	0.0045
Beef muscle	1	3.9	4.4	<0.48				<0.000027	<0.000047	<0.25	0.00020
Sheep muscle	2	1.1	1.3	<0.30				<0.000020	0.000046	<0.19	<0.000071
Sheep muscle max		1.5	1.7	<0.32				<0.000022	0.000058	<0.20	0.000089
Sheep offal	2	0.73	1.2	<0.26				<0.000068	0.00054	<0.20	0.00045
Sheep offal max		0.86	1.5	<0.33				0.000088	0.00070	<0.22	0.00056

<sup>a</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima.

If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>b</sup> Except for milk where units are Bq l<sup>-1</sup>

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

**Table 2.14. Concentrations of radionuclides in surface waters from West Cumbria, 2015**

Location	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>								
		<sup>3</sup> H	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	Gross alpha	Gross beta
Ehen Spit beach <sup>a</sup>	4	150	<0.31	<0.035	<0.31	<0.26	<0.0028	<0.0037	<3.1	10
River Ehen (100m downstream of sewer outfall)	4	<7.4	<0.27	<0.11	<0.28	<0.23	<0.0028	<0.0015	<0.075	0.49
River Calder (downstream)	4	<3.4	<0.22	<0.055	<0.23	<0.19	<0.0028	<0.0012	<0.021	0.15
River Calder (upstream)	4	<3.5	<0.26	<0.027	<0.27	<0.21	<0.0031	<0.0018	<0.017	0.070
Wast Water	1	<4.0	<0.33			<0.32			<0.030	0.069
Ennerdale Water	1	<4.1	<0.11		<0.11	<0.09			<0.033	0.039
Devoke Water	1	<3.3	<0.09		<0.09	<0.08			<0.028	0.030
Thirlmere	1	<3.1	<0.30			<0.24			<0.026	0.058

<sup>a</sup> The concentration of <sup>99</sup>Tc was <0.15 Bq l<sup>-1</sup>

**Table 2.15. Concentrations of radionuclides in road drain sediments from Whitehaven and Seascale, 2015**

Location	No. of sampling observations	Mean radioactivity concentration (dry), Bq kg <sup>-1</sup>						
		<sup>60</sup> Co	<sup>90</sup> Sr	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>238</sup> Pu	<sup>239</sup> Pu+ <sup>240</sup> Pu	<sup>241</sup> Am
Seascale SS 204	1	<1.5	<2.0	<1.4	140	2.9	22	32
Seascale SS 233	1	<1.7	<2.0	<1.7	170	3.2	25	28
Seascale SS 209	1	<0.42	<2.4	<0.37	10	1.1	8.5	13
Seascale SS 232	1	<1.2	<2.0	<1.1	34	4.6	28	37
Seascale SS 231	1	<0.34	<2.0	<0.32	30	6.4	39	57
Whitehaven SS 201	1	<3.0	<2.5	<3.0	21	<0.33	1.4	2.4

**Table 2.16. Doses from artificial radionuclides in the Irish Sea, 2007-2015**

Group	Exposure, mSv per year									
	2007	2008	2009	2010	2011	2012	2013	2014	2015	
Northern Ireland	0.015	0.017	0.012	0.010	0.010	0.011	0.010	0.009	0.008	
Dumfries and Galloway	0.060	0.047	0.047	0.040	0.040	0.046	0.044	0.045	0.037	
Whitehaven	0.009	0.009	0.011	0.010	0.010	0.013	0.010	0.012	0.017	
Sellafield (5 year average consumption)	0.24	0.23	0.20	0.18	0.15	0.14	0.12	0.089	0.084	
Morecambe Bay	0.037	0.042	0.041	0.046	0.034	0.034	0.036	0.032	0.031	
North Wales	0.014	0.018	0.015	0.013	0.014	0.014	0.013	0.018	0.014	

**Table 2.17. Individual radiation exposures, Sellafield, 2015**

Representative person <sup>a</sup>	Exposure, mSv per year							
	Total	Seafood (nuclear industry discharges) <sup>h</sup>	Seafood (other discharges) <sup>i</sup>	Other local food	External radiation from intertidal areas, river banks or fishing gear	Intakes of sediment and water	Gaseous plume related pathways	Direct radiation from site
<b>Total dose – maximum effect of all sources</b>								
<b>Adult mollusc consumers</b>	<b>0.42<sup>e</sup></b>	<b>0.061</b>	<b>0.35</b>	<b>-</b>	<b>0.017</b>	<b>-</b>	<b>-</b>	<b>-</b>
<b>Total dose – maximum effect of gaseous release and direct radiation sources</b>								
<b>Local child inhabitants (0 - 0.25km)</b>	<b>0.008<sup>b</sup></b>	<b>-</b>	<b>-</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>-</b>	<b>&lt;0.005</b>	<b>0.005</b>
<b>Total dose – maximum effect of liquid release source</b>								
<b>Adult mollusc consumers</b>	<b>0.42<sup>e</sup></b>	<b>0.061</b>	<b>0.35</b>	<b>-</b>	<b>0.017</b>	<b>-</b>	<b>-</b>	<b>-</b>
<b>Source specific doses</b>								
<b>Seafood consumers</b>								
Local seafood consumers (habits averaged 2011-15)	0.37 <sup>f</sup>	0.051	0.29	-	0.033	-	-	-
Local seafood consumers (habits for 2015)	0.52 <sup>g</sup>	0.061	0.42	-	0.037	-	-	-
Whitehaven – seafood consumers	0.017	0.017	-	-	-	-	-	-
Dumfries and Galloway – seafood and wildfowl consumers	0.038	0.030	-	-	0.009	-	-	-
Morecambe Bay – seafood consumers	0.031	0.012	-	-	0.019	-	-	-
Northern Ireland – seafood consumers	0.008	0.007	-	-	<0.005	-	-	-
North Wales – seafood consumers	0.014	0.007	-	-	0.007	-	-	-
<b>Other groups</b>								
Ravenglass Estuary, marsh users	0.013	-	-	-	0.010	<0.005	-	-
Fishermen – handling nets or pots <sup>c</sup>	0.10	-	-	-	0.10	-	-	-
Bait diggers and shellfish collectors <sup>c</sup>	0.064	-	-	-	0.064	-	-	-
Ribble Estuary – houseboats	0.054	-	-	-	0.054	-	-	-
Barrow Houseboats	0.050	-	-	-	0.050	-	-	-
Local infant consumers of locally grown food at Ravenglass	0.017 <sup>b</sup>	-	-	0.017	-	-	-	-
Local infant consumers of locally grown food at LLWR near Drigg	<0.005 <sup>b</sup>	-	-	<0.005	-	-	-	-
Infant inhabitants and consumers of locally grown food	0.012 <sup>b</sup>	-	-	0.012	-	-	<0.005	-
<b>Groups with average consumption or exposure</b>								
Average seafood consumer in Cumbria	<0.005	<0.005	-	-	-	-	-	-
Average consumer of locally grown food <sup>d</sup>	<0.005	-	-	<0.005	-	-	-	-
Typical visitor to Cumbria	<0.005	<0.005	<0.005	-	<0.005	-	-	-
<b>Recreational user of beaches</b>								
Dumfries and Galloway	0.006	-	-	-	0.006	-	-	-
North Cumbria	0.011	-	-	-	0.011	-	-	-
Sellafield	0.011	-	-	-	0.011	-	-	-
Lancashire	0.007	-	-	-	0.007	-	-	-
North Wales	0.008	-	-	-	0.008	-	-	-
<b>Recreational user of mud/saltmarsh areas</b>								
Dumfries and Galloway	<0.005	-	-	-	<0.005	-	-	-
North Cumbria	0.005	-	-	-	0.005	-	-	-
Sellafield	0.013	-	-	-	0.013	-	-	-
Lancashire	0.006	-	-	-	0.006	-	-	-
North Wales	0.005	-	-	-	0.005	-	-	-

<sup>a</sup> The total dose is the dose which accounts for all sources including gaseous and liquid discharges and direct radiation.

The total dose for the representative person with the highest dose presented.

Other dose values are presented for specific sources, either liquid discharges or gaseous discharges, and their associated pathways.

They serve as a check on the validity of the total dose assessment.

The representative person is an adult unless otherwise stated

<sup>b</sup> Includes a component due to natural sources of radionuclides

<sup>c</sup> Exposure to skin for comparison with the 50 mSv dose limit

<sup>d</sup> Only the adult age group is considered for this assessment

<sup>e</sup> The dose due to nuclear industry discharges was 0.078 mSv

<sup>f</sup> The dose due to nuclear industry discharges was 0.084 mSv

<sup>g</sup> The dose due to nuclear industry discharges was 0.098 mSv

<sup>h</sup> May include a small contribution from LLWR near Drigg

<sup>i</sup> Enhanced naturally occurring radionuclides from Whitehaven



## 3. Research establishments

This section considers the results of monitoring by the Environment Agency, FSA and SEPA near research establishments that hold nuclear site licences.

NDA has ownership of the majority of such sites, with licensed nuclear sites at Harwell and Winfrith in England, and Dounreay in Scotland. Previously Harwell, Winfrith and Dounreay sites were operated by UKAEA. In 2009, Research Sites Restoration Limited (RSRL) and Dounreay Site Restoration Limited (DSRL) (both wholly-owned subsidiaries of UKAEA) became the site licence companies for Harwell and Winfrith, and Dounreay respectively. UKAEA Limited itself was sold to Babcock International Group plc, including its subsidiary companies DSRL and RSRL, as a preliminary to NDA starting the Dounreay Parent Body Organisation (PBO) competition.

In 2012, Babcock Dounreay Partnership (BDP), which has subsequently been renamed as the Cavendish Dounreay Partnership, was awarded the contract to manage the decommissioning and clean-up of the Dounreay site, and became the PBO for Dounreay. In September 2014, NDA formally appointed Cavendish Fluor Partnership (CFP) as the new PBO for RSRL (and Magnox Limited).

In November 2014, ONR received an application to re-license the Harwell and Winfrith sites into a single site licence company alongside the ten Magnox sites. As of 1st April 2015, Harwell and Winfrith sites, previously operated by RSRL, merged to be part of Magnox Limited.

All of the nuclear licensed sites have reactors that are at different stages of decommissioning. Discharges of radioactive waste are largely related to decommissioning and decontamination operations and the nuclear related research that is undertaken. Tenants, or contractors, such as Nuvia Limited carry out some of this work.

Regular monitoring of the environment was undertaken in relation to all sites, which included the effects of discharges from neighbouring sites and tenants where appropriate, e.g. the Vulcan Naval Reactor Test Establishment (NRTE) adjacent to the Dounreay site.

The medium-term trends in discharges, environmental concentrations and doses at Dounreay, Harwell and Winfrith have been considered in a summary report (Environment Agency, FSA, NIEA and SEPA, 2010b).

Other minor research sites considered in this section are the non-nuclear site at Culham, Oxfordshire and the Imperial College Reactor Centre near Ascot, Berkshire.

### Key points

- *Total doses* for the representative person were less than 2 per cent of the dose limit, for sites that were assessed
- Harwell and Winfrith sites were re-licensed to Magnox Limited in 2015
- Doses, discharges, environmental concentrations and dose rates in 2015 were broadly similar to those in recent years

#### Dounreay, Highland

- *Total dose* for the representative person decreased in 2015

#### Harwell, Oxfordshire

- *Total dose* for the representative person was similar to that in 2015
- Liquid discharges to Lydebank Brook and to the sewer decreased in 2015

#### Winfrith, Dorset

- *Total dose* for the representative person increased in 2015
- Liquid discharges of alpha-emitting radionuclides were the lowest reported value, and tritium increased, in 2015

### 3.1 Dounreay, Highland



The Dounreay site was opened in 1955 to develop research reactors. Three reactors were built on the site; the Prototype Fast Reactor, the Dounreay Fast Reactor and the Dounreay Materials Test Reactor. All three are now

closed and undergoing decommissioning. It is currently expected that final remediation of the site will be achieved by the interim end state date in 2029 (NDA, 2016b).

From 2005, NDA became responsible for the UK's civil nuclear liabilities which included those at UKAEA Dounreay, and UKAEA became a contractor to NDA. In common with other NDA sites, UKAEA prepared a long term decommissioning plan known as the Lifetime Plan.

NDA's Strategy includes a summary of the PBO competition process. Part of this process required the transfer of the three existing radioactive waste disposal authorisations from UKAEA to a new site licence company (Dounreay Site Restoration Limited, DSRL), before DSRL took over the site management contract. In 2012, Babcock Dounreay Partnership (BDP), which has subsequently been renamed as the Cavendish Dounreay Partnership, was awarded the contract to manage the decommissioning and clean-up of the Dounreay site, and became the PBO for Dounreay.

In 2013, SEPA granted DSRL's authorisation for a Low Level Radioactive Waste disposal facility adjacent to the site. The facility began accepting waste for disposal in April 2015.

During 2015, DSRL notified SEPA of two issues relating to compliance reporting with respect to the monthly disposal reports which are submitted to SEPA. As a result, SEPA has written to DSRL conveying its expectation that DSRL will undertake a detailed review of the site's arrangements for the compilation and checking of the monthly disposal reports as part of DSRL's Environmental Improvement Programme.

In July 2015, DSRL notified SEPA of the discovery of holes in the ventilation ducting associated with the on-site laboratories. As the holes within the ducting were downstream of any abatement, no additional radioactivity would have been discharged to the environment. It was subsequently established that the ventilation system was in an extremely poor condition. As a result, SEPA sent a Warning Letter to DSRL in relation to the failure to keep the ventilation system in good repair and associated failures in the maintenance and inspection regime, and the failure to review the implications on existing infrastructure when delays to programmes occur.

In 2015, radioactive waste discharges from Dounreay were made by DSRL under an authorisation granted by SEPA. The quantities of both gaseous and liquid discharges were generally similar to those in 2014 (Appendix 2).

The most recent habits survey was conducted in 2013 (Papworth *et al.*, 2014).

## Doses to the public

The *total dose* from all pathways and sources of radiation was 0.010 mSv (Table 3.1), or 1 per cent of the dose limit, and down from 0.012 mSv in 2014. The representative person was a local adult who spends a large amount of time over sediments, and a change from that in 2014 (an adult consuming wild fruit and nuts at high-rates). The decrease in *total dose* and a change in the representative person was mostly due to a lower estimate of direct radiation from the site in 2015.

The trend in *total dose* over the period 2004 – 2015 is given in Figure 3.1. The variations in the earlier years were

due to changes in caesium-137 concentrations in game meat and the type of game sampled, but *total doses* were low. The change in *total dose* from 2013 was mostly due to the contribution of goats' milk not being included in the assessment (which has been assessed prior to 2013), as milk samples have not been available in most recent years.


Source specific assessments for external pathways (both for Geo occupants, who regularly visit Oigin's Geo, and fishermen), give exposures of less than the *total dose* in 2015 (Table 3.1). In 2015, the dose to a consumer of terrestrial foodstuffs was 0.014 mSv or approximately 1 per cent of the dose limit for members of the public of 1 mSv. Adults were identified as the most exposed age group (as in previous years). The overall decrease in dose from 0.016 mSv (in 2014) was mostly due to a lower caesium-137 concentration in honey sample in 2015. The dose to a consumer of fish and shellfish, including external exposure from occupancy over local beaches, was 0.011 mSv. The increase in dose from less than 0.005 mSv (in 2014) was mostly due to higher gamma dose rates over sand in 2015.

## Gaseous discharges and terrestrial monitoring

DSRL is authorised by SEPA to discharge radioactive gaseous wastes to the local environment via stacks to the atmosphere. Monitoring conducted in 2015 included the sampling of air, freshwater, grass, soil and locally grown terrestrial foods including meat and vegetables as well as wild foods. As there are no dairy cattle herds in the Dounreay area, no milk samples were collected from cattle. Goats' milk samples (which have been analysed in previous years) were not sampled, as no milk sample was available in 2015. The sampling locations for the terrestrial (and marine) monitoring programmes are shown in Figure 3.2 (Dounreay) and Figure 3.3 (north of Scotland). The results for terrestrial samples and radioactivity in air are given in Tables 3.2(a) and (c) and generally show low concentrations of radioactivity. In 2015, low concentrations of caesium-137, strontium-90, cobalt-60, europium-155, uranium, plutonium and americium-241 are reported in most samples (many reported as less than values). The maximum caesium-137 concentrations in honey and wild mushrooms (7 and 3.1 Bq kg<sup>-1</sup>, respectively) were positively detected again in 2015, but were much lower than the values (33 and 18 Bq kg<sup>-1</sup>, respectively) reported in 2014. Activity concentrations in air samples at locations near to the site are reported as, or close to, less than values.

## Liquid waste discharges and aquatic monitoring

Low level liquid waste is routed via a Low Level Liquid Effluent Treatment Plant (LLLETP). The effluent is discharged to sea (Pentland Firth) via a pipeline terminating 600 metres offshore at a depth of about 24 metres. The



**Figure 3.1.** Total dose at research establishments, 2004–2015  
 (Small doses less than or equal to 0.005 mSv are recorded as being 0.005 mSv)




Figure 3.2. Monitoring locations at Dounreay, 2015 (not including farms)




Figure 3.3. Monitoring locations and concentrations of technetium-99 in seaweed in the north of Scotland, 2015 (not including farms)

discharges also include groundwater pumped from the Dounreay Shaft, surface water runoff, leachate from the low level solid waste disposal facility, and a minor contribution from the adjoining reactor site (Vulcan NRTE), which is operated by the MoD's Defence Equipment and Support organisation.

Routine marine monitoring included sampling of seafood, around the Dounreay outfall in the North Atlantic, and other materials further afield from the outfall, as well as the measurement of beta and gamma dose rates. Seafood samples from within the zone covered by a FEPA\* Order are collected under consent granted in 1997 by the Scottish Office.

Crabs were sampled from the outfall area, together with mussels and winkles from areas along the coastline. Additionally, seawater, sediment and seaweed were sampled as indicator materials. The results for marine samples, and gamma and beta dose rates, are given in Tables 3.2(a) and (b). Activity concentrations were generally low in 2015 and generally similar to those in recent years. In 2015, plutonium radionuclides and americium-241 concentrations were elevated in cod, in comparison to those in recent years. Overall gamma dose rates over sand and soil were generally similar in 2015 (in comparison to 2014), although higher dose rates were measured over sand at Sandside Bay and Castletown Harbour. Technetium-99 concentrations in seaweed remained at the expected levels for this distance from Sellafield and were similar to those in 2014. Figure 3.3 also gives time trend information for technetium-99 concentrations (from Sellafield) in seaweed at Sandside Bay (location shown in Figure 3.2), Kinlochbervie and Burwick. They show an overall decline in concentrations over the period at all three locations. Beta dose measurements are reported as less than values (Table 3.2(b)).


During 2015, DSRL continued vehicle-based monitoring of local public beaches for radioactive fragments in compliance with the requirements of the authorisation granted by SEPA. In 2015, 16 fragments were recovered from Sandside Bay and 8 from the Dounreay foreshore. The caesium-137 activity measured in the fragments recovered from Sandside Bay ranged between 5.5 kBq and 130 kBq (similar to ranges observed in recent years). During beach monitoring on the west foreshore, the presence of caesium-137 contamination was detected on a piece of natural flagstone and on a stone, in January 2016 and February 2016, respectively. Both the flagstone and the stone were recovered and brought onto the Dounreay site. It is believed that the contamination on these items is associated with residual historic contamination from the Castlegate Seep.

\* The FEPA Order was made in 1997 following the discovery of fragments of irradiated nuclear fuel on the seabed near Dounreay, by UKAEA, and prohibits the harvesting of seafoods within a 2 km radius of the discharge pipeline.

The previously conducted offshore survey work provided data on repopulation rates of particles to areas of the seabed previously cleared of particles. This work has improved the understanding of particle movements in the marine environment. The Dounreay Particles Advisory Group (DPAG) completed its work following the production of its Fourth Report (DPAG, 2008). Since the work of DPAG was concluded, the Particles Retrieval Advisory Group (Dounreay) (PRAG (D)) has published reports in March 2010 and March 2011 (PRAG (D), 2010; 2011). In March 2016, PRAG(D) published a further report into the retrieval of offshore particles. This was produced following an extensive research and monitoring programme in 2012 (PRAG (D), 2016). The report considers the extent and effectiveness of the offshore recovery programme to reduce the numbers of particles. The report concludes that any noticeable change in the rate or radioactive content of the particles arriving on the nearest public beach (Sandside Bay) will take a number of years to assess and recommends that in the interim the monitoring of beach local beaches be continued.

In 2007, FSA reviewed the Dounreay FEPA Order. A risk assessment, that was peer-reviewed by PHE, indicated that the food chain risk was very small (FSA, 2009). The FEPA Order was reviewed with regard to ongoing work to remove radioactive particles from the seabed and the food chain risk. In 2009, FSA in Scotland announced that the FEPA Order would remain in place, and be reviewed again when the seabed remediation work was complete.

### 3.2 Harwell, Oxfordshire



The site at Harwell was established in 1946 as Britain's first Atomic Energy Research Establishment and is situated approximately 5 km southwest of the town of Didcot. The Harwell nuclear licensed site forms part of Harwell

Oxford, a science, innovation and business campus. The nuclear licensed site originally accommodated five research reactors of various types. Two of the reactors have been completely removed, and the fuel has been removed from the remaining three reactors. Decommissioning at the Harwell site is well underway. It is expected that decommissioning of all redundant buildings on the site will be completed by 2027 (DECC and NDA, 2014). Final site clearance is expected to be achieved by 2064

† DPAG was set up in 2000, and PRAG (D) thereafter, to provide independent advice to SEPA and UKAEA on issues relating to the Dounreay fragments.

(NDA, 2016b). Since 1st April 2015, the Harwell site has been operated by Magnox Limited on behalf of the NDA.

In September 2015, a habits survey was conducted to determine the consumption and occupancy rates by members of the public (Clyne *et al.*, 2016a). Small decreases in both the fish and crustacean consumption rates have been observed, together with an increase in the occupancy rate, in comparison with those of the previous survey in 2007. Revised figures for consumption rates, together with occupancy rates, are provided in Appendix 1 (Table X2.2).

## Doses to the public

The *total dose* from all pathways and sources of radiation was 0.017 mSv in 2015 (Table 3.1), which was less than 2 per cent of the dose limit, and similar to the value of 0.016 mSv in 2014. The dominant contribution to this dose was direct radiation from the site and the representative person was an adult living near to the site, and was a change from that in 2014 (a prenatal child). The trend in *total dose* over the period 2004 – 2015 is given in Figure 3.1. The *total doses* remained broadly similar from year to year, and were low.

Source specific assessments for a high-rate consumer of terrestrial foods, and for an angler, give exposures that were less than the *total dose* (Table 3.1).

## Gaseous discharges and terrestrial monitoring

Gaseous wastes are discharged via stacks to the local environment. The monitoring programme sampled milk and fruit. Sampling locations at Harwell and in other parts of the Thames catchment are shown in Figure 3.4. The results of the terrestrial monitoring programme in 2015 are shown in Table 3.3(a). The results for tritium analyses in terrestrial samples are reported as less than values.

## Liquid waste discharges and aquatic monitoring

Regulated discharges from Harwell to the River Thames at Sutton Courtenay ceased in 2013 and these wastes are now discharged to sewers serving the Didcot STW; treated effluent subsequently enters the River Thames at Long Wittenham. RSRL completed the decommissioning of the treated waste effluent discharge point at Sutton Courtenay in 2014. Discharges of surface water effluent from the Harwell site are made via the Lydebank Brook, north of the site, which is a permitted route. Discharges to both the Lydebank Brook and the sewer decreased, in comparison to releases in 2014. Figure 3.5 shows trends of discharges over time (2000 – 2015) for cobalt-60 and caesium-137. There was an overall reduction in the discharges over the

whole period, particularly for cobalt-60. Caesium-137 discharges in 2015 were the lowest releases for many years.

The aquatic monitoring programme is directed at consumers of freshwater fish and occupancy close to the liquid discharge point. Tritium and cobalt-60 concentrations in all aquatic samples, and caesium-137 concentrations in fish and freshwater, are reported as less than values. The concentrations of all radionuclides in flounder from the lower reaches of the Thames (from Beckton) are reported as less than values. Caesium-137 concentrations in sediments continued to be enhanced above background levels (excluding those close to Sutton Courtenay, reported as a less than value) in 2015, but were small in terms of any radiological effect. In 2015, sediments were also analysed for iodine-125 (not undertaken in recent years). Iodine-125 was positively detected in samples collected from Appleford, Sutton Courtenay and Day's Lock. The reported values are believed to originate from the therapeutic use of this radionuclide in a local hospital. Concentrations of transuranic elements in sediments are mostly reported as less than values. In 2015, gamma dose rates (where comparisons can be made) were generally higher than those in 2014. However, the increases were small and had no significant effect on doses to the representative person.

## 3.3 Winfrith, Dorset



The Winfrith site is located near Winfrith Newburgh. It was established by in 1957 as an experimental reactor research and development site. At various times there have been nine research and development reactors. The last

operational reactor at Winfrith closed in 1995. Seven of the reactors have been decommissioned and dismantled. It is expected that final decommissioning of both remaining reactors (which commenced in 2014) be completed by 2023 and the site to be de-licensed by 2050 (NDA, 2016b). Since 1st April 2015, the Winfrith site has been operated by Magnox Limited on behalf of the NDA. The most recent habits survey undertaken for Winfrith was in 2003 (McTaggart *et al.*, 2004b).

## Doses to the public

In 2015, the *total dose* from all pathways and sources of radiation was 0.014 mSv (Table 3.1), or approximately 1 per cent of the dose limit, and up from less than




Figure 3.4. Monitoring locations at Thames sites, 2015 (not including farms)





Figure 3.5. Trends in liquid discharges of caesium-137 and cobalt-60 from Harwell, Oxfordshire 2000-2015

0.005 mSv (in 2014). The representative person was an adult living near the site (as opposed to an adult consuming fish at high-rates in 2014). The increase in *total dose* was due to a higher estimate of direct radiation from the site in 2015. Trends in *total doses* in the area of the south coast (and the Severn Estuary) over time are shown in Figure 6.1. At Winfrith, *total doses* remained broadly similar from year to year, and were generally very low.

Source specific assessments for a high-rate consumer of locally grown food, and of fish and shellfish, give exposures that were less than 0.005 mSv in 2015 (Table 3.1).

### Gaseous discharges and terrestrial monitoring

Gaseous radioactive waste is discharged via various stacks to the local environment. Discharges of radioactive wastes continued in 2015 at very low rates. The focus of the terrestrial sampling was for the content of tritium and carbon-14 in milk and crops. Local freshwater samples were also analysed. Sampling locations at Winfrith are shown in Figure 3.6. Data for 2015 are given in Table 3.4(a). Results for terrestrial samples provide little indication of an effect due to gaseous discharges.



**Figure 3.6.** Monitoring locations at Winfrith, 2015 (not including farms)

Carbon-14 was detected in locally produced milk, just above expected background concentrations. Low concentrations of tritium were found in surface water to the north of the site, similar to those in previous years. In all cases, tritium, gross alpha and gross beta concentrations in water were below the investigation levels for drinking water in the European Directive 2013/51.

### Liquid waste discharges and aquatic monitoring


Liquid wastes are disposed via a pipeline to deep water in Weymouth Bay. Tritium discharges increased from Winfrith (inner pipeline) in comparison to the (lowest reported) value in 2014. Figure 3.7 shows trends of liquid discharges over time (2000 – 2015) for tritium and alpha emitting radionuclides. Over the period, alpha-emitting radionuclide discharges have generally decreased since the peak in 2003 (although discharges peaked again in 2013). In comparison, tritium discharges have varied more between years, with periodic peaks in releases. Alpha-emitting radionuclides from Winfrith (inner pipeline) were the lowest reported value in 2015.

Analyses of seafood and marine indicator materials and measurements of external radiation over muddy intertidal areas were conducted. Data for 2015 are given in Tables 3.4(a) and (b). Concentrations of radionuclides in the marine environment largely continued at the low levels found in recent years. Gamma dose rates were difficult to distinguish from natural background.

## 3.4 Minor sites


Two minor sites are monitored using a small sampling programme of environmental materials. The results, given in the following sections, show that there was no detected impact on the environment in 2015 due to operation of these sites.

### 3.4.1 Culham, Oxfordshire



Culham Centre for Fusion Energy (CCFE), based at the Culham Science Centre, is the UK's national laboratory for fusion research. CCFE hosts an experimental fusion reactor, the Joint European Torus (JET), owned and operated by UKAEA

(under contract from Euratom). Although not currently designated, NDA understands that the intention of Government is to designate that part of the Culham Site occupied by JET facilities as an NDA site at an appropriate time after JET operation ceases. NDA would then take responsibility for the decommissioning programme that is expected to take 10 years to complete. The length of future operations is uncertain, but the assumption is that



**Figure 3.7.** Trends in liquid discharges of tritium and alpha emitting radionuclides from Winfrith, Dorset 2000-2015

operations will continue until 2018 and the facility is then decommissioned (DECC and NDA, 2014).

*Total dose* is not determined at this site, in this report, because an integrated habits survey has not been undertaken. The source specific dose, from using the River Thames directly as drinking water downstream of the discharge point at Culham in 2015, was estimated to be much less than 0.005 mSv (Table 3.1).

Monitoring of soil and grass around Culham and of sediment and water from the River Thames was undertaken in 2015. Locations and data are shown in Figure 3.4 and Table 3.5, respectively. In previous years, the main effect of the site's operation was the increased tritium concentrations found in grass collected near the site perimeter. In 2015, measurements of tritium in all samples are reported as less than values. Unlike in recent years, strontium-90 was detected at a very low concentration (just above the LoD) in the soil sample. Overall, no effects due to site operation were detected. The measured concentrations of caesium-137 in the River Thames sediment are not attributable to Culham but were due to past discharges from Harwell, nuclear weapons testing fallout from the 1950's and 1960's and the Chernobyl reactor accident in 1986.

### 3.4.2 Imperial College Reactor Centre, Ascot, Berkshire

The licensed reactor at Imperial College is a minor site with very low radioactive discharges, and is monitored using a small sampling programme for environmental materials.

The Reactor Centre provided facilities for the University and other organisations for research and commercial purposes. Imperial College undertook a review of the future of the Reactor Centre at Silwood Park which concluded that the reactor should be closed and decommissioned for financial reasons. The reactor was de-fuelled in the summer of 2014. The aim is that the reactor will be dismantled over a period of ten years with eventual de-licensing of the site by 2023 (DECC and NDA, 2014). In January 2015, Imperial College applied to the ONR for consent to carry out the decommissioning of the reactor. ONR undertook a public consultation on the submitted environmental statement and a decision to grant consent was issued in July 2015 (ONR, 2015).

In 2015, gaseous and aqueous discharges were very low (Appendix 2). Monitoring of the environmental effects involved the analysis of grass and crop (potato) samples by gamma-ray spectrometry. Activity concentrations in both samples, from radioactive discharges, are reported as less than values.

**Table 3.1. Individual doses – research sites, 2015**

Site	Representative person <sup>a</sup>	Exposure, mSv per year						
		Total	Fish and Shellfish	Other local food	External radiation from intertidal areas, river banks or fishing gear	Intakes of sediment and water	Gaseous plume related pathways	Direct radiation from site
Culham								
Source specific dose	Drinkers of river water	<0.005	-	-	-	<0.005	-	-
<b>Total dose – all sources</b>	<b>Adult occupants over sediment</b>	<b>0.010</b>	<b>&lt;0.005</b>			<b>0.010</b>	<b>-</b>	
Source specific doses	Seafood consumers	0.011	<0.005	-	0.010	-	-	-
	Geo occupants <sup>b</sup>	<0.005	-	-	<0.005	-	-	-
	Inhabitants and consumers of locally grown food	0.014	-	0.014	-	-	<0.005	-
Harwell								
<b>Total dose – all sources</b>	<b>Local adult inhabitants (0 - 0.25km)</b>	<b>0.017<sup>c</sup></b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>&lt;0.005</b>	<b>0.017</b>
Source specific doses	Anglers	<0.005	<0.005	-	<0.005	-	-	-
	Infant inhabitants and consumers of locally grown food	<0.005 <sup>c</sup>	-	<0.005	-	-	<0.005	-
Winfrith								
<b>Total dose – all sources</b>	<b>Prenatal children of local inhabitants (0.5 - 1km)</b>	<b>0.014</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>-</b>	<b>&lt;0.005</b>	<b>0.014</b>
Source specific doses	Seafood consumers	<0.005	<0.005	-	<0.005	-	-	-
	Infant inhabitants and consumers of locally grown food	<0.005	-	<0.005	-	-	<0.005	-

- <sup>a</sup> The total dose is the dose which accounts for all sources including gaseous and liquid discharges and direct radiation. The total dose for the representative person with the highest dose presented. Other dose values are presented for specific sources, either liquid discharges or gaseous discharges, and their associated pathways. They serve as a check on the validity of the total dose assessment. The representative person is an adult unless otherwise stated
- <sup>b</sup> People who visit Oigin's Geo, a coastal feature to the east of Dounreay
- <sup>c</sup> Includes a component due to natural sources of radionuclides

**Table 3.2(a). Concentrations of radionuclides in food and the environment near Dounreay, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			<sup>3</sup> H	<sup>60</sup> Co	<sup>65</sup> Zn	<sup>90</sup> Sr	<sup>95</sup> Nb	<sup>99</sup> Tc	<sup>137</sup> Cs
<b>Marine samples</b>									
Cod	Scrabster	2		<0.10	<0.20			<0.40	0.28
Crabs	Pipeline	2		<0.10	<0.19	0.10		<0.27	0.43
Crabs	Strathy	2		<0.10	<0.21			<0.58	
Crabs	Melvich Bay	2		<0.10	<0.23			<0.64	0.18
Winkles	Brims Ness	4		<0.10	<0.26	<0.10		<0.77	
Winkles	Sandside Bay	4		<0.12	<0.30	<0.10		<8.7	0.65
Mussels	Echnaloch Bay	4		<0.11	<0.29			<0.87	1.8
<i>Fucus vesiculosus</i>	Brims Ness	4		<0.10	<0.18			<0.52	
<i>Fucus vesiculosus</i>	Sandside Bay	4		<0.10	<0.22			<0.83	10
<i>Fucus vesiculosus</i>	Burwick Pier	4		<0.10	<0.21			<0.47	4.3
Sediment	Oigin's Geo	3		<0.10	<0.46			<4.8	
Sediment	Brims Ness	1		<0.10	<0.23			<0.34	
Sediment	Sandside Bay	1		<0.10	<0.24			<0.33	
Sediment	Melvich Bay	1		<0.10	<0.12			<0.21	
Sediment	Strathy	1		<0.10	<0.20			<0.39	
Sediment	Rennibister	1		<0.10	<0.33			<0.71	
Seawater	Brims Ness	2	<1.0	<0.10	<0.10			<0.10	
Seawater	Sandside Bay	2	<1.0	<0.10	<0.10			<0.10	
Spume	Oigin's Geo	1		<0.10	<0.19				0.53

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			<sup>154</sup> Eu	<sup>155</sup> Eu	<sup>238</sup> Pu	<sup>239</sup> Pu+ <sup>240</sup> Pu	<sup>241</sup> Am	Gross alpha	Gross beta
<b>Marine samples</b>									
Cod	Scrabster	2	<0.10	<0.15	0.0026	0.0082	0.012		
Crabs	Pipeline	2	<0.10	<0.16	<0.033	<0.033	<0.030	<1.3	140
Crabs	Strathy	2	<0.10	<0.14	<0.00080	0.0098	0.0084		
Crabs	Melvich Bay	2	<0.11	<0.19	<0.00070	<0.00080	0.0012		
Winkles	Brims Ness	4	<0.12	<0.19	<0.062	0.23	0.17		
Winkles	Sandside Bay	4	<0.12	<0.21	0.0087	0.046	0.049		
Mussels	Echnaloch Bay	4	<0.13	<0.22	0.010	0.056	0.048		
<i>Fucus vesiculosus</i>	Brims Ness	4	<0.10	<0.12			<0.10	2.6	360
<i>Fucus vesiculosus</i>	Sandside Bay	4	<0.10	<0.15			<0.10	<3.0	440
<i>Fucus vesiculosus</i>	Burwick Pier	4	<0.10	<0.13			<0.10		
Sediment	Oigin's Geo	3	<0.18	<0.36	0.65	3.0	6.0		
Sediment	Brims Ness	1	0.21	<0.25	1.5	8.7	11		
Sediment	Sandside Bay	1	<0.12	<0.24	2.3	11	7.1		
Sediment	Melvich Bay	1	<0.10	<0.14	0.39	1.6	1.7		
Sediment	Strathy	1	<0.12	<0.24	0.20	2.0	2.4		
Sediment	Rennibister	1	<0.21	0.92	0.51	2.3	<0.0013		
Seawater	Brims Ness	2	<0.10	<0.10			<0.10		
Seawater	Sandside Bay	2	<0.10	<0.10			<0.10		
Spume	Oigin's Geo	1	<0.10	<0.18	0.14	0.69	1.0		

**Table 3.2(a). continued**

Material	Location or selection <sup>b</sup>	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			<sup>3</sup> H	<sup>90</sup> Sr	<sup>95</sup> Nb	<sup>129</sup> I	<sup>137</sup> Cs	<sup>155</sup> Eu	<sup>234</sup> U
<b>Terrestrial samples</b>									
Beef muscle		1	<5.0	<0.10	<0.29	<0.25	0.06		<0.050
Beef offal		1	<5.0	<0.10	<0.67	<0.13	<0.05		<0.050
Broccoli		1	<5.0	0.17	<0.15	<0.05	0.50		
Cabbage		1	<5.0	0.17	<0.12	<0.06	0.09		
Eggs		1	<5.0	<0.10	<0.30	<0.10	<0.05		
Honey		1	<5.0	0.52	<0.19	<1.7	7.0		
Lamb muscle		1	<5.0	<0.10	<0.33	<0.25	0.31		<0.050
Mint		1	<5.0	0.87	<0.45	<0.08	0.16		
Potatoes		1	<5.0	<0.10	<0.25	<0.05	0.10		
Rabbit		1	<5.0	<0.10	<0.61	<0.38	<0.06		
Rosehips		1	<5.0	0.58	<0.34	<0.05	0.15		
Turnips		1	<5.0	<0.10	<0.12	<0.06	<0.05		
Wild mushrooms		2	<5.0	<0.13	<0.32	<0.11	1.6		
Wild mushrooms	max			0.15	<0.49		3.1		
Grass		6	<5.0	0.21	<0.46	<0.05	<0.08		<0.050
Grass	max			0.34	<0.68		0.15		
Soil		6	<5.0	0.99	<0.35	<0.55	19	1.8	28
Soil	max			1.4	<0.89	<0.64	36	2.1	54
Freshwater	Loch Calder	1	<1.0		<0.02		<0.01		
Freshwater	Loch Shurrery	1	<1.0		<0.01		<0.01		
Freshwater	Loch Baligill	1	<1.0		<0.02		<0.01		
Freshwater	Heldale Water	1	<1.0		<0.03		<0.01		

Material	Location or selection <sup>b</sup>	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			<sup>235</sup> U	<sup>238</sup> U	<sup>238</sup> Pu	<sup>239</sup> Pu+ <sup>240</sup> Pu	<sup>241</sup> Am	Gross alpha	Gross beta
<b>Terrestrial samples</b>									
Beef muscle		1	<0.050	<0.050	<0.050	<0.050	<0.050		
Beef offal		1	<0.050	<0.050	<0.050	<0.050	<0.050		
Broccoli		1			<0.050	<0.050	<0.050		
Cabbage		1			<0.050	<0.050	<0.050		
Eggs		1			<0.050	0.012	<0.050		
Honey		1			<0.050	<0.050	<0.050		
Lamb muscle		1	<0.050	<0.050	<0.050	<0.050	<0.050		
Mint		1			<0.050	<0.050	<0.050		
Potatoes		1			<0.050	<0.050	<0.050		
Rabbit		1			<0.050	<0.050	<0.050		
Rosehips		1			<0.050	<0.050	<0.050		
Turnips		1			<0.050	<0.050	<0.050		
Wild mushrooms		2			<0.050	<0.076	<0.050		
Wild mushrooms	max					0.10			
Grass		6	<0.050	<0.050	<0.050	<0.050	<0.053		
Grass	max						0.068		
Soil		6	1.2	27	<0.050	0.37	0.22		
Soil	max		2.7	50		0.62	0.34		
Freshwater	Loch Calder	1					<0.01	<0.010	0.060
Freshwater	Loch Shurrery	1					<0.01	0.015	0.070
Freshwater	Loch Baligill	1					<0.01	<0.010	0.050
Freshwater	Heldale Water	1					<0.01	<0.010	0.060

<sup>a</sup> Except for seawater and freshwater where units are Bq l<sup>-1</sup>, and for soil and sediment where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'Max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

**Table 3.2(b). Monitoring of radiation dose rates near Dounreay, 2015**

Location	Material or ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Sandside Bay	Sand	2	0.066
Sandside Bay	Winkle bed	2	0.14
Oigin's Geo	Rocks	2	0.12
Brims Ness	Sediment	2	0.090
Melvich	Salt marsh	2	0.061
Melvich	Sediment	2	<0.051
Strathy	Sand	2	<0.048
Thurso riverbank	Sediment	2	0.079
Achvarasdal	Grass	2	0.077
Thurso Park	Grass	2	0.071
Borrowston Mains	Grass	2	0.079
East of Dounreay	Grass	2	0.081
Castletown Harbour	Sand	2	0.078
Dunnet Bay	Sand	2	0.054
<b>Mean beta dose rates</b>			$\mu\text{Sv h}^{-1}$
Sandside Bay	Sediment	2	<1.0
Oigin's Geo	Sediment	2	<1.0
Thurso riverbank	Sediment	2	<1.0
Castletown Harbour	Lobster pots	2	<1.0

**Table 3.2(c). Radioactivity in air near Dounreay, 2015**

Location	No. of sampling observations	Mean radioactivity concentration, $\text{mBq m}^{-3}$		
		$^{137}\text{Cs}$	Gross alpha	Gross beta
Shebster	12	<0.011	<0.011	<0.20
Reay	12	<0.0099	<0.012	<0.20
Balmore	12	<0.0099	0.015	<0.21

**Table 3.3(a). Concentrations of radionuclides in food and the environment near Harwell, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			<sup>3</sup> H	<sup>60</sup> Co	<sup>131</sup> I	<sup>137</sup> Cs	<sup>238</sup> Pu
<b>Freshwater samples</b>							
Flounder	Woolwich Reach	1	<25	<0.06	<0.26	<0.06	
Sediment	Appleford	4 <sup>E</sup>		<0.33	16	7.2	<0.33
Sediment	Outfall (Sutton Courtenay)	4 <sup>E</sup>		<0.57	2.1	<7.6	<0.18
Sediment	Day's Lock	4 <sup>E</sup>		<0.37	3.4	8.1	<0.19
Sediment	Lydebank Brook	4 <sup>E</sup>		<1.3		4.9	<0.22
Freshwater	Day's Lock	4 <sup>E</sup>	<3.0	<0.31		<0.25	
Freshwater	Lydebank Brook	4 <sup>E</sup>	<3.2	<0.25		<0.21	
Freshwater	R Thames (above discharge point)	4 <sup>E</sup>	<3.1	<0.22		<0.19	
Freshwater	R Thames (below discharge point)	4 <sup>E</sup>	<3.3	<0.23		<0.19	

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>			
			<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am	Gross alpha	Gross beta
<b>Freshwater samples</b>						
Flounder	Woolwich Reach	1		<0.16		
Sediment	Appleford	4 <sup>E</sup>	0.44	<1.5	<120	290
Sediment	Outfall (Sutton Courtenay)	4 <sup>E</sup>	0.17	<0.67	<130	290
Sediment	Day's Lock	4 <sup>E</sup>	0.32	<1.1	<130	250
Sediment	Lydebank Brook	4 <sup>E</sup>	0.47	<1.1	<130	360
Freshwater	Day's Lock	4 <sup>E</sup>			<0.042	0.24
Freshwater	Lydebank Brook	4 <sup>E</sup>			<0.058	0.18
Freshwater	R Thames (above discharge point)	4 <sup>E</sup>			<0.059	0.28
Freshwater	R Thames (below discharge point)	4 <sup>E</sup>			<0.044	0.23

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>	
			Organic <sup>3</sup> H	<sup>3</sup> H
<b>Terrestrial samples</b>				
Milk		2	<3.2	<3.2
Milk	max		<3.8	<3.8
Strawberries		1	<4.1	<4.1
Grass		1	<3.2	<3.2

<sup>a</sup> Except for milk where units are Bq l<sup>-1</sup>, and for sediment where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima.

If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 3.3(b). Monitoring of radiation dose rates near Harwell, 2015**

Location	Ground type	No. of sampling observations	μGy h <sup>-1</sup>
<b>Mean gamma dose rates at 1m over substrate</b>			
Appleford	Grass and mud	2	0.072
Sutton Courtenay	Mud	2	0.077
Day's Lock	Mud	1	0.068
Day's Lock	Grass	1	0.064

**Table 3.4(a). Concentrations of radionuclides in food and the environment near Winfrith, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			<sup>14</sup> C	<sup>60</sup> Co	<sup>99</sup> Tc	<sup>137</sup> Cs	<sup>238</sup> Pu
<b>Marine samples</b>							
Plaice	Weymouth Bay	1		<0.05		0.09	
Crabs	Lulworth Banks	1	31	<0.06		<0.05	
Scallops	Lulworth Ledges	1		<0.14		<0.11	0.00036
Seaweed	Lulworth Cove	1 <sup>E</sup>		<0.61	<0.54	<0.40	
Seaweed	Bognor Rock	2 <sup>E</sup>		<0.53	0.31	<0.40	
Seawater	Lulworth Cove	1 <sup>E</sup>		<0.24		<0.20	
Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm + <sup>244</sup> Cm	Gross alpha
<b>Marine samples</b>							
Plaice	Weymouth Bay	1		<0.05			
Crabs	Lulworth Banks	1		<0.05			
Scallops	Lulworth Ledges	1	0.0031	0.00065	*	0.000011	
Seaweed	Lulworth Cove	1 <sup>E</sup>		<0.58			
Seaweed	Bognor Rock	2 <sup>E</sup>		<0.49			
Seawater	Lulworth Cove	1 <sup>E</sup>		<0.30		<2.5	14
Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			Organic <sup>3</sup> H	<sup>3</sup> H	<sup>14</sup> C	<sup>137</sup> Cs	Gross alpha
<b>Terrestrial samples</b>							
Milk		2	<3.4	<3.4	18	<0.07	
Milk	max		<3.7	<3.7			
Carrot		1	<3.1	<3.1	13	<0.04	
Grass		1	<4.2	<4.2	19	<0.06	
Sediment	North of site (Stream A)	1 <sup>E</sup>				0.95	<110
Sediment	R Frome (upstream)	1 <sup>E</sup>				1.3	<94
Sediment	R Frome (downstream)	1 <sup>E</sup>				2.4	<97
Sediment	R Win, East of site	1 <sup>E</sup>				0.54	<97
Freshwater	North of site (Stream A)	2 <sup>E</sup>		9.7		<0.25	<0.059
Freshwater	R Frome (upstream)	2 <sup>E</sup>		<3.0		<0.21	<0.034
Freshwater	R Frome (downstream)	2 <sup>E</sup>		<3.1		<0.23	<0.035
Freshwater	R Win, East of site	2 <sup>E</sup>		<3.1		<0.25	<0.064

\* Not detected by the method used

<sup>a</sup> Except for milk and freshwater where units are Bq l<sup>-1</sup>, and for sediment where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 3.4(b). Monitoring of radiation dose rates near Winfrith, 2015**

Location	Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Weymouth Bay	Sand and stones	1	0.056
Red Cliffe Point to Black Head	Shingle	1	0.051
Osmington Mills	Pebbles	1	0.057
Ringstead Bay	Sand and shingle	1	0.053
Durdle Door	Shingle	1	0.048
St Oswald's Head	Pebbles and sand	1	0.053
Lulworth Cove	Sand	1	0.056
Kimmeridge Bay	Pebbles	1	0.077
Swanage Bay 1	Sand	1	0.060
Swanage Bay 2	Sand	1	0.057
Swanage Bay 3	Sand	1	0.058
Poole Harbour	Sand and stones	1	0.052

**Table 3.5. Concentrations of radionuclides in the environment near Culham, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>							
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>90</sup> Sr	<sup>137</sup> Cs	Gross alpha	Gross beta	
Freshwater	River Thames (upstream)	2	<3.1					<0.19	<0.054	0.32
Freshwater	River Thames (downstream)	2	<3.2					<0.21	<0.059	0.39
Grass	1 km East of site perimeter	1	<9.0	<3.5	<1.7	<0.46		<0.83		210
Sediment	River Thames (upstream)	2						12		
Sediment	River Thames (downstream)	2						36		
Soil	1 km East of site perimeter	1	<6.1	<8.4	<5.3	4.0		3.5		270

<sup>a</sup> Except for freshwater where units are Bq l<sup>-1</sup>, and for sediment and soil where dry concentrations apply

## 4. Nuclear power stations

### Key points

- *Total doses* for the representative person were less than 6 per cent of the dose limit for all sites assessed
- Electricity production continued at one Magnox station (Wylfa), one PWR station (Sizewell B) and seven AGR stations in 2015
- Discharges, environmental concentrations and dose rates in 2015 were broadly similar to those in 2014
- Concentrations of radiocaesium and transuranic elements were enhanced around some sites. These were mainly due to discharges from Sellafield

### Berkeley, Gloucestershire and Oldbury, South Gloucestershire

- *Total dose* for the representative person was low and unchanged in 2015
- Gaseous discharges of tritium and carbon-14 from Berkeley, and liquid discharges of tritium and caesium-137 from Oldbury, increased in 2015

### Bradwell, Essex

- *Total dose* for the representative person increased in 2015
- New source specific assessment undertaken to determine dose to houseboat dwellers
- Enhanced monitoring undertaken in 2015

### Chapelcross, Dumfries and Galloway

- *Total dose* for the representative person increased in 2015
- Gaseous discharges of "all other nuclides" increased in 2015

### Dungeness, Kent

- *Total dose* for the representative person decreased in 2015
- Gaseous discharges of tritium, sulphur-35, and argon-41, and liquid discharges of tritium and sulphur-35, increased from Dungeness B in 2015

### Hartlepool, County Durham

- *Total dose* for the representative person decreased in 2015
- Liquid discharges of tritium and sulphur-35, decreased in 2015

### Heysham, Lancashire

- *Total dose* for the representative person was unchanged in 2015
- Gaseous discharges of carbon-14 and argon-41 increased at Heysham 2, and liquid discharges of tritium decreased from Heysham 1 and Heysham 2, in 2015

### Hinkley Point, Somerset

- *Total dose* for the representative person decreased in 2015
- Gaseous discharges of carbon-14 and argon-41, and liquid discharges of tritium and sulphur-35, decreased from Hinkley B in 2015

### Hunterston, North Ayrshire

- Gaseous discharges of carbon-14 increased from Hunterston B in 2015

### Sizewell, Suffolk

- *Total dose* for the representative person was generally unchanged in 2015
- Liquid discharges of caesium-137 and tritium decreased from Sizewell A and Sizewell B, respectively, in 2015

### Torness, East Lothian

- *Total dose* for the representative person was unchanged in 2015

### Trawsfynydd, Gwynedd

- *Total dose* for the representative person was similar in 2015
- Gaseous discharges of tritium, and liquid discharges of tritium and "other radionuclides", decreased in 2015
- Concentration of caesium-137 in brown trout was the lowest reported value in 2015

### Wylfa, Isle of Anglesey

- *Total dose* for the representative person increased in 2015
- Power generation from the Magnox station ceased in December 2015
- Gaseous and liquid discharges of tritium decreased, and gaseous discharges of carbon-14, sulphur-35 and argon-41 increased, in 2015

This section considers the results of environment and food monitoring by the Environment Agency, FSA and SEPA from nuclear power stations. There is a total of 19 nuclear power stations at 14 locations, nine in England (Berkeley, Oldbury, Bradwell, Calder Hall, Dungeness, Hartlepool, Heysham, Hinkley Point and Sizewell), three in Scotland (Chapelcross, Hunterston and Torness) and two in Wales (Trawsfynydd and Wylfa). Some of these stations are being decommissioned.

Eleven of the 19 nuclear power stations are older Magnox power stations, owned by NDA. NDA (set up under the Energy Act 2004) is a non-departmental public body, with a remit to secure the decommissioning and clean-up of the UK's civil public sector nuclear licensed sites. Only one Magnox station (Wylfa) continued to generate electricity in 2015, others are in the process of de-fuelling or decommissioning. In March 2016, NDA published a business plan which summarises the programme of work at each of the sites during 2016/19 (NDA, 2016b).

In March 2014, NDA announced that the Cavendish Fluor Partnership, a joint venture between Cavendish Nuclear (a wholly-owned subsidiary of Babcock International Group plc) and Fluor Corporation, had been selected to take ownership of Magnox Limited. In 2013, Magnox Limited managed ten nuclear sites and was owned and operated by Energy Solutions on behalf of NDA. In September 2014, NDA formally appointed Cavendish Fluor Partnership (CFP) as the new PBO for and Magnox Limited (and RSRL). In November 2014, ONR received an application to relicence the ten Magnox sites into a single site licence company alongside the Harwell and Winfrith sites. As of 1st April 2015, Harwell and Winfrith sites, previously operated by RSRL, merged to be part of Magnox Limited. Concurrently the EPR 2010 radioactive substances permits for each site were also transferred to Magnox Limited.

Calder Hall is being decommissioned; it is operated by Sellafield Limited and discharges from this Magnox power station are considered in Section 2 because it is located at Sellafield.

Seven AGR power stations and one Pressurised Water Reactor (PWR) power station were owned and operated by EDF Energy Nuclear Generation Limited in 2015; these are Dungeness B, Hartlepool, Heysham 1 and 2, Hinkley Point B and Sizewell B Power Stations in England, and Hunterston B and Torness Power Stations in Scotland. All of these were generating electricity during 2015.

Gaseous and liquid discharges from each of the power stations are regulated by the Environment Agency and NRW in England and Wales, respectively and by SEPA in Scotland. In 2015, gaseous and liquid discharges were below regulated limits for each of the power stations (see Appendix 2). Independent monitoring of the environment around each of the power stations is conducted by the FSA and the Environment Agency in England and Wales, and by


SEPA in Scotland. In Wales, this is conducted on behalf of NRW and the Welsh Government.

The medium-term trends in dose, discharges and environmental concentrations at these sites have been considered in a summary report (Environment Agency, FSA, NIEA and SEPA, 2010b).

The sites are grouped in this Section according to whether they are in England, Scotland or Wales.

## ENGLAND

### 4.1 Berkeley, Gloucestershire and Oldbury, South Gloucestershire



Berkeley and Oldbury are both Magnox power stations. Berkeley Power Station is situated on the eastern bank of the River Sever and was powered by two Magnox reactors. Berkeley was the first commercial power

station in the UK to enter into decommissioning. It ceased electricity generation in 1989 and de-fuelling was completed in 1992. Decommissioning is still in progress and radioactive wastes are still generated by these operations. The Berkeley site will enter the Care and Maintenance phase by the year 2021. Thereafter, the current plan is to de-license the Berkeley site (released from regulatory control). Final site clearance is expected to commence in 2070 and be achieved by 2079 (NDA, 2016a).

Oldbury Power Station is located on the south bank of the River Sever close to the village of Oldbury-on-Severn and has two Magnox reactors. Electricity generation started in 1967 and ceased in 2012. De-fuelling was completed in 2016 and the site is now focusing on the retrieval, processing, storage and dispatch of waste. The Oldbury site will enter the Care and Maintenance phase by the year 2027. Thereafter, the current plan is to de-license the Oldbury site (released from regulatory control). Final site clearance is expected to commence in 2092 and be achieved by 2101 (NDA, 2016a).

Berkeley and Oldbury sites are considered together for the purposes of environmental monitoring because the effects from both sites contribute to the same area. The most recent habits survey was undertaken in 2014 (Clyne *et al.*, 2015).

## Doses to the public

The *total dose* from all pathways and sources of radiation is assessed to have been less than 0.005 mSv in 2015 (Table 4.1), which was less than 0.5 per cent of the dose limit, and unchanged from 2014. In 2015, the representative person was a 1 year-old infant consuming milk at high-rates. The trend in the *total dose* over the period 2004 – 2015 is given in Figure 4.1. Any longer-term variations in *total doses* with time are attributable to changes in the contribution from direct radiation from the site.

The source specific assessment for a high-rate consumer of fish and shellfish gives an exposure that was also less than 0.005 mSv in 2015 (Table 4.1). The dose to a consumer of fish and shellfish includes external radiation, a component due to the tritium historically originating from GE Healthcare Limited at Cardiff, and a component of the dose resulting from an increased tritium dose coefficient (see Appendix 1). The dose to a high-rate consumer of locally grown foods was estimated to be 0.005 mSv. The small increase in dose (from < 0.005 mSv in 2014) was due to higher carbon-14 concentrations in milk in 2015. As in 2014, an assessment was undertaken to determine the external exposure for houseboat dwellers in the vicinity of the Berkeley and Oldbury sites (as identified in the recent habits survey). The estimated dose was 0.006 mSv for this activity (Table 4.1), and down from 0.022 mSv in 2014. The lower value in 2015 was due to a decrease in the dose rates at Sharpness, in comparison to those in 2014. The estimate for this pathway is determined as a cautious value (and therefore not included in the *total dose* assessment), because gamma dose rate measurements used were not necessarily representative of the types of ground type and houseboat location (as identified in the habits survey).


## Gaseous discharges and terrestrial monitoring

The Berkeley and Oldbury sites discharge gaseous radioactive wastes via separate stacks to the atmosphere. Discharges of tritium, and carbon-14 to a lesser extent, increased from Berkeley, in comparison to releases in 2014. The focus of the terrestrial sampling was for the content of tritium, carbon-14 and sulphur-35 in milk and crops. Local freshwater samples were also analysed. Data for 2015 are given in Table 4.2(a). Sulphur-35 was detected at very low levels in one terrestrial food sample (wheat). Carbon-14 was detected in locally produced foods at concentrations close to background values. Carbon-14 concentrations in foodstuffs (milk) increased by small amounts in comparison to those in 2014. Tritium, gross alpha and gross beta concentrations in surface water were below the investigation levels for drinking water in the European Directive 2013/51.

## Liquid waste discharges and aquatic monitoring

Liquid radioactive wastes are discharged to the Severn Estuary. Discharges of tritium, and to a lesser extent caesium-137, increased from Oldbury, in comparison to releases in 2014. Analyses of seafood and marine indicator materials and measurements of external radiation were conducted over muddy intertidal areas. Measurements of tritium in seafood were made in order to monitor the additional local effects of historical discharges from the GE Healthcare Limited radiopharmaceutical plant in Cardiff (see Section 6). Data for 2015 are given in Tables 4.2(a) and (b). Most of the artificial radioactivity detected was due to caesium-137, representing the combined effect of discharges from the sites, other nuclear establishments discharging into the Bristol Channel and weapons testing, and possibly a small Sellafield-derived component. There is now limited evidence to suggest that caesium-137 concentrations in sediment have been generally decreasing over the last decade (Figure 4.2). As in recent years, the tritium concentration in fish is reported as a less than value. In previous years, the levels of tritium in seafood have been relatively high and were likely to be mainly due to historical discharges from GE Healthcare Limited, Cardiff. Very small concentrations of other radionuclides were detected but, taken together, were of low radiological significance. Gamma dose rates were generally similar, although rates at Oldbury (south and south west) and Sharpness decreased by small amounts, in comparison to those in 2014.


### 4.2 Bradwell, Essex



The Bradwell site is located on the south side of the Blackwater Estuary. This Magnox power station ceased electricity production in 2002 after 40 years of operation, and de-fuelling was completed in 2006. The focus for the

site is the completion of decommissioning projects. Bradwell will be the first Magnox site in England to enter the Care and Maintenance phase, which is currently planned for 2019. The Care and Maintenance phase is expected to last for around 70 years. Thereafter, the plan is for final site clearance to commence in 2083 and be achieved by 2092 (NDA, 2016a).

During 2014, the Environment Agency carried out a review of their own environmental monitoring programme, which concluded that the Bradwell programme was consistent with the published technical guidance (Environment Agency, FSA and SEPA, 2010). However, in



**Figure 4.1.** Total dose at nuclear power stations, 2004-2015  
 (Small doses less than or equal to 0.005 mSv are recorded as being 0.005 mSv)




Figure 4.2. Caesium-137 concentration in marine sediments near nuclear power stations between 2004-2015

response to requests from local stakeholders to enhance the environmental monitoring, whilst treatment of Intermediate Level Waste (Fuel Element Debris) was being carried out, the Environment Agency expanded the size of their environmental monitoring programme. These enhancements to the environmental monitoring started at the beginning of 2015 and continued throughout the year. These enhancements are only planned to remain in place whilst this waste treatment is being carried out; thereafter it will then revert to the baseline requirements.

In August 2015, a habits survey was conducted to determine the consumption and occupancy rates by members of the public (Clyne et al., 2016b). Small decreases in both the fish and crustacean consumption rates and the occupancy rate have been observed, together with an increase in the mollusc consumption rate, in comparison with those of the previous survey in 2007. Revised figures for consumption rates, together with occupancy rates, are provided in Appendix 1 (Table X2.2).

## Doses to the public

The *total dose* from all pathways and sources of radiation was 0.017 mSv in 2015 (Table 4.1), which was less than 2 per cent of the dose limit for members of the public of 1 mSv, and up from less than 0.005 mSv (in 2014). The representative person was a prenatal child of local inhabitants in 2015, and a change from that in 2014 (adult consumer of fish). The increase in *total dose* was due to a higher estimate of direct radiation from the site in 2015. The trend in *total dose* over the period 2004 – 2015 is given in Figure 4.1. Any significant variations in *total dose* with time were attributed to changes in the estimate of direct radiation.

The source specific assessments for a high-rate consumer of locally grown foods, and of fish and shellfish give exposures that were less than the *total dose* in 2015 (Table 4.1). An additional source specific assessment was undertaken in 2015 to determine the external exposure based on the 2015 enhanced monitoring. The estimated dose was also less than 0.005 mSv. This estimate is determined as a cautious value (using maximising assumptions for gamma dose rates and occupancy rates) and therefore not included in the *total dose* assessment.

## Gaseous discharges and terrestrial monitoring

This power station is permitted to discharge gaseous wastes to the local environment via stacks to the atmosphere. Terrestrial sampling is similar to that for other power stations including analyses of milk and crop samples for tritium, carbon-14 and sulphur-35. Samples of water are also taken from a coastal ditch and public supplies. In 2015, a number of grass samples were also collected and analysed, to enhance the environmental

monitoring programme. Data for 2015 are given in Table 4.3(a). Activity concentrations were low in terrestrial samples, although some small enhancements of carbon-14 concentrations were measured in food samples. In grass samples, tritium, carbon-14 and caesium-137 concentrations were all reported as less than values. Tritium, gross alpha and gross beta concentrations in freshwater (public supplies) were below the investigation levels for drinking water in the European Directive 2013/51. As in previous years, the gross beta activities in water from the coastal ditch continued to be enhanced above background levels, and these were in excess of the WHO screening level for drinking water (1 Bq l<sup>-1</sup>). Tritium concentrations in coastal ditches were similar to those in recent years, with positively detected values substantially below the EU reference level for tritium of 100 Bq l<sup>-1</sup>. The water in the ditches is not known to be used as a source of drinking water.

## Liquid waste discharges and aquatic monitoring

Liquid wastes are discharged into the River Blackwater estuary. Aquatic sampling was directed at consumption of locally caught fish and shellfish and external exposure over intertidal sediments. Seaweeds were also analysed as an environmental indicator material. In 2015, a number of sediment and seawater samples were also collected and analysed, and measurement of gamma dose rates, to enhance the environmental monitoring programme. Data for 2015 are given in Tables 4.3(a) and (b). Low concentrations of artificial radionuclides were detected in aquatic materials as a result of discharges from the station, discharges from Sellafield and weapons testing. In seawater samples, tritium, caesium-137 and americium-241 concentrations were all reported as less than values. Due to the low concentrations detected, it is generally difficult to attribute the results to a particular source; however, concentrations (including those for the 2015 enhanced sediment monitoring) were generally similar in magnitude to those results reported in recent years. There is an overall decline in caesium-137 concentrations in sediments over the last decade (Figure 4.2), and the activity concentration in 2015 was similar in comparison to the (lowest reported) value in 2014. The technetium-99 detected in seaweeds at Bradwell was likely to be due to the long distance transfer of Sellafield derived activity. Gamma dose rates on beaches, and those taken for the enhanced monitoring in 2015, were difficult to distinguish from natural background.

### 4.3 Dungeness, Kent



The Dungeness power stations are located on the south Kent coast between Folkestone and Rye. There are two separate A and B nuclear power stations on neighbouring sites; the A station was powered by two Magnox reactors

and the B station has two AGRs. Discharges are made via separate and adjacent outfalls and stacks, but for the purposes of environmental monitoring these are considered together. Dungeness A ceased generating electricity in 2006. De-fuelling of both Magnox reactors was completed in 2012. The Dungeness A site will enter the Care and Maintenance phase by the year 2027. Final site clearance is expected to commence in 2087 and be achieved by 2097 (NDA, 2016a). Dungeness B is expected to continue electricity generation until 2028 (previously scheduled for cessation in 2018). EDF Energy must continue to demonstrate that the station complies with the Safety Cases which are reviewed by ONR to enable continued operation to 2028. The most recent habits survey was undertaken in 2010 (Clyne *et al.*, 2011b).

#### Doses to the public

In 2015, the *total dose* from all pathways and sources of radiation was 0.014 mSv (Table 4.1), or approximately 1 per cent of the dose limit of 1 mSv, and down from 0.021 mSv in 2014. The decrease in *total dose* (from 2014) was mostly attributed to a lower estimate of direct radiation from the site in 2015. As in recent years, this is almost entirely due to direct radiation from the site. An adult living near to the site was the representative person. The trend in *total dose* over the period 2004 – 2015 is given in Figure 4.1. *Total doses* ranged between 0.015 and 0.63 mSv over the time period and were dominated by direct radiation. Following the shut-down of the Magnox reactors in 2006, this dose has significantly declined.

Source specific assessments for a high-rate consumer of locally grown foodstuffs, for a local bait digger (who consumes large quantities of fish and shellfish and spends long periods of time in the location being assessed), and for a houseboat occupant give exposures that were less than the *total dose* (Table 4.1). The doses to a high-rate consumer of locally grown foods and local bait digger were estimated to be 0.005 mSv and 0.006 mSv, respectively. The dose to a houseboat dweller from external exposure was less than 0.005 mSv and a small decrease from 0.005 mSv (in 2014) was because gamma dose rates were measured on different types of ground type (at Rye Bay)

in 2015. In 2014 the ground type comprised of sand and shingle, whereas in 2015 it comprised of two ground types (both mud and sand and sand and shingle).

#### Gaseous discharges and terrestrial monitoring

Discharges of tritium, sulphur-35 and argon-41 increased, by a small amount, from Dungeness B, in comparison to releases in 2014. The focus of the terrestrial sampling was analyses of tritium, carbon-14 and sulphur-35 in milk and crops. The results of monitoring for 2015 are given in Tables 4.4(a). Activity concentrations in many terrestrial foods are reported as less than values (or close to the less than value). As in previous years, low concentrations of sulphur-35 were positively detected in some locally produced foods and carbon-14 was detected in milk above background values. Tritium, gross alpha and gross beta concentrations in freshwater were below the investigation levels for drinking water in the European Directive 2013/51.

#### Liquid waste discharges and aquatic monitoring

Discharges of tritium, and sulphur-35 to a lesser extent, increased from Dungeness B, in comparison to releases in 2014. Marine monitoring included gamma dose rate measurements, and analysis of seafood and sediments. The results of monitoring for 2015 are given in Tables 4.4(a) and (b). Caesium-137 concentrations in seafood and marine materials are attributable to discharges from the stations and to weapon test fallout with a long distance contribution from Sellafield and La Hague. Due to the low concentrations detected, it is generally difficult to attribute the results to a particular source. The low concentrations of transuranic nuclides in scallops were typical of levels expected at sites remote from Sellafield. No tritium was detected in seafood in 2015. Caesium-137 concentrations in sediment have remained low over the last decade (Figure 4.2); the apparent increase in 2010 was due to the inclusion of a value ( $<5.8 \text{ Bq kg}^{-1}$ ) which was reported as a less than value. Gamma dose rates were generally difficult to distinguish from the natural background.

## 4.4 Hartlepool, County Durham



Hartlepool Power Station is situated on the mouth of the Tees estuary, on the north east coast of England. This station, which is powered by twin AGRs, began operation in 1983. It is estimated that its power generation will

continue until at least 2024 (extended from 2019). The most recent habits survey was undertaken in 2014 (Garrod *et al.*, 2015b).

### Doses to the public

The *total dose* from all pathways and sources of radiation was 0.022 mSv in 2015 (Table 4.1), which was approximately 2 per cent of the dose limit, and down from 0.027 mSv in 2014. The decrease in *total dose* (from 2014) was mostly attributed to a lower estimate of direct radiation from the site in 2015. The representative person was an adult spending time living near to the site whose dose was from direct radiation (from the site) and external exposure from activity in sand and sediment on local beaches. The trend in *total dose* over the period 2004 – 2015 is given in Figure 4.1. *Total doses* remained broadly similar from year to year, and were low.

Source specific assessments for both high-rate consumers of locally grown foodstuffs, and of fish and shellfish, give exposures that were less than the *total dose* (Table 4.1). The dose to a local fish and shellfish consumer, including external radiation but excluding naturally occurring radionuclides, was 0.020 mSv. The dose in 2014 was 0.012 mSv, and the reason for the increase in 2015 is the same as that contributing to the maximum *total dose*.

Since 2012, a source specific assessment has been undertaken to determine the exposure from naturally occurring radionuclides, as a consequence of the reported polonium-210 concentrations in mollusc samples. As in previous years, winkle samples were collected in 2015 for South Gare (inside the Tees Estuary entrance) consisted of a mixture including some winkles from the estuary entrance near Paddy's Hole. The area in the close proximity of Paddy's Hole was unlikely to sustain a high-rate consumption of winkles, as it is an extremely localised area which contains oil and other wastes. In addition, the most recent habits survey undertaken in 2008 did not identify any consumption of molluscs from Paddy's Hole. However, in the event that some of these molluscs were a constituent of the diet of a high-rate consumer of fish and shellfish, the dose from naturally occurring

radionuclides was considered. In 2015, the polonium-210 (and lead-210) concentration was not enhanced above background, and therefore no additional exposure contributed to the dose, in addition to that from artificial radionuclides. This estimate assumes that the median concentrations for naturally occurring radionuclides at background (Appendix 1, Table X4.1) be subtracted from the total concentrations as measured in 2015.

### Gaseous discharges and terrestrial monitoring

Gaseous radioactive waste is discharged via stacks to the local environment. Analyses of tritium, carbon-14, sulphur-35 and gamma emitters were made in milk and crop samples. Samples of water are also taken from a borehole and public water supplies. Data for 2015 are given in Table 4.5(a). The effects of gaseous disposals from the site were not easily detectable in foodstuffs, although small enhancements of sulphur-35 concentrations (reported as just above the less than value) were measured one food sample (barley). Carbon-14 was detected in locally produced milk at concentrations close to expected background values. Tritium, gross alpha and gross beta concentrations in freshwater were below the investigation levels for drinking water in the European Directive 2013/51.

### Liquid waste discharges and aquatic monitoring

Regulated discharges of radioactive liquid effluent are made to Hartlepool Bay with a minor component being discharged directly to the River Tees. Discharges of tritium and sulphur-35 decreased in 2015, in comparison to those in 2014. Results of the aquatic monitoring programme conducted in 2015 are shown in Tables 4.5(a) and (b). As in previous years, small enhancements of carbon-14 concentrations, above expected background, were observed in seafood samples. Enhancements are most likely to be due to carbon-14 discharges from a nearby non-nuclear site since carbon-14 discharges from the power station are low. Technetium-99 analysis in seaweed is used as a specific indication of the far-field effects of disposals to sea from Sellafield. Concentrations in seaweed were low and much less than the peak observed in 1998 (see also Figure 2.9). They are less than 1 per cent of the equivalent concentrations near Sellafield. Iodine-131 was again positively detected in seaweed samples collected around the mouth of the River Tees Estuary in 2015. The detected values, as in previous years, are believed to originate from the therapeutic use of this radionuclide in a local hospital. Detectable concentrations of radiocaesium and transuranics were mainly due to disposals from Sellafield and to weapon test fallout. However, caesium-137 concentrations in sediment have remained low for a number of years (Figure 4.2). In 2015, the reported polonium-210 and lead-210 concentrations

in winkles from South Gare (consisting of a mixture including some winkles collected from the estuary entrance near Paddy's Hole) were 13 Bq kg<sup>-1</sup> and 1.2 Bq kg<sup>-1</sup>, respectively and values expected due to natural sources. The polonium-210 concentration was lower in comparison to those concentrations in recent years (enhanced above background). Overall, gamma dose rates in 2015 were generally higher over sand (e.g. Seaton, Sands, Seaton Carew and Carr House) in comparison to those in 2014.

## 4.5 Heysham, Lancashire



Heysham Power Station is situated on the Lancashire coast to the south of Morecambe and near the port of Heysham. This establishment comprises two separate nuclear power stations, both powered by two AGRs. It is

estimated that Heysham 1 and 2 will continue to generate electricity until at least 2024 and 2030, respectively (extended from 2019 and 2023). Disposals of radioactive waste from both stations are made under permit via separate outfalls to Morecambe Bay and via stacks, but for the purposes of environmental monitoring both stations are considered together. The most recent habits survey was undertaken in 2011 (Garrod *et al.*, 2012).

### Doses to the public

The *total dose* from all pathways and sources of radiation was 0.023 mSv in 2015 (Table 4.1), or approximately 2 per cent of the dose limit for members of the public, and unchanged from 2014. The representative person was an adult who was a high-rate consumer of molluscs. The trend in *total dose* over the period 2004 – 2015 is given in Figure 4.1. Any changes in *total doses* from 2004 – 2010 were attributable to environmental variability (in measurements of gamma dose rates); thereafter (2011–2013) relatively lower *total doses* were estimated due to a lower occupancy rate over local beaches.

Source specific assessments for high-rate terrestrial food consumption, and from external exposure for turf cutting over salt marsh, give exposures that were less than the *total dose* (Table 4.1). The estimated dose from terrestrial food consumption, and turf cutting over salt marsh, in 2015 were less than 0.005 mSv and 0.016 mSv, respectively (both unchanged from 2014). The dose to a local fisherman, who was considered to consume a large amount of seafood and was exposed to external radiation over intertidal areas, was 0.031 mSv in 2015, which was

approximately 3 per cent of the dose limit for members of the public of 1 mSv (Table 4.1), and similar to that in 2014 (0.032 mSv).

### Gaseous discharges and terrestrial monitoring

In 2015, discharges of carbon-14 and argon-41 increased (by small amounts) at Heysham 2, in comparison to releases in 2014. The monitoring programme for the effects of gaseous disposals was similar to that for other power stations. Data for 2015 are given in Table 4.6(a). The effects of gaseous disposals were difficult to detect in 2015. Carbon-14 was detected in locally produced milk at concentrations close to expected background values. In 2015, the maximum caesium-137 concentration in milk was positively detected (unlike in previous years). Small enhancements of concentrations of sulphur-35 were measured in some food samples (reported as just above the less than value), but activities of cobalt-60 are all reported as less than values.

### Liquid waste discharges and aquatic monitoring

Discharges of tritium decreased (by small amounts) from Heysham 1 and Heysham 2, compared with those in 2014. The monitoring programme for the effects of liquid disposals included sampling of fish, shellfish, sediment, seawater and measurements of gamma dose rates and for completeness the data considered in this section include all of those for Morecambe Bay. A substantial part of the programme is in place in order to monitor the effects of Sellafield disposals. The results for 2015 are given in Tables 4.6(a) and (b). In general, similar levels to those for 2014 were observed and the effect of liquid disposals from Heysham was difficult to detect above the Sellafield background. Concentrations of tritium in flounder, shrimps and mussels were not sufficiently high to demonstrate that any originated as a result of discharges from Heysham. Concentrations of technetium-99 in marine samples remained at levels typical of recent years, caused by discharges from Sellafield. In 2015, strontium-90 concentrations were detected at low levels (close to, or just above, the less than value) in food samples. Gamma dose rates over intertidal sediment were generally similar to measurements in recent years.

## 4.6 Hinkley Point, Somerset



The Hinkley Point Power Station sites are situated on the Somerset coast, west of the River Parrett estuary. There are two separate A and B stations that include two Magnox reactors and two AGRs, respectively. Hinkley Point A

started electricity generation in 1965 and ceased in 2000. This station completed de-fuelling in 2004 and is undergoing decommissioning. The Hinkley Point A site will enter the Care and Maintenance phase by the year 2025. Final site clearance is expected to commence in 2081 and be achieved by 2090 (NDA, 2016a). It is estimated that power generation will continue at Hinkley Point B until at least 2023. A single environmental monitoring programme covers the effects of the two power stations.

In 2013, the Secretary of State for Energy and Climate Change granted a planning consent order to EDF Energy to build and operate Hinkley Point C and associated development. The decision follows the submission of EDF Energy's application to the Infrastructure Planning Commission (now the Planning Inspectorate) in 2011. More information concerning decisions on environmental permit applications for the proposed nuclear power plant can be found at: <https://www.gov.uk/government/collections/hinkley-point>. In 2014, ONR published its assessment of a Pre-Construction Safety Report submitted by NNB GenCo Limited for the Hinkley Point C licensed site: <http://www.onr.org.uk/hinkley-point-c/assessment-reports.htm#preconstruction>.

In April 2015, NNB GenCo Limited informed ONR, due to the continued uncertainty surrounding the Final Investment Decision date, that the EDF board had requested that the project review its spend profile. NNB GenCo Limited announced a pause to the overall programme for the Hinkley Point C Project and has reviewed its current activities to ensure that they are commensurate with the revised project plan. Until it is ready to fully remobilise the project, NNB GenCo is currently focussing its resources on activities which support preparation for start of construction.

The most recent habits survey was conducted in 2010 (Clyne *et al.*, 2011a).

### Doses to the public

In 2015, the *total dose* from all pathways and sources of radiation was 0.016 mSv (Table 4.1), or less than 2 per cent

of the dose limit, and down from 0.022 mSv in 2014. An adult who spent a large amount of time over sediments was the representative person. The decrease in *total dose* was mostly due to lower dose rates over sediment in 2015, in comparison to those in 2014. The trend in *total dose* over the period 2004 – 2015 is given in Figure 4.1. In 2010, the decrease in *total dose* (and continued thereafter) was attributed to relatively lower gamma dose rates over local beaches.

A source specific assessment for a high-rate consumer of locally grown food gives an exposure that was less than the *total dose* (Table 4.1). The dose to this consumer was less than 0.005 mSv in 2015. The decrease in dose (from 0.010 mSv in 2014) was due to lower carbon-14 concentrations in milk. The dose to a local fisherman, who consumed a large amount of seafood and was exposed to external radiation over intertidal areas, was 0.023 mSv in 2015, which was approximately 2 per cent of the dose limit for members of the public of 1 mSv. The decrease in dose (from 0.032 mSv in 2014) was mostly due lower dose rates over mud and rock at Stolford in 2015. This estimate also includes the effects of discharges (historical) of tritium and carbon-14 from Cardiff and uses an increased tritium dose coefficient (see Appendix 1).

### Gaseous discharges and terrestrial monitoring

Gaseous radioactive waste is discharged via separate stacks to the local environment. Discharges of carbon-14 and argon-41 from Hinkley Point B decreased by a small amount, in comparison to releases in 2014; other discharges of radionuclides were broadly comparable (including those from Hinkley Point A). Analyses of milk and crops were undertaken to measure activity concentrations of tritium, carbon-14, sulphur-35 and gamma emitters. Local reservoir water samples were also taken and analysed. Data for 2015 are given in Table 4.7(a). Activity concentrations of tritium and gamma emitters (including caesium-137) in terrestrial materials are reported as less than values. Sulphur-35 from Hinkley Point B was detected at low concentrations in some of the food samples (excluding milk). Carbon-14 was detected in locally produced milk at concentrations close to expected background values in 2015, and decreased in comparison to those in 2014. Tritium, gross alpha and gross beta concentrations in reservoir water were below the investigation levels for drinking water in the European Directive 2013/51.


### Liquid waste discharges and aquatic monitoring

Regulated discharges of radioactive liquid effluent from both power stations are made via separate outfalls into the Bristol Channel. Discharges of tritium and sulphur-35 from Hinkley Point B decreased by a small amount in 2015,

in comparison to releases in 2014. Analyses of seafood and marine indicator materials and measurements of external radiation were conducted over intertidal areas. Measurements of tritium and carbon-14 are made primarily to establish the local effects of historical discharges from the GE Healthcare Limited plant at Cardiff.

The environmental results for 2015 are given in Tables 4.7 (a) and (b). Activity concentrations observed in seafood and other materials (including tritium in fish and shellfish) from the Bristol Channel were generally similar to those in recent years. Concentrations of other radionuclides in the aquatic environment represent the combined effect of releases from these stations, plus other establishments that discharge into the Bristol Channel. Other contributors to the aquatic environment are Sellafield, weapons testing and Chernobyl fallout. Due to the low concentrations detected, it is generally difficult to attribute the results to a particular source. The concentrations of transuranic nuclides in seafoods were of negligible radiological significance. There is now limited evidence to suggest that caesium-137 concentrations in sediment have been generally decreasing over the most recent years (Figure 4.2). Overall, gamma dose rates over intertidal sediment in 2015 were generally similar, although the dose rates at Stolford (mud and rocks) decreased, in comparison to those in 2014.

## 4.7 Sizewell, Suffolk



The two Sizewell Power Stations are located on the Suffolk coast, near Lowestoft. Sizewell A is a Magnox twin reactor site. Sizewell B, powered by one reactor, is the only commercial PWR power station in the UK. The B station began

operation in 1995 and it is estimated that it will end power generation by 2035. Sizewell A power station ceased electricity generation in 2006. De-fuelling commenced in 2007 and was completed in 2014. The Sizewell A site will enter the Care and Maintenance phase by the year 2027. Final site clearance is expected to commence in 2088 and be achieved by 2097 (NDA, 2016a).

In June 2015, a habits survey was conducted to determine the consumption and occupancy rates by members of the public (Garrod *et al.*, 2016). Decreases in the fish, crustacean and mollusc consumption rates have been observed, together with an increase in the occupancy rate, in comparison with those of the previous survey in 2010. External exposure rates due to occupancy on a boat over mud decreased significantly. Revised figures for

consumption rates, together with occupancy rates, are provided in Appendix 1 (Table X2.2).

## Doses to the public

The *total dose* from all pathways and sources was 0.021 mSv in 2015 (Table 4.1) or approximately 2 per cent of the dose limit, and similar to the value of 0.020 mSv in 2013. As in recent years, the dominant contribution to *total dose* was from direct radiation and the representative person was an adult living in the vicinity of the site. Dose from this pathway has reduced by a factor of three since Sizewell A ceased generation in 2006. The trend in *total dose* over the period 2004 – 2015 is given in Figure 4.1. The *total dose* declined at the end of 2006, following the closure of the Magnox reactors at Sizewell A, thereafter any variations were due to the change in the contribution from direct radiation from the site.

Source specific assessments for both a high-rate consumer of locally grown foodstuffs, and of fish and shellfish, and of external exposure for houseboat occupancy, give exposures that were less than the *total dose* in 2015 (Table 4.1). The dose to a consumer of locally grown foods was less than 0.005 mSv. The dose to a houseboat dweller from external exposure was less than 0.005 mSv in 2015 (as in 2014) because of the significant decrease in occupancy rates from the revised habits data.

## Gaseous discharges and terrestrial monitoring

Gaseous wastes are discharged via separate stacks to the local environment. The results of the terrestrial monitoring in 2015 are shown in Table 4.8 (a). Gamma-ray spectrometry and analysis of tritium, carbon-14 and sulphur-35 in milk and crops generally showed very low concentrations of artificial radionuclides near the power stations in 2015. Carbon-14 concentrations were detected in locally produced milk, just above expected background concentrations, whilst sulphur-35 was positively detected at a very low concentration in one food sample (barley). Tritium concentrations in local freshwater were all low, although those measured at the Leisure Park were positively detected above the less than value (as in 2014 and 2010). Tritium, gross alpha and gross beta concentrations in surface water were below the investigation levels for drinking water in the European Directive 2013/51.


## Liquid waste discharges and aquatic monitoring

Regulated discharges of radioactive liquid effluent are made via outfalls to the North Sea. Caesium-137 and tritium discharges decreased from Sizewell A and Sizewell B, respectively, in comparison to those in 2014. The

decrease in tritium discharges, over period of sustained power generation (there was no forced or re-fuelling outage in 2015), was attributed to removal of the secondary sources from the Sizewell B reactor core. In the aquatic programme, analysis of seafood, sediment, and seawater, and measurements of gamma dose rates were conducted in intertidal areas. Data for 2015 are given in Tables 4.8(a) and (b). Concentrations of artificial radionuclides were low and mainly due to the distant effects of Sellafield discharges and to weapons testing. Tritium concentrations in seafood are all reported as less than values. Caesium-137 concentrations in sediment have remained low over the last decade (Figure 4.2). Overall, gamma radiation dose rates over intertidal areas were difficult to distinguish from the natural background, although the dose rates at Southwold Harbour increased in comparison to those in 2014 (most likely due to natural variation).

## SCOTLAND

### 4.8 Chapelcross, Dumfries and Galloway



Chapelcross was Scotland's first commercial nuclear power station and has four Magnox reactors located near the town of Annan in Dumfries and Galloway. After 45 years of continuous operation, electricity generation ceased

in 2004 and the station has been undergoing decommissioning. De-fuelling of the reactors began in 2008 and was completed during 2013. The major hazards remaining on the site will now be addressed early during decommissioning. The site will enter the Care and Maintenance phase by the year 2028. Current plans are for the site to be de-licensed (released from regulatory control). Final site clearance is expected to commence in 2085 and be achieved by 2095 (NDA, 2015).

Habits surveys have been undertaken to investigate aquatic and terrestrial exposure pathways. In 2015, a habits survey was conducted to determine the consumption and occupancy rates by members of the public (SEPA, *in press*). Increases in the fish, mollusc and crustacean consumption rates and the handling rate for fishing gear have been observed, together with decreases in the wildfowl consumption and occupancy rates, in comparison with those of the previous survey in 2010. Revised figures for consumption rates, together with occupancy rates, are provided in Appendix 1 (Table X2.2). In 2012, a habits survey was also conducted to determine the consumption

and occupancy rates by members of the public on the Dumfries and Galloway coast (Garrod *et al.*, 2013a). The results of this survey are used to determine the potential exposure pathways relating to permitted liquid discharges from the Sellafield nuclear licensed site in Cumbria (see Section 2.3.1).

### Doses to the public

The *total dose* from all pathways and sources of radiation is assessed to have been 0.022 mSv in 2015 (Table 4.1), which was approximately 2 per cent of the dose limit, and up from 0.014 mSv in 2014. In 2015, an infant consuming milk was the representative person, and was a change from that in 2014 (adult who spent a large amount of time over sediments). The increase in *total dose*, and change in the representative person, (from 2014) was mostly attributed to the inclusion of a less than value for americium-241 activity in food in the 2015 assessment. In line with the rules on use of the results for dose calculations, americium-241 was included because detectable activity was observed in one reported sample (soil, maximum) from the terrestrial environment (unlike in 2014) in 2015. The trend in *total dose* over the period 2004 – 2015 is given in Figure 4.1. *Total doses* remained broadly similar from year to year, and were low.

Source specific assessments for a high-rate consumer of locally grown food and seafood (crustaceans), and for a salmon and wildfowl consumer, give exposures that were less than the *total dose* in 2015 (Table 4.1). The dose for the terrestrial food consumer was estimated to be 0.017 mSv in 2015. The reason for the increase in dose (from 0.008 mSv in 2014) is the same as that contributing to the maximum *total dose* in 2015. The dose for the salmon, mollusc and wildfowl consumer was 0.009 mSv in 2015 and similar to that in 2014 (0.010 mSv); this change includes a smaller contribution from gamma dose rates (mostly due to lower dose rates over salt marsh at the pipeline and Dornoch Brow), and includes a larger contribution from americium-241 in molluscs (from the revision of the habits information), in 2015.

A consideration of the discharges from Chapelcross indicates that they contribute a very small fraction of the dose to the local population from seafood consumption and occupancy over salt marsh; the greater proportion of the dose can be attributed to the discharges from Sellafield.

### Gaseous discharges and terrestrial monitoring

Gaseous radioactive waste is discharged via stacks to the local environment. In 2015, discharges of "all other nuclides" increased by a small amount, in comparison to releases in 2014.

Terrestrial monitoring consisted of the analysis of a variety of foods, including milk, fruit, crops and game, as well as grass, soil and freshwater samples, for a range of radionuclides. Air samples at three locations were also monitored to investigate the inhalation pathway.

The results of terrestrial food and air monitoring in 2014 are given in Tables 4.9(a) and (c). The activity concentrations of radionuclides in milk and grass were generally similar to those observed in previous years. As in 2014, carbon-14 concentrations in milk were similar to the default values used to represent background levels. Sulphur-35 and americium-241 concentrations in all terrestrial food samples are reported as less than values in 2015. In previous years, the tritium results in terrestrial samples have shown the effects of discharges from Chapelcross. In 2015, tritium concentrations over the range of food samples were all reported as less than values, whilst most other samples are reported as just above or as less than values. As in recent years, the level of tritium was measured above the detection limit in one freshwater sample (Gullielands Burn). Activity concentrations in air samples at locations near to the site are reported as less than values (or close to the less than value) (Table 4.9(c)).

### Liquid waste discharges and aquatic monitoring

Radioactive liquid effluents are discharged to the Solway Firth. Samples of seawater and *Fucus vesiculosus*, as environmental indicators, were collected in addition to seafood, sediments and dose rates. Data for 2015 are given in Tables 4.9(a) and (b). Concentrations of artificial radionuclides in marine materials in the Chapelcross vicinity are mostly due to the effects of Sellafield discharges and are consistent with values expected at this distance from Sellafield. Concentrations of most radionuclides remained at similar levels to those detected in recent years. Low concentrations of cobalt-60, europium-154 and europium-155 were positively detected (just above the less than value) in sediment samples. As in previous years, concentrations of caesium-137, plutonium radionuclides and americium-241 were enhanced in sediment samples, although the plutonium radionuclides and americium-241 concentrations taken close to the pipeline were much lower (in comparison to those in recent years), in 2015. Technetium-99 concentrations in seaweed and mollusc samples were higher, by small amounts, in comparison to those in 2014, whilst concentrations in biota are reported as less than values in 2015. Concentrations of caesium-137 in sediments, largely due to Sellafield, are generally in decline over the last decade (Figure 4.2). In 2015, gamma dose rates (where comparisons can be made) were generally similar to those in 2014; albeit with small decreases over salt marsh at the pipeline and Dornoch Brow. Measurements of the contact beta dose rate on stake nets are reported as less than values in 2015.

Between 1992 and 2009, a number of particles were found at the end of the discharge outfall consisting of lime-scale originating from deposits within the pipeline. Magnox Limited continues to monitor this area frequently and no particles were found during 2015 (as for the interim years). The relining of the pipeline and grouting at strategic points, which was undertaken in 2009/2010, has reduced the potential for particles to be released.

## 4.9 Hunterston, North Ayrshire



Hunterston Power Station is located on the Ayrshire coast near West Kilbride. At this location there are two separate nuclear power stations – Hunterston A and Hunterston B.

Hunterston A was powered

by twin Magnox reactors until it ceased electricity production in 1990 and is now being decommissioned by Magnox Limited. Defueling was completed in 1995. Decommissioning activities have focussed on two key areas during 2015, these were: the ongoing draining and cleaning of the cartridge (nuclear fuel) cooling pond; and making progress towards ensuring that all higher activity waste is stored in a passively safe manner. The site will enter the Care and Maintenance phase by the year 2022. Current plans are for the Hunterston A site to be de-licensed (released from regulatory control) with final site clearance to commence in 2071 and be achieved by 2080 (NDA 2016a).

The majority of the radioactivity in liquid effluent discharged from the Hunterston A site over the last few years has arisen from the cartridge cooling pond. The draining of the cartridge cooling pond was largely completed during 2015 and accordingly discharges of effluent from the cooling pond are coming to an end. This is reflected in the quantity of radioactivity discharged in liquid effluent from the site which is approximately 50 per cent lower than last year.

In terms of safe management of legacy higher activity waste, Magnox Limited are in the process of constructing and commissioning two new facilities at Hunterston A; the solid intermediate level waste encapsulation plant (SILWE) and the wet intermediate level waste recovery and encapsulation plant (WILREP). The legacy higher activity waste, which is present at the Hunterston A site, will be processed through these facilities and made passively safe by encapsulating it in a grout mixture. The encapsulated waste will then be transferred to the Intermediate Level Radioactive Waste Store (ILWS) for storage.

Hunterston B is powered by a pair of AGRs. Due to issues with the boiler tubes a number of years ago, the station is limited to approximately 80 per cent of its design output. The life of the station has been extended twice, and the current end of generation is set for 2023.

In late 2013, EDF Energy applied to SEPA to vary Hunterston B's authorisation in order to allow radioactive waste to be disposed of by transfer to any waste permitted person, both within the UK and overseas, and to be able to accept radioactive waste from other EDF Energy stations for the purposes of bulking up low volume wastes before final disposal. The proposed variation does not include any changes to the station's authorised limits for discharges to the environment. During 2015, the application was still being determined by SEPA.

Hunterston B exceeded its authorised tidal window by approximately 26 minutes on one occasion in November 2015, due to human error. The discharge was stopped, whilst the tide was still ebbing, and there has been no discernible environmental impact as a result of the exceedance. SEPA subsequently issued the station with a Warning Letter in response to this contravention of the station's authorisation.

Environmental monitoring in the area considers the effects of both Hunterston A and Hunterston B sites together. The most recent habits survey was undertaken in 2012, to determine the consumption and occupancy rates by members of the public (Rumney *et al.*, 2013a).

## Doses to the public

The *total dose* from all pathways and sources of radiation is assessed to have been 0.025 mSv in 2015 (Table 4.1), which was less than 3 per cent of the dose limit, and similar to 0.021 mSv in 2014. The apparent increase in dose was attributed to the inclusion of the less than value for americium-241 activity in food in the 2015 assessment. In line with the rules on use of the results for dose calculations, americium-241 was included because detectable activity was observed in other samples (soil and grass) from the terrestrial environment in 2015. The actual doses received were likely to have been less than 0.025 mSv because the assessment includes less than values for radionuclide concentrations in food. The representative person was a 1 year-old infant consuming milk at high-rates in 2015 (as opposed to a prenatal child exposed to direct radiation in 2014). The trend in *total dose* over the period 2004 – 2015 is given in Figure 4.1. The decrease in *total dose* in recent years reflected a downward trend in the reported direct radiation.

Source specific assessments for both a high-rate consumer of locally grown food, and of local seafood, give exposures that were less than the *total dose* in 2015 (Table 4.1). The estimated dose to a terrestrial food consumer was 0.019 mSv in 2015, which was less than 2 per cent of the

dose limit for members of the public of 1 mSv. The reason for the apparent increase in dose from 0.009 mSv (in 2014) was the same as that contributing to the maximum *total dose*. The dose to a fish and shellfish consumer was less than 0.005 mSv.

## Gaseous discharges and terrestrial monitoring

Gaseous discharges are made via separate discharge points from the Hunterston A and Hunterston B stations. Discharges of carbon-14 increased by a small amount from Hunterston B, in comparison to those releases in 2014. There is a substantial terrestrial monitoring programme which includes the analyses of a comprehensive range of wild and locally produced foods. In addition, air, freshwater, grass and soil are sampled to provide background information. The results of terrestrial food and air monitoring in 2015 are given in Tables 4.10(a) and (c). The concentrations of radionuclides in air, milk, crops and fruit were generally low and, where comparisons can be drawn, similar to those in previous years. The caesium-137 concentration in honey was positively detected ( $2.7 \text{ Bq kg}^{-1}$ ) in 2015 (slightly elevated in comparison to recent years), whilst sulphur-35 was detected at a very low concentration in one food sample (wild blackberries). Some carbon-14 concentrations in foodstuffs were higher than the default values used to represent background levels (including honey, vegetables and domestic/wild fruit). Activity concentrations in air at locations near to the site are reported as less than values (or close to the less than value) (Table 4.10(c)).

## Liquid waste discharges and aquatic monitoring

Authorised liquid discharges from both Hunterston stations are made to the Firth of Clyde via the Hunterston B station's cooling water outfall. The main part of the aquatic monitoring programme consists of sampling of fish and shellfish and the measurement of gamma and beta dose rates on the foreshore. Samples of sediment, seawater and seaweed are analysed as environmental indicator materials.

The results of aquatic monitoring in 2015 are shown in Tables 4.10(a) and (b). The concentrations of artificial radionuclides in the marine environment are predominantly due to Sellafield discharges, the general values being consistent with those to be expected at this distance from Sellafield. The reported concentrations of technetium-99 from Sellafield in crabs and lobsters around Hunterston continued to remain low in 2015 and were generally similar to those reported in previous years. Small concentrations (above the less than value) of activation products (e.g. cobalt-60 and silver-110m) were also detected in seaweed, which were likely to have originated from the site, but these were of negligible radiological significance. Gamma

and beta dose rates were generally similar to those in 2014. Caesium-137 concentrations in sediment have remained low over the last decade (Figure 4.2).

#### 4.10 Torness, East Lothian



Torness Power Station is located near Dunbar on the east coast of Scotland. This station, which is powered by two AGRs, began operation at the end of 1987 and it is currently scheduled to cease generation in 2030

(extended from 2023). Disposals and discharges of radioactive waste from the site are made in accordance with RSA 93 authorisation issued to the site by SEPA in 2007.

In December 2013, EDF Energy applied to SEPA to vary the authorisation for Torness in order to allow radioactive waste to be disposed of by transfer to any waste permitted person, both within the UK and overseas, and to be able to accept radioactive waste from other EDF Energy stations for the purposes of bulking up low volume wastes before final disposal. The proposed variation does not include any changes to the site's authorised limited for discharges to the local environment. SEPA continued its determination of the application during 2015.

EDF Energy is continuing with its programme to reduce carbon deposition within the reactor and has continued to inject carbonyl sulphide (COS) into both reactors during 2015. This process was started in 2011 and after the initial expected increase in sulphur-35 levels discharged to the local environment, via the liquid and gaseous routes, the levels have stabilised. In 2015, discharges of sulphur-35 to the local environment were similar to those in 2014 and remained within the authorised limits. The gaseous and liquid discharges from the site are given in Appendix 2.

The most recent habits survey was undertaken in 2011 (Clyne *et al.*, 2013b).

#### Doses to the public

In 2015, the *total dose* from all pathways and sources of radiation was 0.020 mSv (Table 4.1) or 2 per cent of the dose limit, and unchanged from the previous four years. Direct radiation was the dominant contributor to the dose and the representative person was an adult. The trend in *total dose* over the period 2004 – 2015 is given in

Figure 4.1. *Total doses* remained broadly similar from year to year, and were low.

Source specific assessments for both a high-rate consumer of locally grown foods and of local fish and shellfish give exposures that were less than the *total dose* in 2015 (Table 4.1). The estimated dose to a terrestrial food consumer was 0.006 mSv, which was approximately 0.5 per cent of the dose limit for members of the public of 1 mSv, and unchanged from recent years. The dose to a fish and shellfish consumer was less than 0.005 mSv.

#### Gaseous discharges and terrestrial monitoring

A variety of foods, including milk, crops, fruit, and game as well as grass, soil and freshwater samples, were measured for a range of radionuclides. Air sampling at three locations was undertaken to investigate the inhalation pathway. The results of terrestrial food and air monitoring in 2015 are given in Tables 4.11(a) and (c). Activity concentrations in many terrestrial foods are reported as less than values (or close to the less than value). As in recent years, the effects of discharges from the power station were not observed for concentrations of sulphur-35, which are reported as less than values in terrestrial foods and environmental indicator materials. In 2015, caesium-137 in honey (not sampled in recent years) was positively detected at a low concentration (4.3 Bq kg<sup>-1</sup>). Americium-241 concentrations, measured by gamma-ray spectrometry, are also reported as less than values. Measured concentrations of radioactivity in air at locations near to the site are reported as less than values (or close to the less than value) (Table 4.11(c)).

#### Liquid waste discharges and aquatic monitoring

Liquid radioactive wastes are discharged to the Firth of Forth. Samples of seawater, sediment and *Fucus vesiculosus*, as useful environmental indicators, were collected in addition to seafood. Measurements were also made of gamma dose rates over intertidal areas, supported by analyses of sediment, and beta dose rates on fishing gear.

The results of the aquatic monitoring in 2015 are shown in Tables 4.11(a) and (b). Concentrations of artificial radionuclides were mainly due to the distant effects of Sellafield discharges and to weapon testing and Chernobyl fallout. In 2015, americium-241 concentrations in nephrops (from Dunbar Bay) were similar to those in previous years (although elevated in 2014). As in recent years, a few very low concentrations of activation products were detected in environmental indicator samples. These were likely to have originated from the station. Technetium-99 concentrations in marine samples were similar to those in 2014. In 2015, a caesium-137

concentration was slightly elevated in a sediment sample (Heckies Hole). Overall, caesium-137 concentrations in sediments have remained low over the last decade (Figure 4.2). Beta radiation from fishermen's pots are reported as less than values. Gamma dose rates over intertidal areas were generally indistinguishable from natural background and were similar to those measured in recent years.

## WALES

### 4.11 Trawsfynydd, Gwynedd



Trawsfynydd Power Station is located inland, on the northern bank of a lake in the heart of Snowdonia National Park, North Wales and was powered by twin Magnox reactors. Trawsfynydd ceased to generate electricity in 1991.

De-fuelling of the reactors was completed in 1995 and the station is being decommissioned. The focus for the site is now the completion of decommissioning projects. The Trawsfynydd site continues to prepare for entry into the Care and Maintenance phase. Final site clearance is expected to commence in 2078 and be achieved by 2083 (NDA, 2016a). The most recent habits survey was undertaken in 2005 (Tipple *et al.*, 2006).

#### Doses to the public

The *total dose* from all pathways and sources of radiation was 0.014 mSv in 2015 (Table 4.1), which was approximately 1 per cent of the dose limit, and similar to that in 2014 (0.013 mSv). The representative person in 2015 was an infant living near to the site, whose dose was from the consumption of milk at high-rates and direct radiation (from the site). The trend in *total dose* over the period 2004 – 2015 is given in Figure 4.1. *Total doses* remained broadly similar from year to year, and were low.

A source specific assessment for an angler (who consumes large quantities of fish and spends long periods of time in the location being assessed) gives an exposure that was less than the *total dose* in 2015 (Table 4.1). The dose to an angler was 0.007 mSv in 2015, which was approximately 0.5 per cent of the dose limit for members of the public of 1 mSv, and unchanged from 2014. The observed activity concentrations in lake sediments are used as the basis for external radiation calculations in view of the difficulty in establishing the increase in measured dose rates above natural background levels. The dose to an

infant consuming terrestrial food was 0.016 mSv, or less than 2 per cent of the dose limit. The dose in 2014 was 0.025 mSv, and the decrease in dose was mostly due to a lower reported less than value for the americium-241 concentration in milk in 2015, in comparison to that in 2014.

#### Gaseous discharges and terrestrial monitoring

Discharges of tritium decreased by a small amount, in comparison to releases in 2014. The results of the terrestrial programme, for local milk, fruit and grass samples in 2015, are shown in Table 4.12(a). Results from surveys, for activity concentrations in sheep samples, are available in earlier RIFE reports (e.g. Environment Agency, FSA, NIEA, NRW and SEPA, 2014). Concentrations of activity in all terrestrial samples were low. Carbon-14 concentrations in foodstuffs (milk and blackberries) were higher than the default values used to represent background levels. As in previous years, measured activities for caesium-137 are reported as less than values (or close to the less than value). The most likely source of small amounts of caesium-137 is fallout from Chernobyl and weapon tests, though it is conceivable that a small contribution may be made by re-suspension of lake activity. In recognition of this potential mechanism, monitoring of transuranic radionuclides was also conducted in a food sample (and in animal samples in previous years). In 2015, detected activities in fruit were low and generally similar to observations in other areas of England and Wales, where activity was attributable to weapon test fallout. There was no evidence of re-suspension of activity in sediment from the lake shore contributing to increased exposure from transuranic radionuclides in 2015.

#### Liquid waste discharges and aquatic monitoring

Discharges of liquid radioactive waste are made to a freshwater lake making the power station unique in UK terms. Discharges of tritium and “other radionuclides” decreased by a small amount in comparison to releases in 2014. The aquatic monitoring programme was directed at consumers of freshwater fish caught in the lake and external exposure over the lake shoreline; the important radionuclides are radiocaesium and, to a lesser extent, strontium-90. Freshwater and sediment samples are also analysed. Habits surveys have established that the species of fish regularly consumed are brown and rainbow trout. Most brown trout are indigenous to the lake but rainbow trout are introduced from a hatchery. Because of the limited period that they spend in the lake, introduced fish generally exhibit lower radiocaesium concentrations than indigenous fish.

Data for 2015 are given in Tables 4.12(a) and (b). The majority of activity concentrations in fish and sediments

result from historical discharges. The concentration of caesium-137 in fish (brown trout) is the lowest reported value, and concentrations in water are reported as less than values, in 2015. Concentrations in the water column are predominantly maintained by processes that release activity (such as remobilisation) from near surface sediments. Low concentrations of other radionuclides including transuranics were also detected, particularly in lake sediments; in previous years' monitoring, it has been demonstrated that these concentrations increase with depth beneath the sediment surface. Overall, caesium-137 concentrations in the lake sediments were decreased (by small amounts), in comparison to those in 2014. In 2015, sediment concentrations of strontium-90, plutonium-239+240 and americium-241 at one location (fish farm) were lower than those in 2014, but overall, sediment activity concentrations in 2015 were similar to those in other recent years. Strontium-90 and transuranic concentrations in fish continued to be very low in 2015 and it is the effects of caesium-137 that dominate the fish consumption and external radiation pathways.

In the lake itself, there remains clear evidence of activity concentrations from the site's liquid discharges. However, gamma dose rates found on the shoreline (where anglers fish) were difficult to distinguish from background levels, and rates in 2015 were similar to those in 2014. The predominant radionuclide was caesium-137. The time trends of concentrations of caesium-137 in sediments and discharges are shown in Figure 4.3. A substantial decline in levels was observed in the late 1990s in line with reducing discharges. In the earlier part of the last decade, the observed concentrations were mainly affected by sample variability. In the latter part of the last decade, with sustained reductions in discharges of caesium-137, there was a general progressive decrease in these concentrations in sediments, with the lowest concentrations reported in 2010. In years thereafter, there has been an overall small increase in activity concentrations, but activities generally decreasing again from the small peak in discharge in 2012.

#### 4.12 Wylfa, Isle of Anglesey



Wylfa Power Station is located on the north coast of Anglesey and has two Magnox reactors (Reactor 1 and 2). It was the last and largest power station of its type to be built in the UK and commenced electricity

generation in 1971 and ceased in December 2015, after a five-year lifetime extension (Reactor 1). Reactor 2 at the Wylfa site ceased generating electricity in 2012. In 2013, a

decision report was published by ONR, issuing consent for Magnox Limited to start decommissioning Wylfa power station within five years (ONR, 2013). Reactor defueling started in 2015. Final site clearance is expected to commence in 2096 and be achieved by 2105 (NDA, 2016a). The most recent habits survey was undertaken in 2013 (Garrod *et al.*, 2014).

#### Doses to the public

The *total dose* from all pathways and sources of radiation was 0.013 mSv in 2015 (Table 4.1), which was approximately 1 per cent of the dose limit, and up from 0.007 mSv in 2014. In 2015, the representative person was an infant living near to the site, and a change from that in 2014 (an adult who spends a large amount of time over sediments). The increase in *total dose* (from 2014) and change in the representative person was mostly due to a higher estimate of direct radiation from the site in 2015. The trend in *total dose* over the period 2004 – 2015 is given in Figure 4.1. *Total doses* remained broadly similar from year to year, and were generally very low.


A source specific assessment for a high-rate consumer of both locally grown foods gives an exposure that was less than the *total dose* in 2015 (Table 4.1). The dose to a consumer of locally grown foods was less than 0.005 mSv and unchanged from 2014. The dose to a high-rate consumer of fish and shellfish (including external radiation) was 0.007 mSv. The reason for the small decrease in dose in 2015 (from 0.010 mSv in 2014) was mostly because gamma dose rates were measured on different types of ground type from one year to the next and, to a lesser extent, a reduction in the reported less than value for americium-241 in fish in 2015.

#### Gaseous discharges and terrestrial monitoring

Discharges of tritium decreased, and carbon-14, sulphur-35 and argon-41 increased, in comparison to releases in 2014. The focus of the terrestrial sampling was for the content of tritium, carbon-14 and sulphur-35 in milk and crops. Data for 2015 are given in Table 4.13(a). Sulphur-35 was detected at a very low concentration in food samples (barley and potatoes). Carbon-14 was detected in locally produced milk at concentrations close to expected background concentrations. Tritium, gross alpha and gross beta concentrations in surface water (public supply) were below the investigation levels for drinking water in the European Directive 2013/51.

#### Liquid waste discharges and aquatic monitoring

Discharges of tritium decreased, in comparison to releases in 2014. The monitoring programme for the effects of



**Figure 4.3.** Caesium-137 liquid discharge from Trawsfynydd and concentration in sediment in Trawsfynydd lake, 1995-2015

liquid disposals included sampling of seafood, sediment, seawater and measurements of gamma dose rates. The results of the programme in 2015 are given in Tables 4.13 (a) and (b). The data for artificial radionuclides related to the Irish Sea continue to reflect the distant effects of Sellafield discharges. The activity concentrations in 2015 were similar to those in 2014, including technetium-99

derived from Sellafield. Caesium-137 concentrations in sediment have remained low over the last decade (Figure 4.2). Where comparisons can be made (from similar ground types and locations), gamma dose rates were generally similar in comparison to those in recent years.

**Table 4.1. Individual doses – nuclear power stations, 2015**

Site	Representative person <sup>a</sup>	Exposure, mSv per year					
		Total	Fish and shellfish	Other local food	External radiation from intertidal areas or the shoreline	Gaseous plume related pathways	Direct radiation from site
<b>England</b>							
Berkeley and Oldbury							
<b>Total dose – all sources</b>	<b>Infant milk consumers</b>	<b>&lt;0.005</b>	-	<b>&lt;0.005</b>	-	-	-
Source specific doses	Seafood consumers	<0.005	<0.005	-	<0.005	-	-
	Houseboat occupants	0.006	-	-	0.006	-	-
	Infant inhabitants and consumers of locally grown food	0.005	-	0.005	-	<0.005	-
Bradwell							
<b>Total dose – all sources</b>	<b>Prenatal children of local inhabitants (0 - 0.25km)</b>	<b>0.017</b>	-	<b>&lt;0.005</b>	-	<b>&lt;0.005</b>	<b>0.017</b>
Source specific doses	Seafood consumers	<0.005	<0.005	-	<0.005	-	-
	Infant inhabitants and consumers of locally grown food	<0.005	-	<0.005	-	<0.005	-
Dungeness							
<b>Total dose – all sources</b>	<b>Local adult inhabitants (0.5 - 1km)</b>	<b>0.014</b>	<b>&lt;0.005</b>	-	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>0.013</b>
Source specific doses	Seafood consumers	0.006	<0.005	-	<0.005	-	-
	Houseboat occupants	<0.005	-	-	<0.005	-	-
	Infant inhabitants and consumers of locally grown food	0.005	-	<0.005	-	<0.005	-
Hartlepool							
<b>Total dose – all sources</b>	<b>Local adult inhabitants (0 - 0.25km)</b>	<b>0.022</b>	<b>&lt;0.005</b>	-	<b>0.012</b>	<b>&lt;0.005</b>	<b>0.010</b>
Source specific doses	Seafood consumers <sup>b</sup>	0.020	<0.005	-	0.020	-	-
	Infant inhabitants and consumers of locally grown food	<0.005	-	<0.005	-	<0.005	-
Heysham							
<b>Total dose – all sources</b>	<b>Adult mollusc consumers</b>	<b>0.023</b>	<b>0.012</b>	-	<b>0.010</b>	-	-
Source specific doses	Seafood consumers	0.031	0.012	-	0.019	-	-
	Turf cutters	0.016	-	-	0.016	-	-
	Infant inhabitants and consumers of locally grown food	<0.005	-	<0.005	-	<0.005	-
Hinkley Point							
<b>Total dose – all sources</b>	<b>Adult occupants over sediment</b>	<b>0.016</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>0.015</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>
Source specific doses	Seafood consumers	0.023	<0.005	-	0.022	-	-
	Infant inhabitants and consumers of locally grown food	<0.005	-	<0.005	-	<0.005	-
Sizewell							
<b>Total dose – all sources</b>	<b>Local adult inhabitants (0 - 0.25km)</b>	<b>0.021</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>0.020</b>
Source specific doses	Seafood consumers	<0.005	<0.005	-	<0.005	-	-
	Houseboat occupants	<0.005	-	-	<0.005	-	-
	Infant inhabitants and consumers of locally grown food	<0.005	-	<0.005	-	<0.005	-

**Table 4.1. continued**

Site	Representative person <sup>a</sup>	Exposure, mSv per year					
		Total	Fish and shellfish	Other local food	External radiation from intertidal areas or the shoreline	Gaseous plume related pathways	Direct radiation from site
<b>Scotland</b>							
Chapelcross							
<b>Total dose – all sources</b>	<b>Infant milk consumers</b>	<b>0.022</b>	<b>&lt;0.005</b>	<b>0.022</b>	<b>&lt;0.005</b>	-	-
Source specific doses	Salmon, mollusc and wildfowl consumers	0.009	<0.005	<0.005	<0.005	-	-
	Crustacean consumers	<0.005	<0.005	-	-	-	-
	Infant inhabitants and consumers of locally grown food	0.017	-	0.016	-	<0.005	-
Hunterston							
<b>Total dose – all sources</b>	<b>Infant milk consumers</b>	<b>0.025</b>	-	<b>0.025</b>	-	-	-
Source specific doses	Seafood consumers	<0.005	<0.005	-	<0.005	-	-
	Infant inhabitants and consumers of locally grown food	0.019	-	0.017	-	<0.005	-
Torness							
<b>Total dose – all sources</b>	<b>Local adult inhabitants (0.5 - 1km)</b>	<b>0.020</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	<b>0.020</b>
Source specific doses	Seafood consumers	<0.005	<0.005	-	<0.005	-	-
	Infant inhabitants and consumers of locally grown food	0.006	-	0.006	-	<0.005	-
<b>Wales</b>							
Trawsfynydd							
<b>Total dose – all sources</b>	<b>Infant local inhabitants (0.25 - 0.5km)</b>	<b>0.014</b>	-	<b>0.008</b>	-	<b>&lt;0.005</b>	<b>0.006</b>
Source specific doses	Anglers	0.007	0.005	-	<0.005	-	-
	Infant inhabitants and consumers of locally grown food	0.016	-	0.016	-	<0.005	-
Wylfa							
<b>Total dose – all sources</b>	<b>Infant local inhabitants (0.25 - 0.5km)</b>	<b>0.013</b>	-	<b>&lt;0.005</b>	-	<b>&lt;0.005</b>	<b>0.012</b>
Source specific doses	Seafood consumers	0.007	<0.005	-	0.005	-	-
	Infant inhabitants and consumers of locally grown food	<0.005	-	<0.005	-	<0.005	-

<sup>a</sup> The total dose is the dose which accounts for all sources including gaseous and liquid discharges and direct radiation. The total dose for the representative person with the highest dose presented. Other dose values are presented for specific sources, either liquid discharges or gaseous discharges, and their associated pathways. They serve as a check on the validity of the total dose assessment. The representative person is an adult unless otherwise stated

<sup>b</sup> Excluding possible enhancement of naturally occurring radionuclides. See Section 4

**Table 4.2(a). Concentrations of radionuclides in food and the environment near Berkeley and Oldbury nuclear power stations, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>3</sup> H	<sup>14</sup> C	<sup>99</sup> Tc	<sup>137</sup> Cs	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu
<b>Marine samples</b>								
Salmon	Beachley	1				0.19		
Mullet	River Severn	1	<25				0.24	
Elvers	River Severn	1				<0.10		
Shrimps	Guscar	2	25	29		0.33	0.00026	0.0018
Seaweed	Pipeline	2 <sup>E</sup>			<1.7	<0.72		
Sediment	Hills Flats	2 <sup>E</sup>				19		
Sediment	1 km south of Oldbury	2 <sup>E</sup>				21		
Sediment	2 km south west of Berkeley	2 <sup>E</sup>				24		
Sediment	Sharpness	2 <sup>E</sup>				20		
Seawater	Local beach	2 <sup>E</sup>				<0.19		

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm + <sup>244</sup> Cm	Gross alpha	Gross beta	
<b>Marine samples</b>								
Salmon	Beachley	1	<0.08					
Mullet	River Severn	1	<0.05					
Elvers	River Severn	1	<0.08					
Shrimps	Guscar	2	0.0019	*	0.000025			
Seaweed	Pipeline	2 <sup>E</sup>	<0.76					
Sediment	Hills Flats	2 <sup>E</sup>	<0.75					
Sediment	1 km south of Oldbury	2 <sup>E</sup>	<0.79					
Sediment	2 km south west of Berkeley	2 <sup>E</sup>	<0.80					
Sediment	Sharpness	2 <sup>E</sup>	<0.76					
Seawater	Local beach	2 <sup>E</sup>	<0.28				<1.6	6.5

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>137</sup> Cs	Gross alpha	Gross beta
<b>Terrestrial samples</b>								
Milk		4	<3.5	19	<0.21	<0.07		
Milk	max		<4.1	24	<0.28			
Apples		1	<2.8	14	<0.10	<0.05		
Wheat		1	<5.0	78	0.40	<0.04		
Freshwater	Gloucester and Sharpness Canal	2 <sup>E</sup>	<3.1		<0.38	<0.19	<0.058	<0.24
Freshwater	Public supply	2 <sup>E</sup>	<3.1		<0.25	<0.19	<0.048	<0.18

\* Not detected by the method used

<sup>a</sup> Except for milk and water where units are Bq l<sup>-1</sup>, and for sediment where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 4.2(b). Monitoring of radiation dose rates near Berkeley and Oldbury nuclear power stations, 2015**

Location	Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
1 km south of Oldbury	Mud and salt marsh	2	0.078
2 km south west of Oldbury	Mud and salt marsh	2	0.072
Guscar Rocks	Mud	1	0.081
Guscar Rocks	Mud and salt marsh	1	0.079
Lydney Rocks	Mud	2	0.096
Sharpness	Mud and salt marsh	2	0.072
Hills Flats	Mud and salt marsh	1	0.079
Hills Flats	Salt marsh	1	0.076

**Table 4.3(a). Concentrations of radionuclides in food and the environment near Bradwell nuclear power station, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>3</sup> H	<sup>90</sup> Sr	<sup>99</sup> Tc	<sup>137</sup> Cs	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu
<b>Marine samples</b>								
Bass	Pipeline	1				0.30		
Lobsters	West Mersea	1				0.05		
Native oysters	Blackwater Estuary	1				<0.11	0.00034	0.0019
Sea beet	Tollesbury	1			0.78	<0.05		
Seaweed	Waterside	2 <sup>E</sup>			1.3	<0.65		
Samphire	West Mersea	1 <sup>E</sup>			<0.24	<0.57		
Sediment	Bradwell Pipeline	2 <sup>E</sup>		<2.1		3.8		
Sediment	Bradwell Marina	2 <sup>E</sup>				2.5		
Sediment	Waterside	2 <sup>E</sup>		<3.5		5.2		
Sediment	Steeple	2 <sup>E</sup>				8.9		
Sediment	Maylandsea Bay	2 <sup>E</sup>				12		
Sediment	Blackwater	2 <sup>E</sup>				13		
Sediment	N side Blackwater Estuary	2 <sup>E</sup>		<2.0		7.8		
Sediment	Osea Causeway	2 <sup>E</sup>				2.5		
Sediment	Maldon	2 <sup>E</sup>		<2.0		14		
Sediment	Maldon (2)	2 <sup>E</sup>				41		
Sediment	Heybridge	2 <sup>E</sup>				13		
Sediment	Strood Channel	2 <sup>E</sup>				11		
Sediment	Tollesbury Boatyard	2 <sup>E</sup>				11		
Sediment	Tollesbury saltwater pool	2 <sup>E</sup>				11		
Sediment	West Mersea Beach Huts	2 <sup>E</sup>		<2.0		1.3		
Sediment	West Mersea Boatyard	2 <sup>E</sup>		<2.0		4.9		
Sediment	Pyefleet	2 <sup>E</sup>				6.9		
Sediment	Rowhedge	2 <sup>E</sup>				10		
Sediment	Alresford Creek	2 <sup>E</sup>				5.9		
Sediment	Brightlingsea Bateman's Tower	2 <sup>E</sup>				<0.18		
Sediment	Brightlingsea saltwater pool	2 <sup>E</sup>				<0.21		
Sediment	St Osyth boat lake	2 <sup>E</sup>				13		
Seawater	Bradwell Pipeline	2 <sup>E</sup>	<3.5			<0.21		
Seawater	Bradwell Marina	2 <sup>E</sup>				<0.20		
Seawater	Steeple	2 <sup>E</sup>				<0.23		
Seawater	Maylandsea Bay	2 <sup>E</sup>				<0.22		
Seawater	Blackwater	2 <sup>E</sup>				<0.19		
Seawater	Osea Causeway	2 <sup>E</sup>				<0.27		
Seawater	Maldon	2 <sup>E</sup>	<3.3			<0.30		
Seawater	Heybridge	2 <sup>E</sup>				<0.19		
Seawater	Strood Channel	2 <sup>E</sup>	<3.2			<0.26		
Seawater	Tollesbury Boatyard	2 <sup>E</sup>				<0.22		
Seawater	Tollesbury saltwater pool	2 <sup>E</sup>				<0.26		
Seawater	Pyefleet	2 <sup>E</sup>	<3.3			<0.26		
Seawater	Rowhedge	2 <sup>E</sup>				<0.22		
Seawater	Alresford Creek	2 <sup>E</sup>				<0.19		
Seawater	Brightlingsea Bateman's Tower	2 <sup>E</sup>				<0.26		
Seawater	Brightlingsea saltwater pool	2 <sup>E</sup>				<0.19		
Seawater	St Osyth boat lake	2 <sup>E</sup>				<0.22		

**Table 4.3(a). continued**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm + <sup>244</sup> Cm	Gross alpha	Gross beta
<b>Marine samples</b>							
Bass	Pipeline	1	<0.19				
Lobsters	West Mersea	1	<0.04				
Native oysters	Blackwater Estuary	1	0.0012	*	0.000027		
Sea beet	Tollesbury	1	<0.11				
Seaweed	Waterside	2 <sup>E</sup>	<0.63				
Samphire	West Mersea	1 <sup>E</sup>	<0.57				
Sediment	Bradwell Pipeline	2 <sup>E</sup>	<0.61				
Sediment	Bradwell Marina	2 <sup>E</sup>	<0.73			380	990
Sediment	Waterside	2 <sup>E</sup>	<1.1				
Sediment	Steeple	2 <sup>E</sup>	<1.2			380	900
Sediment	Maylandsea Bay	2 <sup>E</sup>	<1.6			300	930
Sediment	Blackwater	2 <sup>E</sup>	<1.6			370	910
Sediment	N side Blackwater Estuary	2 <sup>E</sup>	<1.5				
Sediment	Osea Causeway	2 <sup>E</sup>	<0.44			<120	360
Sediment	Maldon	2 <sup>E</sup>	<1.1				
Sediment	Maldon (2)	2 <sup>E</sup>	<1.7			310	1000
Sediment	Heybridge	2 <sup>E</sup>	<1.5			290	820
Sediment	Strood Channel	2 <sup>E</sup>	<1.6			260	840
Sediment	Tollesbury Boatyard	2 <sup>E</sup>	<1.6			310	1000
Sediment	Tollesbury saltwater pool	2 <sup>E</sup>	<0.54			<120	410
Sediment	West Mersea Beach Huts	2 <sup>E</sup>	<0.46				
Sediment	West Mersea Boatyard	2 <sup>E</sup>	<1.0				
Sediment	Pyefleet	2 <sup>E</sup>	<1.6			350	880
Sediment	Rowhedge	2 <sup>E</sup>	<1.7			310	1000
Sediment	Alresford Creek	2 <sup>E</sup>	<1.4			250	670
Sediment	Brightlingsea Bateman's Tower	2 <sup>E</sup>	<0.25			<120	<180
Sediment	Brightlingsea saltwater pool	2 <sup>E</sup>	<0.27			<130	<140
Sediment	St Osyth boat lake	2 <sup>E</sup>	<1.3			260	870
Seawater	Bradwell Pipeline	2 <sup>E</sup>	<0.28			<3.9	12
Seawater	Bradwell Marina	2 <sup>E</sup>	<0.29			<4.0	13
Seawater	Steeple	2 <sup>E</sup>	<0.29			<4.2	18
Seawater	Maylandsea Bay	2 <sup>E</sup>	<0.29			<4.2	14
Seawater	Blackwater	2 <sup>E</sup>	<0.28			<4.3	14
Seawater	Osea Causeway	2 <sup>E</sup>	<0.30			<4.1	16
Seawater	Maldon	2 <sup>E</sup>	<0.31			<1.5	6.0
Seawater	Heybridge	2 <sup>E</sup>	<0.28			<3.6	15
Seawater	Strood Channel	2 <sup>E</sup>	<0.30			<3.9	16
Seawater	Tollesbury Boatyard	2 <sup>E</sup>	<0.28			<3.8	15
Seawater	Tollesbury saltwater pool	2 <sup>E</sup>	<0.31			<4.8	15
Seawater	Pyefleet	2 <sup>E</sup>	<0.28			<3.6	14
Seawater	Rowhedge	2 <sup>E</sup>	<0.27			<4.6	11
Seawater	Alresford Creek	2 <sup>E</sup>	<0.29			<4.1	13
Seawater	Brightlingsea Bateman's Tower	2 <sup>E</sup>	<0.31			<3.2	11
Seawater	Brightlingsea saltwater pool	2 <sup>E</sup>	<0.28			<4.1	14
Seawater	St Osyth boat lake	2 <sup>E</sup>	<0.28			<3.7	13

**Table 4.3(a). continued**

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>137</sup> Cs	Gross alpha	Gross beta
<b>Terrestrial samples</b>								
Milk		2	<2.8	19				
Milk	max							
Cabbage		1	<3.5	14				
Lucerne		1	<6.8	21				
Grass	Maldon	2 <sup>E</sup>	<9.2	<11		<0.99	<3.9	290
Grass	Strood Channel	2 <sup>E</sup>	<8.5	<4.6		<1.0	<3.0	210
Grass	Tollesbury	2 <sup>E</sup>	<10	<6.4		<1.2	<5.3	190
Grass	West Mersea, near Sewage Works	2 <sup>E</sup>	<9.7	<8.3		<1.2	<2.8	180
Grass	St Osyth	2 <sup>E</sup>	<10	<7.0		<0.91	<2.9	170
Freshwater	Public supply, N side Estuary	1 <sup>E</sup>	<4.1		<0.33	<0.32	<0.060	0.28
Freshwater	Public supply, S side Estuary	1 <sup>E</sup>	<3.3		<0.28	<0.30	<0.073	0.43
Freshwater	Coastal ditch 1	1 <sup>E</sup>	<3.1		<2.2	<0.30	<0.65	3.0
Freshwater	Coastal ditch 2	1 <sup>E</sup>	3.3		<3.2	<0.25	<0.64	2.7
Freshwater	Coastal ditch 3	1 <sup>E</sup>	3.5		<2.4	<0.26	<0.58	7.7
Freshwater	Coastal ditch 4	1 <sup>E</sup>	4.9		<0.32	<0.20	<0.53	6.9

\* Not detected by the method used

<sup>a</sup> Except for milk and water where units are Bq l<sup>-1</sup>, and for sediment where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima.

If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 4.3(b). Monitoring of radiation dose rates near Bradwell, 2015**

Location	Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Bradwell Beach	Sand and pebbles	2	0.068
Beach opposite power station, N side of estuary	Mud and salt marsh	2	0.069
Bradwell Marina	Mud	2	0.078
Waterside	Mud and shingle	1	0.058
Waterside	Mud and pebbles	1	0.060
Steeple	Mud	1	0.085
Steeple	Mud and pebbles	1	0.098
Maylandsea Bay	Mud	1	0.066
Maylandsea Bay	Mud and shingle	1	0.055
Blackwater	Mud and shingle	1	0.061
Blackwater	Mud and pebbles	1	0.068
Osea Causeway	Mud and sand	1	0.059
Osea Causeway	Mud and pebbles	1	0.073
Maldon	Mud	2	0.069
Maldon	Mud and salt marsh	2	0.067
Maldon (2)	Grass	2	0.070
Heybridge	Mud and shingle	1	0.059
Heybridge	Mud and pebbles	1	0.064
Strood Channel	Mud and salt marsh	2	0.065
Strood Channel	Grass	2	0.077
Tollesbury	Grass	2	0.085
Tollesbury boatyard	Mud and salt marsh	1	0.073
Tollesbury boatyard	Mud and pebbles	1	0.082
Tollesbury saltwater pool	Mud and sand	1	0.053
Tollesbury saltwater pool	Mud and pebbles	1	0.068
West Mersea beach huts	Mud	1	0.060
West Mersea beach huts	Mud and sand	1	0.066
West Mersea boatyard	Mud and shells	2	0.058
West Mersea near Sewage Works	Grass	2	0.073
Pyefleet	Mud and pebbles	1	0.072
Pyefleet	Shingle	1	0.067
Rowhedge	Mud and salt marsh	2	0.069
Alresford Creek	Mud and salt marsh	2	0.063
Brightlingsea Bateman's Tower	Mud and shingle	1	0.050
Brightlingsea Bateman's Tower	Pebbles and sand	1	0.058
Brightlingsea saltwater pool	Pebbles and sand	2	0.050
St Osyth	Grass	2	0.069
St Osyth boat lake	Mud	1	0.061
St Osyth boat lake	Mud and pebbles	1	0.070

**Table 4.4(a). Concentrations of radionuclides in food and the environment near Dungeness nuclear power stations, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			Organic <sup>3</sup> H	<sup>3</sup> H	<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>99</sup> Tc
<b>Marine samples</b>								
Cod	Pipeline	1	<25	<25		<0.06		0.18
Sole	Pipeline	1	<25	<25		<0.04		0.07
Shrimps	Pipeline	1	<25	<25		<0.08		<0.07
Scallops	Pipeline	1	<25	<25	32	<0.06	<0.026	<0.05
Sea kale	Dungeness Beach	1				<0.08		<0.08
Seaweed	Folkestone	2 <sup>E</sup>				<0.65	<0.45	<0.47
Sediment	Rye Harbour 1	2 <sup>E</sup>				<0.50		<1.0
Sediment	Camber Sands	2 <sup>E</sup>				<0.22		<0.18
Sediment	Pilot Sands	2 <sup>E</sup>				<0.26		<0.19
Seawater	Dungeness South	2 <sup>E</sup>		<3.0		<0.22		<0.20

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm + <sup>244</sup> Cm	Gross alpha
<b>Marine samples</b>								
Cod	Pipeline	1			<0.17			
Sole	Pipeline	1			<0.11			
Shrimps	Pipeline	1			<0.20			
Scallops	Pipeline	1	0.00039	0.0022	0.00094	*	0.000032	
Sea kale	Dungeness Beach	1			<0.06			
Seaweed	Folkestone	2 <sup>E</sup>			<0.46			
Sediment	Rye Harbour 1	2 <sup>E</sup>	<0.27	0.33	<0.78			610
Sediment	Camber Sands	2 <sup>E</sup>			<0.40			
Sediment	Pilot Sands	2 <sup>E</sup>			<0.34			
Seawater	Dungeness South	2 <sup>E</sup>			<0.27		<3.3	14

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>60</sup> Co	<sup>137</sup> Cs	Gross alpha
<b>Terrestrial Samples</b>								
Milk		2	<3.2	21	<0.16	<0.06	<0.07	
Milk	max				<0.23	<0.07		
Potatoes		1	<3.2	20	0.40	<0.04	<0.08	
Wheat		1	<5.8	83	0.50	<0.08	<0.07	
Freshwater	Long Pits	2 <sup>E</sup>	<3.2		<0.23	<0.20	<0.19	<0.023 0.071
Freshwater	Pumping station Well number 1	1 <sup>E</sup>	<3.2		<0.26	<0.33	<0.27	<0.011 0.098
Freshwater	Pumping station Well number 2	1 <sup>E</sup>	<3.0		<0.23	<0.32	<0.25	<0.014 0.11
Freshwater	Reservoir	1 <sup>E</sup>	<3.0		<0.23	<0.21	<0.18	<0.019 0.073

\* Not detected by the method used

<sup>a</sup> Except for milk and water where units are Bq l<sup>-1</sup>, and for wheat and sediment where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima.

If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 4.4(b). Monitoring of radiation dose rates near Dungeness nuclear power stations, 2015**

Location	Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Littlestone-on-Sea	Sand and shingle	2	0.049
Greatstone-on-Sea	Mud and sand	2	0.054
Dungeness East	Sand	2	0.054
Dungeness South	Shingle	2	0.046
Jurys Gap	Mud and sand	1	0.045
Jurys Gap	Sand and shingle	1	0.065
Rye Bay	Mud and sand	1	0.056
Rye Bay	Sand and shingle	1	0.054

**Table 4.5(a). Concentrations of radionuclides in food and the environment near Hartlepool nuclear power station, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>							
			Organic		<sup>14</sup> C	<sup>60</sup> Co	<sup>99</sup> Tc	<sup>131</sup> I	<sup>137</sup> Cs	<sup>210</sup> Pb
<b>Marine samples</b>										
Cod	Pipeline	1	<25	<25	29	<0.05		<0.94	0.32	
Crabs	Pipeline	1	<25	<25	37	<0.04		<1.1	0.06	
Winkles	South Gare	2	<26	<25	32	<0.04		<0.33	0.09	1.2
Seaweed	Pilot Station	2 <sup>E</sup>				<0.49	3.1	4.7	<0.37	
Sediment	Old Town Basin	2 <sup>E</sup>				<0.39			<2.0	
Sediment	Seaton Carew	2 <sup>E</sup>				<0.27			<0.21	
Sediment	Paddy's Hole	2 <sup>E</sup>				<0.33			1.4	
Sediment	North Gare	2 <sup>E</sup>				<0.36			<0.26	
Sediment	Greatham Creek	2 <sup>E</sup>				<0.64			2.6	
Sea coal	Old Town Basin	2 <sup>E</sup>				<0.39			<0.53	
Sea coal	Carr House Sands	2 <sup>E</sup>				<0.47			<0.84	
Seawater	North Gare	2 <sup>E</sup>		<3.6		<0.25			<0.20	

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>							
			<sup>210</sup> Po	<sup>238</sup> Pu	<sup>239</sup> Pu+ <sup>240</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm+ <sup>244</sup> Cm	Gross alpha	Gross beta
<b>Marine samples</b>										
Cod	Pipeline	1				<0.06				
Crabs	Pipeline	1				<0.11				
Winkles	South Gare	2	13	0.0026	0.020	0.010	*	*		
Seaweed	Pilot Station	2 <sup>E</sup>				<0.44				
Sediment	Old Town Basin	2 <sup>E</sup>				<0.59				
Sediment	Seaton Carew	2 <sup>E</sup>				<0.36				
Sediment	Paddy's Hole	2 <sup>E</sup>				<0.55				
Sediment	North Gare	2 <sup>E</sup>				<0.36				
Sediment	Greatham Creek	2 <sup>E</sup>				<0.74				
Sea coal	Old Town Basin	2 <sup>E</sup>				<0.52				
Sea coal	Carr House Sands	2 <sup>E</sup>				<0.52				
Seawater	North Gare	2 <sup>E</sup>				<0.28			<4.5	12

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>							
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>60</sup> Co	<sup>137</sup> Cs	Gross alpha	Gross beta	
<b>Terrestrial samples</b>										
Milk		2	<2.8	19	<0.21	<0.06				
Milk	max		<3.0	20	<0.28					
Barley		1	<5.5	91	0.90	<0.06				
Potatoes		1	<3.0	24	<0.10	<0.10				
Freshwater	Public supply	2 <sup>E</sup>	<3.0		<0.24	<0.25	<0.22	<0.22	<0.085	0.18
Freshwater	Borehole, Dalton Piercy	2 <sup>E</sup>	<3.3		<0.25	<0.27	<0.22	<0.22	<0.18	0.15

\* Not detected by the method used

<sup>a</sup> Except for milk and water where units are Bq l<sup>-1</sup>, and for sediment and sea coal where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima.

If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 4.5(b). Monitoring of radiation dose rates near Hartlepool nuclear power station, 2015**

Location	Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Fish Sands	Sand	2	0.072
Old Town Basin	Mud and sand	1	0.075
Old Town Basin	Sand	1	0.082
Carr House	Sand	1	0.074
Carr House	Sand and seacoal	1	0.076
Seaton Carew	Sand	2	0.069
Seaton Sands	Sand	2	0.071
North Gare	Pebbles and sand	2	0.072
Paddy's Hole	Sand and stones	1	0.19
Paddy's Hole	Pebbles and sand	1	0.18
Greatham Creek bird hide	Mud	1	0.086
Greatham Creek bird hide	Mud and salt marsh	1	0.093

**Table 4.6(a). Concentrations of radionuclides in food and the environment near Heysham nuclear power stations, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>							
			Organic		<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>99</sup> Tc	<sup>137</sup> Cs	<sup>155</sup> Eu
			<sup>3</sup> H	<sup>3</sup> H						
<b>Marine samples</b>										
Flounder	Morecambe	2	<28	<35	55	<0.10	<0.025	<0.48	4.3	<0.15
Shrimps	Morecambe	2	35	42	85	<0.06	<0.063	0.52	2.5	<0.14
Winkles <sup>b</sup>	Middleton Sands	2	39	<25	61	<0.09	0.12	6.8	2.6	<0.18
Mussels <sup>c</sup>	Morecambe	2	66	64	85	<0.08	0.21	41	2.2	<0.13
Wildfowl	Morecambe	1				<0.08			0.64	<0.13
Seaweed	Half Moon Bay	2 <sup>E</sup>				<0.72		140	3.1	
Sediment	Half Moon Bay	2 <sup>E</sup>				<0.51			33	
Sediment	Pott's Corner	2 <sup>E</sup>				<0.43			11	
Sediment	Morecambe Central Pier	2 <sup>E</sup>				<0.28			3.0	
Sediment	Red Nab Point	2 <sup>E</sup>				<0.50			33	
Sediment	Sunderland Point	4 <sup>E</sup>				<0.50			63	
Sediment	Conder Green	4 <sup>E</sup>				<0.58			63	
Sediment	Sand Gate Marsh	4 <sup>E</sup>				<0.48			77	
Seawater	Heysham Harbour	2 <sup>E</sup>		31		<0.28			<0.23	

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>							Gross alpha	Gross beta
			<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm + <sup>244</sup> Cm			
<b>Marine samples</b>											
Flounder	Morecambe	2	0.00034	0.0022		0.0046	*	0.000039			
Shrimps	Morecambe	2	0.0039	0.024		0.041	*	*			
Winkles <sup>b</sup>	Middleton Sands	2	0.21	1.3	6.5	2.5	*	0.0041		88	
Mussels <sup>c</sup>	Morecambe	2	0.31	1.8	9.5	3.5	*	*		65	
Wildfowl	Morecambe	1				<0.08					
Seaweed	Half Moon Bay	2 <sup>E</sup>				<1.6					
Sediment	Half Moon Bay	2 <sup>E</sup>				59					
Sediment	Pott's Corner	2 <sup>E</sup>				9.8					
Sediment	Morecambe Central Pier	2 <sup>E</sup>				1.2					
Sediment	Red Nab Point	2 <sup>E</sup>				48					
Sediment	Sunderland Point	4 <sup>E</sup>				96			380	780	
Sediment	Conder Green	4 <sup>E</sup>				76			370	870	
Sediment	Sand Gate Marsh	4 <sup>E</sup>				70			240	650	
Seawater	Heysham Harbour	2 <sup>E</sup>				<0.29			<2.9	9.4	

Material	Location or selection <sup>d</sup>	No. of sampling observations <sup>e</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					Gross alpha	Gross beta
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>60</sup> Co	<sup>137</sup> Cs		
<b>Terrestrial samples</b>									
Milk		2	<3.2	18	<0.21	<0.05	<0.12		
Milk	max		<3.3	19	<0.28		0.17		
Potatoes		1	<3.3	20	0.40	<0.05	<0.05		
Wheat		1	<5.1	78	0.80	<0.09	0.25		
Freshwater	Lancaster	2 <sup>E</sup>	<3.0		<0.54	<0.23	<0.20	<0.023	<0.042

\* Not detected by the method used

<sup>a</sup> Except for milk and water where units are Bq l<sup>-1</sup>, and for sediment where dry concentrations apply

<sup>b</sup> The concentration of <sup>210</sup>Po was 9.7 Bq kg<sup>-1</sup>

<sup>c</sup> The concentration of <sup>210</sup>Po was 38 Bq kg<sup>-1</sup>

<sup>d</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima.

If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>e</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>f</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 4.6(b). Monitoring of radiation dose rates near Heysham nuclear power stations, 2015**

Location	Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Greenodd Salt Marsh	Salt marsh	1	0.082
Greenodd Salt Marsh	Grass	1	0.078
Sand Gate Marsh	Salt marsh	3	0.088
Sand Gate Marsh	Grass	1	0.083
High Foulshaw	Mud	1	0.087
High Foulshaw	Sand and mud	1	0.077
High Foulshaw	Grass and mud	1	0.082
High Foulshaw	Salt marsh	1	0.099
Arnside 1	Mud and sand	2	0.082
Arnside 1	Sand	1	0.094
Arnside 1	Grass and sand	1	0.081
Arnside 2	Salt marsh	4	0.098
Morecambe Central Pier	Sand	2	0.076
Half Moon Bay	Sand	1	0.079
Half Moon Bay	Sand and stones	1	0.087
Red Nab Point	Sand	2	0.086
Middleton Sands	Sand	2	0.078
Sunderland	Mud and sand	1	0.093
Sunderland	Salt marsh	3	0.091
Sunderland Point	Mud and sand	4	0.10
Colloway Marsh	Salt marsh	4	0.12
Lancaster	Grass	4	0.082
Aldcliffe Marsh	Salt marsh	4	0.092
Conder Green	Mud	1	0.095
Conder Green	Salt marsh	3	0.090

**Table 4.7(a). Concentrations of radionuclides in food and the environment near Hinkley Point nuclear power stations, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			Organic <sup>3</sup> H	<sup>3</sup> H	<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>99</sup> Tc
<b>Marine samples</b>								
Cod	Stolford	2	<27	36	28	<0.05		0.28
Shrimps	Stolford	1	50	59	37	<0.05		0.20
Limpets	Stolford	1	26	51	29	<0.10		0.19
Seaweed	Pipeline	2 <sup>E</sup>				<0.73	4.6	<0.56
Sediment	Watchet Harbour	2 <sup>E</sup>				<0.57	<2.0	2.3
Sediment	Pipeline	2 <sup>E</sup>				<0.36	<2.0	3.0
Sediment	Stolford	2 <sup>E</sup>				<0.53	<2.0	15
Sediment	Stearl Flats	2 <sup>E</sup>				<0.42	<2.0	5.5
Sediment	River Parrett	2 <sup>E</sup>				<1.2	<2.0	20
Sediment	Weston-Super-Mare	2 <sup>E</sup>				<0.37	<2.0	1.2
Sediment	Burnham-On-Sea	2 <sup>E</sup>				<0.34	<2.5	1.0
Sediment	Kilve	2 <sup>E</sup>				<0.32	<2.0	1.1
Sediment	Blue Anchor Bay	2 <sup>E</sup>				<0.41	<2.0	2.1
Seawater	Pipeline	2 <sup>E</sup>				<0.31	<0.040	<0.24

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>238</sup> Pu	<sup>239</sup> Pu+ <sup>240</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm+ <sup>244</sup> Cm	Gross alpha
<b>Marine samples</b>								
Cod	Stolford	2			<0.15			
Shrimps	Stolford	1	0.00015	0.00071	0.00060	*	*	
Limpets	Stolford	1			<0.08			
Seaweed	Pipeline	2 <sup>E</sup>			<0.72			
Sediment	Watchet Harbour	2 <sup>E</sup>			<0.59			
Sediment	Pipeline	2 <sup>E</sup>			<0.50			
Sediment	Stolford	2 <sup>E</sup>			<0.72			
Sediment	Stearl Flats	2 <sup>E</sup>			<0.60			
Sediment	River Parrett	2 <sup>E</sup>			<1.2			
Sediment	Weston-Super-Mare	2 <sup>E</sup>			<0.42			
Sediment	Burnham-On-Sea	2 <sup>E</sup>			<0.39			
Sediment	Kilve	2 <sup>E</sup>			<0.45			
Sediment	Blue Anchor Bay	2 <sup>E</sup>			<0.55			
Seawater	Pipeline	2 <sup>E</sup>			<0.27		<3.1	13

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>60</sup> Co	<sup>137</sup> Cs	Gross alpha
<b>Terrestrial samples</b>								
Milk		2	<2.8	19	<0.30	<0.05	<0.05	
Milk	max			20	<0.33	<0.06	<0.06	
Honey		1	<4.2	68	0.30	<0.13	<0.14	
Potatoes		1	<3.2	21	0.30	<0.06	<0.09	
Wheat		1	<6.1	85	1.0	<0.05	<0.04	
Freshwater	Durleigh Reservoir	2 <sup>E</sup>	<2.9		<0.52	<0.33	<0.27	<0.030 0.18
Freshwater	Ashford Reservoir	2 <sup>E</sup>	<3.2		<0.45	<0.20	<0.18	<0.039 0.11

\* Not detected by the method used

<sup>a</sup> Except for milk and water where units are Bq l<sup>-1</sup> and for sediment and soil where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 4.7(b). Monitoring of radiation dose rates near Hinkley Point nuclear power stations, 2015**

Location	Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Weston-Super-Mare	Sand and mud	4	0.067
Burnham	Sand and mud	3	0.062
Burnham	Sand	1	0.066
River Parrett	Mud	1	0.087
River Parrett	Grass and mud	1	0.076
River Parrett	Mud and stones	2	0.072
Stear Flats	Mud	2	0.079
Stear Flats	Sand	1	0.083
Stear Flats	Mud and pebbles	1	0.072
Stolford	Mud	2	0.093
Stolford	Mud and rock	2	0.088
Hinkley Point	Mud and rock	2	0.091
Hinkley Point	Sand and shingle	1	0.088
Hinkley Point	Rock and sand	1	0.094
Kilve	Sand and mud	1	0.090
Kilve	Mud and rock	2	0.088
Kilve	Rock and sand	1	0.094
Watchet Harbour	Rock and mud	3	0.10
Watchet Harbour	Rock and sand	1	0.11
Blue Anchor Bay	Sand and mud	4	0.081

**Table 4.8(a). Concentrations of radionuclides in food and the environment near Sizewell nuclear power stations, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			Organic <sup>3</sup> H	<sup>3</sup> H	<sup>14</sup> C	<sup>137</sup> Cs	<sup>238</sup> Pu
<b>Marine samples</b>							
Cod	Sizewell	1	<25	<25		0.17	
Sole	Sizewell	1	<25	<25		0.10	
Crabs	Sizewell	1	<25	<25		<0.07	
Mussels	River Alde	1	<25	<25	31	<0.04	0.00089 0.0053
Sediment	Rifle range	2 <sup>E</sup>				<0.19	
Sediment	Aldeburgh	2 <sup>E</sup>				<0.23	
Sediment	Southwold	2 <sup>E</sup>				6.7	
Seawater	Sizewell	2 <sup>E</sup>		<3.2		<0.25	

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm + <sup>244</sup> Cm	Gross alpha	Gross beta
<b>Marine samples</b>							
Cod	Sizewell	1	<0.05				
Sole	Sizewell	1	<0.17				
Crabs	Sizewell	1	<0.07				
Mussels	River Alde	1	0.0046	*	0.00014		
Sediment	Rifle range	2 <sup>E</sup>	<0.26				
Sediment	Aldeburgh	2 <sup>E</sup>	<0.30				
Sediment	Southwold	2 <sup>E</sup>	<0.70				870
Seawater	Sizewell	2 <sup>E</sup>	<0.27			<3.2	13

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>137</sup> Cs	Gross alpha	Gross beta
<b>Terrestrial samples</b>								
Milk		2	<2.6	<18	<0.20	<0.07		
Milk	max		<2.7	20	<0.23			
Barley		1	<5.0	91	0.60	<0.05		
Potatoes		1	<3.4	17	<0.10	<0.03		
Freshwater	Nature Reserve	2 <sup>E</sup>	<3.0		<0.26	<0.20	<0.038	0.25
Freshwater	The Meare	2 <sup>E</sup>	<3.1		<0.27	<0.26	<0.031	0.21
Freshwater	Leisure Park	2 <sup>E</sup>	11		<0.26	<0.19	<0.052	0.32

\* Not detected by the method used

<sup>a</sup> Except for milk and water where units are Bq l<sup>-1</sup>, and for sediment where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 4.8(b). Monitoring of radiation dose rates near Sizewell, 2015**

Location	Ground type	No. of sampling observations	µGy h <sup>-1</sup>
<b>Mean gamma dose rates at 1m over substrate</b>			
Sizewell Beach	Pebbles and sand	1	0.052
Sizewell Beach	Sand and shingle	1	0.040
Dunwich	Sand and shingle	2	0.053
Rifle Range	Sand and shingle	2	0.049
Aldeburgh	Sand and shingle	1	0.043
Aldeburgh	Shingle	1	0.054
Southwold Harbour	Mud and salt marsh	2	0.073

**Table 4.9(a). Concentrations of radionuclides in food and the environment near Chapelcross nuclear power station, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>								
			<sup>3</sup> H	<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>95</sup> Nb	<sup>95</sup> Zr	<sup>99</sup> Tc	<sup>106</sup> Ru	<sup>110m</sup> Ag
<b>Marine samples</b>											
Flounder	Inner Solway	2		26	<0.10	<0.10	<0.68	<0.46	<0.25	<0.98	<0.15
Salmon	Inner Solway	1	<5.0		<0.10		<0.44	<0.36		<0.91	<0.11
Sea trout	Inner Solway	1	<5.0		<0.10		<0.94	<0.58		<1.1	<0.17
Shrimps	Inner Solway	2	<5.0		<0.10	<0.10	<0.55	<0.36	<0.16	<0.82	<0.14
Cockles	North Solway	1	<5.0		<0.10		<0.46	<0.35		<0.75	<0.10
Mussels	North Solway	2	<5.0	77	<0.14	0.37	<0.31	<0.29	23	<0.82	<0.15
<i>Fucus vesiculosus</i>	Pipeline	4			<0.10		<0.25	<0.20	62	<0.48	<0.11
<i>Fucus vesiculosus</i>	Brownhouses	2			<0.10		<0.42	<0.25	31	<0.41	<0.10
<i>Fucus vesiculosus</i>	Dornoch Brow	2			<0.10		<0.28	<0.18	37	<0.39	<0.10
Sediment	Priestside Bank	1			<0.10		<0.23	<0.21		<0.64	<0.10
Sediment	Pipeline	4	<5.0		0.43		<0.45	<0.25		<1.1	<0.19
Sediment	Dornoch Brow	1			0.28		<0.24	<0.18		<0.53	<0.10
Sediment	Powfoot	1			<0.10		<0.22	<0.22		<0.74	<0.11
Sediment	Redkirk	1			0.15		<0.22	<0.11		<0.68	<0.12
Sediment	Stormont	1			<0.10		<0.14	<0.13		0.34	0.21
Seawater	Pipeline	2	2.7		<0.10		<0.13	<0.11		<0.26	<0.10

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>									
			<sup>125</sup> Sb	<sup>137</sup> Cs	<sup>154</sup> Eu	<sup>155</sup> Eu	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Pu	<sup>241</sup> Am	Gross alpha	Gross beta
<b>Marine samples</b>												
Flounder	Inner Solway	2	<0.28	<4.6	<0.13	<0.24	<0.0081	<0.0072		0.010		
Salmon	Inner Solway	1	<0.25	0.15	<0.12	<0.21				<0.12		
Sea trout	Inner Solway	1	<0.27	1.9	<0.14	<0.24				<0.14		
Shrimps	Inner Solway	2	<0.22	3.0	<0.11	<0.19	<0.0019	0.012		0.034		
Cockles	North Solway	1	<0.21	0.36	<0.10	<0.19	0.070	0.42		0.60		
Mussels	North Solway	2	<0.26	1.7	<0.14	<0.22	0.46	2.7	8.7	5.1		
<i>Fucus vesiculosus</i>	Pipeline	4	<0.14	2.6	<0.10	<0.21	0.28	1.3		2.5	4.4	
<i>Fucus vesiculosus</i>	Brownhouses	2	<0.14	6.0	<0.10	<0.14	0.52	3.0		5.1	8.5	
<i>Fucus vesiculosus</i>	Dornoch Brow	2	<0.13	2.5	<0.10	0.20	0.41	2.2		3.8	6.7	
Sediment	Priestside Bank	1	<0.22	18	<0.19	0.94	2.1	11		20		
Sediment	Pipeline	4	<0.61	130	<0.42	<0.93	<0.014	0.11		0.23		
Sediment	Dornoch Brow	1	0.38	78	0.22	1.5	6.7	39		71		
Sediment	Powfoot	1	<0.26	31	<0.18	<0.27	21	15		25		
Sediment	Redkirk	1	<0.25	48	<0.21	<0.39	13	72		40		
Sediment	Stormont	1	<0.14	42	<0.10	0.73	4.5	25		44		
Seawater	Pipeline	2	<0.10	<0.10	<0.10	<0.10				<0.10		

**Table 4.9(a). continued**

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>95</sup> Nb	<sup>106</sup> Ru
<b>Terrestrial samples</b>									
Milk		10	<6.0	<16	<0.50	<0.05	<0.10	<0.25	<0.36
Milk	max		<15	18	<0.51			<0.31	<0.39
Apples		2	<5.0	17	<0.50	<0.05	<0.10	<0.12	<0.23
Apples	max			18				<0.16	<0.28
Beef		1	<5.0	42	<0.50	<0.10	<0.10	<0.37	<0.76
Cabbage		2	<5.0	<15	<0.52	<0.05	<0.11	<0.65	<0.51
Cabbage	max				<0.53	<0.06	0.13		<0.54
Cauliflower		1	<5.0	9.3	<0.50	<0.60	<0.10	<0.41	<0.36
Duck		1	<5.0	63	<1.2	<0.17	0.28	<1.7	<1.7
Eggs		1	<5.0	43	<0.69	<0.05	<0.10	<0.44	<0.42
Goose		1	<5.0	36	<1.2	<0.19	<0.10	<2.0	<2.0
Leeks		1	<5.0	<15	<0.50	<0.05	0.15	<0.12	<0.27
Potatoes		1	<5.0	23	<0.50	<0.05	<0.10	<0.19	<0.36
Rosehips		1	<5.0	30	<0.50	<0.05	0.37	<0.48	<0.40
Turnips		1	<5.0	15	<0.50	<0.05	0.12	<0.15	<0.31
Wood pigeon		1	<5.0	35	<0.82	<0.05	<0.10	<0.53	<0.48
Grass		4	<6.1	28	<0.50	<0.06	<0.32	<0.32	<0.40
Grass	max		7.1	45		<0.10	0.71	<0.64	<0.91
Soil		4	<23	18	<1.9	<0.05	1.8	<0.47	<0.47
Soil	max		36	21	<2.2		2.3	0.65	<0.53
Freshwater	Purdomstone	1		1.8		<0.01		<0.02	<0.08
Freshwater	Winterhope	1		1.6		<0.01		<0.01	<0.03
Freshwater	Black Esk	1		<1.0		<0.01		<0.01	<0.07
Freshwater	Gullielands Burn	1		16		<0.01		<0.01	<0.03

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					Gross alpha	Gross beta
			<sup>110m</sup> Ag	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>155</sup> Eu	<sup>241</sup> Am		
<b>Terrestrial samples</b>									
Milk		10	<0.05	<0.05	<0.05		<0.06		
Apples		2	<0.05	<0.05	<0.05		<0.06		
Apples	max						<0.07		
Beef		1	<0.10	<0.10	0.11		<0.11		
Cabbage		2	<0.06	<0.05	<0.05		<0.09		
Cabbage	max						<0.11		
Cauliflower		1	<0.05	<0.05	<0.05		<0.05		
Duck		1	<0.28	<0.17	2.1		<0.32		
Eggs		1	<0.05	<0.05	<0.05		<0.05		
Goose		1	<0.28	<0.19	0.99		<0.26		
Leeks		1	<0.05	<0.05	<0.05		<0.05		
Potatoes		1	<0.05	<0.05	<0.05		<0.05		
Rosehips		1	<0.05	<0.05	0.07		<0.11		
Turnips		1	<0.05	<0.05	<0.05		<0.05		
Wood pigeon		1	<0.08	<0.05	1.1		<0.07		
Grass		4	<0.07	<0.06	<0.08		<0.08	<1.2 290	
Grass	max		<0.12	<0.10	0.18		<0.13	2.0 380	
Soil		4	<0.08	<0.07	7.1	1.4	<0.44	110 1100	
Soil	max		<0.09	<0.06	<8.8		0.72	1300	
Freshwater	Purdomstone	1	<0.01	<0.01	<0.01		<0.01	<0.010 0.040	
Freshwater	Winterhope	1	<0.01	<0.01	<0.01		<0.01	0.012 0.044	
Freshwater	Black Esk	1	<0.01	<0.01	<0.01		<0.01	<0.010 0.81	
Freshwater	Gullielands Burn	1	<0.01	<0.01	<0.01		<0.01	<0.010 0.19	

<sup>a</sup> Except for milk and water where units are Bq l<sup>-1</sup>, and for sediment and soil where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima.

If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

**Table 4.9(b). Monitoring of radiation dose rates near Chapelcross, 2015**

Location	Material or Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Glencaple Harbour	Sand	2	0.078
Priestside Bank	Salt marsh	2	0.066
Powfoot Merse	Sediment	2	0.073
Gullielands	Grass	1	0.081
Seafield	Salt marsh	2	0.080
Woodhead	Grass	1	0.073
East Bretton	Grass	1	0.069
Pipeline	Salt marsh	4	0.083
Dumbretton	Grass	1	0.060
Battlehill	Sediment and rocks	2	0.076
Dornoch Brow	Salt marsh	2	0.077
Dornoch Brow	Sediment	2	0.079
Browhouses	Sediment	2	0.076
Redkirk	Sediment	2	0.070
Stormont	Sand	2	0.071
<b>Mean beta dose rates</b>			$\mu\text{Sv h}^{-1}$
Pipeline 500m east	Sand	1	<1.0
Pipeline 500m west	Sand	1	<1.0
Pipeline	Stake nets	3	<1.0

**Table 4.9(c). Radioactivity in air near Chapelcross, 2015**

Location	No. of sampling observations	Mean radioactivity concentration, $\text{mBq m}^{-3}$		
		$^{137}\text{Cs}$	Gross alpha	Gross beta
Eastriggs	12	<0.010	0.016	<0.20
Kirtlebridge	9	<0.010	0.015	<0.20
Brydekirk	10	<0.010	0.013	<0.20

**Table 4.10(a). Concentrations of radionuclides in food and the environment near Hunterston nuclear power station, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			<sup>3</sup> H	<sup>35</sup> S	<sup>54</sup> Mn	<sup>60</sup> Co	<sup>65</sup> Zn	<sup>99</sup> Tc	<sup>110m</sup> Ag
<b>Marine samples</b>									
Cod	Millport	2			<0.12	<0.10	<0.30		<0.14
Hake	Millport	2			<0.10	<0.10	<0.20		<0.60
Crabs	Millport	2			<0.11	<0.10	<0.23	0.60	<0.11
Nephrops	Millport	2			<0.12	<0.10	<0.31		<0.13
Lobsters	Largs	1			<0.10	<0.10	<0.24	21	<0.10
Squat lobsters	Largs	2			<0.10	<0.10	<0.19	7.7	<0.10
Winkles	Pipeline	2			<0.12	<0.10	<0.33		<0.83
Scallops	Largs	2			<0.13	<0.10	<0.33		<0.13
Oysters	Hunterston	1			<0.10	<0.10	<0.21		<0.10
<i>Fucus vesiculosus</i>	N of pipeline	2			<0.12	0.14	<0.17		<0.10
<i>Fucus vesiculosus</i>	S of pipeline	2			0.98	<0.54	<0.21		0.12
Sediment	Largs	1			<0.10	<0.10	<0.26		<0.11
Sediment	Millport	1			<0.10	<0.10	<0.14		<0.10
Sediment	Gull's Walk	1			<0.10	<0.10	<0.10		<0.10
Sediment	Ardneil Bay	1			<0.10	<0.10	<0.14		<0.10
Sediment	Fairlie	1			<0.10	<0.10	<0.10		<0.10
Sediment	Pipeline	1			<0.10	<0.10	<0.16		<0.10
Sediment	Ardrossan North Bay	1			<0.10	<0.10	<0.10		<0.10
Sediment	Ardrossan South Bay	1			<0.10	<0.10	<0.10		<0.10
Seawater	Pipeline	2	4.2	<0.81	<0.10	<0.10	<0.10		<0.10

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			<sup>125</sup> Sb	<sup>137</sup> Cs	<sup>154</sup> Eu	<sup>155</sup> Eu	<sup>238</sup> Pu	<sup>239</sup> Pu+ <sup>240</sup> Pu	<sup>241</sup> Am
<b>Marine samples</b>									
Cod	Millport	2	<0.23	0.66	<0.11	<0.20			<0.13
Hake	Millport	2	<0.16	0.94	<0.10	<0.15			<0.12
Crabs	Millport	2	<0.19	<0.11	<0.10	<0.17	0.0057	0.023	0.021
Nephrops	Millport	2	<0.25	<0.28	<0.12	<0.21			<0.13
Lobsters	Largs	1	<0.20	0.14	<0.10	<0.17			<0.10
Squat lobsters	Largs	2	<0.16	<0.28	<0.10	<0.15	<0.020	<0.019	0.33
Winkles	Pipeline	2	<0.28	<0.20	<0.14	<0.22	<0.19	0.087	0.065
Scallops	Largs	2	<0.26	<0.11	<0.13	<0.25	<0.037	0.048	<0.083
Oysters	Hunterston	1	<0.21	<0.10	<0.11	<0.20			<0.11
<i>Fucus vesiculosus</i>	N of pipeline	2	<0.15	0.23	<0.10	<0.15			<0.11
<i>Fucus vesiculosus</i>	S of pipeline	2	<0.18	<0.19	<0.11	<0.18			<3.4
Sediment	Largs	1	<0.19	3.8	<0.14	0.70			0.23
Sediment	Millport	1	<0.13	3.2	<0.11	<0.23			0.28
Sediment	Gull's Walk	1	<0.10	6.0	<0.10	0.35			0.51
Sediment	Ardneil Bay	1	<0.12	1.8	<0.10	<0.17			<0.15
Sediment	Fairlie	1	<0.10	4.6	<0.10	0.17			0.37
Sediment	Pipeline	1	<0.14	2.3	<0.11	<0.22			0.18
Sediment	Ardrossan North Bay	1	<0.10	2.9	<0.10	0.13			0.28
Sediment	Ardrossan South Bay	1	<0.10	2.6	<0.10	0.22			0.89
Seawater	Pipeline	2	<0.10	<0.10	<0.10	<0.10			<0.10

**Table 4.10(a). continued**

Material	Selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>95</sup> Nb
<b>Terrestrial Samples</b>								
Milk		3	<5.0	<16	<0.50	<0.05	<0.10	<0.22
Milk	max							<0.23
Apples		2	<5.0	<21	<0.50	<0.05	<0.14	<0.20
Apples	max			27			0.18	<0.23
Beef		1	<5.0	45	<0.54	<0.05	<0.10	<0.23
Cabbage		1	<5.0	<15	<0.50	<0.05	<0.10	<0.12
Carrots		1	<5.0	<15	<0.50	<0.05	<0.10	<0.21
Celery		1	<5.0	6.9	<0.14	<0.05	0.10	<0.18
Honey		1	<5.0	91	<0.70	<0.06	<0.10	<0.22
Leeks		1	<5.0	<15	<0.50	<0.05	0.13	<0.58
Pheasant		1	<5.0	35	<1.3	<0.06	0.13	<1.2
Pork		1	<5.0	37	<0.50	<0.05	<0.10	<0.28
Potatoes		1	<5.0	23	<0.50	<0.05	0.11	<0.11
Rabbit		1	<5.0	26	<1.3	<0.05	<0.10	<0.85
Rosehips		1	<5.0	29	<0.50	<0.05	0.62	<0.10
Wild blackberries		1	<5.0	26	0.86	<0.05	0.59	<0.33
Grass		3	<5.0	24	<0.59	<0.05	<0.33	<0.38
Grass	max			29	<0.77		0.57	<0.39
Soil		3	<5.0	<17	<1.6	<0.05	1.2	<0.73
Soil	max			21	<1.7		2.3	<0.86
Freshwater	Knockenden Reservoir	1	<1.0			<0.01		<0.03
Freshwater	Loch Ascog	1	<1.0			<0.01		<0.01
Freshwater	Munnoch Reservoir	1	<1.0			<0.01		<0.01
Freshwater	Camphill	1	<1.0			<0.01		<0.01
Freshwater	Outerwards	1	<1.0			<0.01		<0.02

Material	Selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>110m</sup> Ag	<sup>137</sup> Cs	<sup>155</sup> Eu	<sup>241</sup> Am	Gross alpha	Gross beta
<b>Terrestrial Samples</b>								
Milk		3	<0.05	<0.05		<0.07		
Milk	max			<0.06		<0.06		
Apples		2	<0.06	<0.16		<0.06		
Apples	max			0.27		<0.07		
Beef		1	<0.05	0.50		<0.08		
Cabbage		1	<0.05	<0.05		<0.05		
Carrots		1	<0.05	<0.05		<0.07		
Celery		1	<0.05	<0.05		<0.06		
Honey		1	<0.11	2.7		<0.11		
Leeks		1	<0.06	<0.05		<0.06		
Pheasant		1	<0.09	0.29		<0.08		
Pork		1	<0.05	0.06		<0.13		
Potatoes		1	<0.05	0.11		<0.06		
Rabbit		1	<0.07	0.10		<0.06		
Rosehips		1	<0.05	0.06		<0.12		
Wild blackberries		1	<0.05	0.29		<0.05		
Grass		3	<0.07	0.71		<0.32	<0.58	320
Grass	max		<0.08	1.9		0.84	<1.1	580
Soil		3	<0.07	25	0.59	<0.30	74	680
Soil	max			32	0.92	0.35	93	1000
Freshwater	Knockenden Reservoir	1	<0.01	<0.01		<0.01	<0.010	0.051
Freshwater	Loch Ascog	1	<0.01	<0.01		<0.01	<0.010	0.098
Freshwater	Munnoch Reservoir	1	<0.01	<0.01		<0.01	<0.010	0.052
Freshwater	Camphill	1	<0.01	<0.01		<0.01	<0.010	0.030
Freshwater	Outerwards	1	<0.01	<0.01		<0.02	<0.010	0.056

<sup>a</sup> Except for milk, seawater and freshwater where units are Bq l<sup>-1</sup> and for sediment and soil where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

**Table 4.10(b). Monitoring of radiation dose rates near Hunterston nuclear power station, 2015**

Location	Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Meigle Bay	Sediment	2	0.056
Largs Bay	Sediment	2	0.062
Kilchatten Bay	Sediment	2	0.050
Millport	Sand	2	<0.047
Gull's Walk	Sediment	2	0.061
Hunterston	Sediment	2	0.057
0.5 km north of pipeline	Sediment	2	0.052
0.5 km south of pipeline	Sediment	2	0.058
Portencross	Grass	1	0.053
Ardneil Bay	Sediment	2	0.049
Ardrossan North Bay	Sediment	2	<0.050
Ardrossan South Bay	Sediment	2	0.057
Milstonford	Grass	1	0.058
Biglies	Grass	1	0.091
<b>Beta dose rates</b>			$\mu\text{Sv h}^{-1}$
Millport	Sand	1	<1.0
0.5 km north of pipeline	Sand	1	<1.0
0.5 km south of pipeline	Sand	1	<1.0

**Table 4.10(c). Radioactivity in air near Hunterston, 2015**

Location	No. of sampling observations	Mean radioactivity concentration, $\text{mBq m}^{-3}$		
		$^{137}\text{Cs}$	Gross alpha	Gross beta
Fencebay	12	<0.010	0.015	<0.20
West Kilbride	11	<0.011	0.014	<0.20
Low Ballees	11	<0.010	0.015	<0.20

**Table 4.11(a). Concentrations of radionuclides in food and the environment near Torness nuclear power station, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>54</sup> Mn	<sup>60</sup> Co	<sup>65</sup> Zn	<sup>99</sup> Tc	<sup>110m</sup> Ag	<sup>137</sup> Cs
<b>Marine samples</b>								
Cod	White Sands	2	<0.10	<0.10	<0.16		<0.10	0.32
Mackerel	Pipeline	1	<0.12	<0.10	<0.31		<0.12	0.18
Crabs <sup>d</sup>	Torness	1	<0.10	<0.10	<0.12	<0.23	<0.10	<0.10
Lobsters	Torness	1	<0.11	<0.10	<0.28	3.5	<0.12	<0.10
Nephrops	Dunbar	2	<0.11	<0.10	<0.20		<0.11	<0.11
Winkles	Pipeline	2	<0.12	0.65	<0.29		1.3	<0.13
<i>Fucus vesiculosus</i>	Pipeline	2	0.38	0.67	<0.23		<0.18	<0.10
<i>Fucus vesiculosus</i>	Thornton Loch	2	0.13	0.3	<0.21	11	<0.10	<0.10
<i>Fucus vesiculosus</i>	White Sands	2	<0.11	<0.10	<0.27		<0.11	<0.10
<i>Fucus vesiculosus</i>	Pease Bay	2	<0.10	<0.10	<0.19		<0.10	<0.10
<i>Fucus vesiculosus</i>	Coldingham Bay	2	<0.23	<0.21	<0.63		<0.24	<0.20
Sediment	Dunbar	1	<0.11	<0.10	<0.36		<0.14	2.7
Sediment	Barns Ness	1	<0.10	<0.10	<0.21		<0.10	1.9
Sediment	Thornton Loch	1	<0.10	<0.10	<0.16		<0.10	1.0
Sediment	Heckies Hole	1	<0.10	<0.10	<0.21		<0.10	13
Sediment	Belhaven Bay	1	<0.10	<0.10	<0.21		<0.10	1.9
Sediment	Coldingham Bay	1	<0.10	<0.10	<0.15		<0.10	0.88
Sediment	Pease Bay	1	<0.10	<0.10	<0.29		<0.12	1.6
Seawater <sup>e</sup>	Pipeline	2	<0.10	<0.10	<0.13		<0.10	<0.10

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>155</sup> Eu	<sup>238</sup> Pu	<sup>239</sup> Pu+ <sup>240</sup> Pu	<sup>241</sup> Am	Gross alpha	Gross beta
<b>Marine samples</b>								
Cod	White Sands	2	<0.12			<0.10		
Mackerel	Pipeline	1	<0.20			<0.12		
Crabs <sup>d</sup>	Torness	1	<0.11			<0.10		
Lobsters	Torness	1	<0.21			<0.12		
Nephrops	Dunbar	2	<0.16	<0.0044	0.011	0.011		
Winkles	Pipeline	2	<0.20			<0.13	<0.98	92
<i>Fucus vesiculosus</i>	Pipeline	2	<0.17			<0.12		
<i>Fucus vesiculosus</i>	Thornton Loch	2	<0.16			<0.11		
<i>Fucus vesiculosus</i>	White Sands	2	<0.17			<0.11		
<i>Fucus vesiculosus</i>	Pease Bay	2	<0.14			<0.10		
<i>Fucus vesiculosus</i>	Coldingham Bay	2	<0.45			<0.29		
Sediment	Dunbar	1	1.2			<0.33		
Sediment	Barns Ness	1	0.95			<0.26		
Sediment	Thornton Loch	1	<0.26			<0.22		
Sediment	Heckies Hole	1	1.6			1.2		
Sediment	Belhaven Bay	1	0.81			<0.40		
Sediment	Coldingham Bay	1	<0.23			<0.21		
Sediment	Pease Bay	1	0.59			<0.19		
Seawater <sup>e</sup>	Pipeline	2	<0.13			<0.10		

**Table 4.11(a). continued**

Material	Location or Selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>90</sup> Sr	<sup>95</sup> Nb
<b>Terrestrial Samples</b>							
Milk		2	<5.0	<17	<0.50	<0.10	<0.26
Milk	max			<18			<0.29
Cabbage		1	<5.0	<15	<0.50	<0.10	<0.40
Carrots		1	<5.0	<15	<0.50	0.12	<0.33
Eggs		1	<5.0	32	<0.81	<0.10	<0.41
Honey		1	<5.0	<15	<0.67	<0.10	<0.09
Leeks		1	<5.0	<15	<0.50	<0.10	<0.48
Partridge		1	<5.0	27	<0.74	<0.10	<0.41
Pheasant		1	<5.0	30	<1.0	<0.10	<0.37
Potatoes		2	<5.0	22	<0.50	<0.10	<0.16
Potatoes	max						<0.20
Pork		1	<5.0	<15	<0.54	<0.10	<0.89
Rabbit		1	<5.0	23	<0.78	<0.10	<1.3
Rosehips		1	<5.0	34	<0.50	0.30	<0.21
Venison		1	<5.0	24	<0.89	<0.10	<0.39
Wild mushrooms		1	<5.0	18	<0.50	0.15	<0.62
Grass		3	<5.0	20	<0.50	<0.12	<0.64
Grass	max			23		0.13	<0.71
Soil	max	3	<5.0	<16	<1.4	1.0	<0.61
Soil	max			18	<2.0	1.9	<0.91
Freshwater	Hopes Reservoir	1	<1.0				<0.02
Freshwater	Thorter's Reservoir	1	<1.0				<0.02
Freshwater	Whiteadder	1	<1.0				<0.02
Freshwater	Thornton Loch Burn	1	<1.0				<0.02

Material	Location or Selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			<sup>137</sup> Cs	<sup>155</sup> Eu	<sup>241</sup> Am	Gross alpha	Gross beta
<b>Terrestrial Samples</b>							
Milk		2	<0.05		<0.07		
Milk	max				<0.08		
Cabbage		1	<0.05		<0.05		
Carrots		1	<0.06		<0.08		
Eggs		1	<0.05		<0.06		
Honey		1	4.3		<0.07		
Leeks		1	<0.05		<0.05		
Partridge		1	<0.05		<0.14		
Pheasant		1	0.06		<0.08		
Potatoes		2	<0.05		<0.07		
Pork		1	<0.05		<0.11		
Rabbit		1	<0.07		<0.14		
Rosehips		1	<0.05		<0.06		
Venison		1	0.19		<0.05		
Wild mushrooms		1	0.34		<0.05		
Grass		3	<0.05		<0.06	2.2	300
Grass	max					3.2	420
Soil		3	7.8	1.4	<0.21	140	950
Soil	max		9.8	1.8	<0.22	160	1100
Freshwater	Hopes Reservoir	1	<0.01		<0.01	<0.010	0.036
Freshwater	Thorter's Reservoir	1	<0.01		<0.01	0.010	0.041
Freshwater	Whiteadder	1	<0.01		<0.01	<0.010	0.033
Freshwater	Thornton Loch Burn	1	<0.01		<0.01	0.016	0.10

<sup>a</sup> Except for milk and seawater where units are Bq l<sup>-1</sup> and for sediment and soil where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>d</sup> The concentration of <sup>14</sup>C was 41 Bq kg<sup>-1</sup>

<sup>e</sup> The concentrations of <sup>3</sup>H and <sup>35</sup>S were <1.8 and <0.53 Bq l<sup>-1</sup> respectively

**Table 4.11(b). Monitoring of radiation dose rates near Torness nuclear power station, 2015**

Location	Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Heckies Hole	Mud	2	0.054
Dunbar Inner Harbour	Rocks	2	0.082
Belhaven Bay	Salt marsh	2	0.052
Barns Ness	Rocks	2	0.051
Skateraw	Sand	2	<0.048
Thornton Loch	Grass and soil	1	0.068
Thornton Loch beach	Sand	2	0.057
Ferneylea	Grass and soil	1	0.064
Pease Bay	Sand	2	0.066
St Abbs Head	Rocks	2	0.10
Coldingham Bay	Sand	2	0.059
West Meikle Pinkerton	Grass and soil	1	0.066
<b>Mean beta dose rates on fishing gear</b>			$\mu\text{Sv h}^{-1}$
Torness	Crab and Lobster Pots	2	<1.0

**Table 4.11(c). Radioactivity in air near Torness, 2015**

Location	No. of sampling observations	Mean radioactivity concentration, $\text{mBq m}^{-3}$		
		$^{137}\text{Cs}$	Gross alpha	Gross beta
Innerwick	11	<0.010	0.018	<0.20
Cockburnspath	12	<0.010	<0.024	<0.21
West Barns	12	<0.010	0.021	<0.21

**Table 4.12(a). Concentrations of radionuclides in food and the environment near Trawsfynydd nuclear power station, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			<sup>3</sup> H	<sup>35</sup> S	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>154</sup> Eu
<b>Freshwater samples</b>									
Brown trout	Trawsfynydd Lake	1			<0.22	0.84	<0.25	27	<0.72
Rainbow trout	Trawsfynydd Lake	1			<0.08		<0.09	5.0	<0.28
Sediment	Lake shore	2 <sup>E</sup>			<0.56	3.3	<0.51	340	
Sediment	Bailey Bridge	2 <sup>E</sup>			<1.1	5.4	<0.94	140	
Sediment	Fish farm	2 <sup>E</sup>			<1.3	<2.0	<1.2	530	
Sediment	Footbridge	2 <sup>E</sup>			<1.1	<2.0	<0.98	220	
Sediment	Cae Adda	2 <sup>E</sup>			<0.61	<2.0	<0.53	200	
Freshwater	Public supply	2 <sup>E</sup>	<3.3	<0.29	<0.23		<0.39	<0.30	
Freshwater	Gwylan Stream	2 <sup>E</sup>	<3.4	<0.27	<0.22		<0.23	<0.20	
Freshwater	Hot Lagoon	2 <sup>E</sup>	<3.3	<0.23	<0.27		<0.28	<0.22	
Freshwater	Afon Prysor	2 <sup>E</sup>	<3.3	<0.75	<0.23		<0.32	<0.27	
Freshwater	Trawsfynydd Lake	2 <sup>E</sup>	<3.3	<0.23	<0.13		<0.13	<0.10	
Freshwater	Afon Tafarn-helyg	2 <sup>E</sup>	<3.4	<0.60	<0.24		<0.30	<0.23	

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm + <sup>244</sup> Cm	Gross alpha	Gross beta
<b>Freshwater samples</b>									
Brown trout	Trawsfynydd Lake	1	<0.000076	0.00016	0.00055	*	*		
Rainbow trout	Trawsfynydd Lake	1			<0.08				
Sediment	Lake shore	2 <sup>E</sup>	<0.39	0.72	1.4				
Sediment	Bailey Bridge	2 <sup>E</sup>	<0.49	0.53	<1.0				
Sediment	Fish farm	2 <sup>E</sup>	2.1	6.3	11				
Sediment	Footbridge	2 <sup>E</sup>	<0.41	<1.2	<1.7				
Sediment	Cae Adda	2 <sup>E</sup>	<0.28	<0.64	<1.8				
Freshwater	Public supply	2 <sup>E</sup>						<0.027	<0.017
Freshwater	Gwylan Stream	2 <sup>E</sup>						<0.025	0.15
Freshwater	Hot Lagoon	2 <sup>E</sup>						<0.031	<0.10
Freshwater	Afon Prysor	2 <sup>E</sup>						<0.026	<0.050
Freshwater	Trawsfynydd Lake	2 <sup>E</sup>						<0.027	<0.066
Freshwater	Afon Tafarn-helyg	2 <sup>E</sup>						<0.026	<0.060

Material	Selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			Total <sup>3</sup> H	<sup>14</sup> C	<sup>90</sup> Sr	<sup>137</sup> Cs	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am
<b>Terrestrial Samples</b>									
Milk		2	<3.1	20	<0.029	<0.06			<0.08
Milk	max		<3.4	21	0.033				<0.10
Blackberries		1	<2.7	18		0.12	0.000020	0.000076	0.00021
Grass		1	<4.7	21		0.59	0.00085	0.0056	0.011

\* Not detected by the method used

<sup>a</sup> Except for milk and water where units are Bq l<sup>-1</sup>, and for sediment where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 4.12(b). Monitoring of radiation dose rates near Trawsfynydd nuclear power station, 2015**

Location	Ground type	No. of sampling observations	μGy h <sup>-1</sup>
<b>Mean gamma dose rates at 1m over substrate</b>			
Footbridge	Grass and stones	1	0.10
Footbridge	Stones	1	0.11
Lake shore	Pebbles and stones	1	0.098
Lake shore	Stones	1	0.10
Bailey Bridge	Stones	1	0.097
Bailey Bridge	Rock and stones	1	0.097
Fish Farm	Stones	1	0.10
Fish Farm	Rock and stones	1	0.098
Cae Adda	Pebbles and stones	1	0.090
Cae Adda	Stones	1	0.089

**Table 4.13(a). Concentrations of radionuclides in food and the environment near Wylfa nuclear power station, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			Organic <sup>3</sup> H	<sup>3</sup> H	<sup>14</sup> C	<sup>99</sup> Tc	<sup>137</sup> Cs	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu
<b>Marine samples</b>									
Plaice	Pipeline	1	<25	<25	36		0.41		
Crabs	Pipeline	1	40	<25	56		0.37		
Lobsters	Pipeline	1	<25	<25	58	16	0.50	0.0053	0.032
Winkles	Cemaes Bay	1	<25	<25	36	4.1	0.19	0.012	0.092
Seaweed	Cemaes Bay	2 <sup>E</sup>				13	<0.36		
Sediment	Cemaes Bay	2 <sup>E</sup>					3.0		
Sediment	Cemlyn Bay West	2 <sup>E</sup>					2.4		
Seawater	Cemaes Bay	2 <sup>E</sup>		<3.3			<0.18		
Seawater	Cemlyn Bay West	2 <sup>E</sup>					<0.22		

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			<sup>241</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm + <sup>244</sup> Cm	Gross alpha	Gross beta
<b>Marine samples</b>								
Plaice	Pipeline	1		<0.11				
Crabs	Pipeline	1		<0.19				
Lobsters	Pipeline	1	0.17	0.13	*	*		120
Winkles	Cemaes Bay	1	0.40	0.13	*	*		79
Seaweed	Cemaes Bay	2 <sup>E</sup>		<0.50				
Sediment	Cemaes Bay	2 <sup>E</sup>		1.3				
Sediment	Cemlyn Bay West	2 <sup>E</sup>		0.95				
Seawater	Cemaes Bay	2 <sup>E</sup>		<0.26			<3.0	9.4
Seawater	Cemlyn Bay West	2 <sup>E</sup>		<0.28			<3.7	14

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					Gross alpha	Gross beta
			<sup>3</sup> H	<sup>14</sup> C	<sup>35</sup> S	<sup>137</sup> Cs			
<b>Terrestrial samples</b>									
Milk		2	<2.6	19	<0.38	<0.06			
Milk	max		<2.9	20	<0.43				
Barley		1	<5.4	110	1.9	<0.07			
Potatoes		1	<3.7	23	0.30	<0.03			
Freshwater	Public supply	1 <sup>E</sup>	<3.2		<0.98	<0.18	<0.023	0.17	

\* Not detected by the method used

<sup>a</sup> Except for milk and water where units are Bq l<sup>-1</sup>, and sediment where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 4.13(b). Monitoring of radiation dose rates near Wylfa nuclear power station, 2015**


Location	Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>			
Cemaes Bay	Sand	2	0.069
Cemlyn Bay West	Rock and shingle	1	0.072
Cemlyn Bay West	Shingle	1	0.075

## 5. Defence establishments

This section considers the results of monitoring by the Environment Agency, FSA and SEPA undertaken routinely near nine defence-related establishments in the UK. In addition, the MoD makes arrangements for monitoring at other defence sites where contamination may occur. The operator at the Atomic Weapons Establishment (AWE) in Berkshire carries out environmental monitoring to determine the effects from low level gaseous discharges at its sites. Monitoring at nuclear submarine berths is also conducted by the MoD (DSTL Radiological Protection Services, 2013).

The medium-term trends in doses, discharges and environmental concentrations at Aldermaston, Devonport, Faslane and Coulport, and Rosyth have been considered in a summary report (Environment Agency, FSA, NIEA and SEPA, 2010b).

### 5.1 Aldermaston, Berkshire



AWE at Aldermaston provides and maintains the fundamental components of the UK's nuclear deterrent (Trident). The site and facilities at Aldermaston remain in Government ownership under a

Government Owned Contractor Operator (GOCO) arrangement. The day-to-day operations and the maintenance of Britain's nuclear stockpile are managed, on behalf of the MoD, by AWE plc. The site is regulated by the Environment Agency to discharge low concentrations of radioactive waste to the environment. The most recent discharge permit was issued in 2012.

A habits survey has been conducted to determine the consumption and occupancy rates by members of the public in the vicinity of the site (Ly *et al.*, 2012). Data for consumption, handling and occupancy rates are provided in Appendix 1 (Table X2.2).

#### Doses to the public

In 2015, the *total dose* from all pathways and sources of radiation was less than 0.005 mSv (Table 5.1), or less than

#### Key points

- *Total doses* for the representative person were similar to or less than 1 per cent of the dose limit at all those sites assessed except at Barrow where the effects of historical discharges from Sellafield were apparent
- Discharges, environmental concentrations and dose rates in 2015 were broadly similar to those in 2014 at all establishments

#### Barrow, Cumbria

- *Total dose* for the representative person was 5 per cent of the dose limit but dominated by effects from Sellafield

0.5 per cent of the dose limit. The representative person was a 1 year-old child consuming local cows' milk at high-rates.

Source specific assessments for high-rate consumers of locally grown foods, for sewage workers and for anglers, give exposures that were also less than 0.005 mSv in 2015 (Table 5.1). Estimates of activity concentrations in fish have been based on shellfish samples from the aquatic monitoring programme for dose determination, and for anglers the assessment has conservatively included consumption of fish at a low rate of 1 kg per year.

#### Gaseous discharges and terrestrial monitoring

Gaseous radioactive waste is discharged via stacks on the site. Discharges in 2015 were generally similar to those reported in 2014 though there was an increase in 'volatile beta' discharges. This was due to a change in operations on the site and there were no detected environmental effects due to the increase.

Samples of milk, terrestrial foodstuffs, grass and soil were taken from locations close to the site (Figure 3.4). Activity concentrations in milk and foodstuffs (Table 5.2(a)) were generally below the limits of detection, as in 2014. The tritium concentrations in milk, foodstuffs, grass and soil are reported as less than values in 2015. Tritium is considered in the dose assessment and is of very low significance. In soil samples, where comparisons can be drawn at the same location, concentrations of caesium-137 were similar to values in 2014. Levels of uranium isotopes also remained similar to 2014. Natural background or weapon test

fallout would have made a significant contribution to the levels detected.

## Liquid waste discharges and aquatic monitoring

Discharges of radioactive liquid effluent are made under permit to the sewage works at Silchester (Figure 3.4), and to the Aldermaston stream. Discharges of alpha and other beta radionuclides to Silchester in 2015 were similar to those reported in 2014; discharges of tritium to Aldermaston Stream were very low and similar to those in previous years. There are two factors behind the longer-term decline in discharges of tritium from Aldermaston (Figure 5.1): the closure and decommissioning of the original tritium facility (the replacement facility uses sophisticated abatement technology that results in the discharge of significantly less tritium into the environment), and the historical contamination of groundwater. The historical contamination has been reduced in recent years by radioactive decay and dilution by natural processes. Environmental monitoring of the River Thames (Pangbourne and Mapledurham) has continued to assess the effect of historical discharges.

Activity concentrations for freshwater, fish, crayfish and sediment samples, and measurements of dose rates, are given in Tables 5.2(a) and (b). The concentrations of artificial radioactivity detected in the Thames catchment were very low and generally similar to those for 2014. Iodine-131 detected in sediments and sludges is most likely to be due to waste discharges from the therapeutic uses of this radionuclide in medicine. Concentrations of tritium in samples are reported as less than values. Activity concentrations of artificial radionuclides in River Kennet shellfish were at very low levels and similar to those reported in 2014. Analyses of radiocaesium and uranium activity levels in River Kennet sediments were broadly consistent with previous years. Gross alpha and beta activities in freshwater samples were below the investigation levels for drinking water in the European Directive 2013/51. Gamma dose rates were below or close to natural background.

### 5.2 Barrow, Cumbria




At Barrow, BAE Systems Marine Limited (BAESM) builds, tests and commissions new nuclear powered submarines. Permitted gaseous and liquid discharges were reported as nil and less than 1 per cent

of the discharge limit, respectively, in 2015. The FSA's terrestrial monitoring is limited to vegetable and grass sampling and the Environment Agency monitors dose rates and analyses sediment samples from local intertidal areas. The latter is directed primarily at the far-field effects of Sellafield discharges. A habits survey was undertaken in 2012 (Garrod *et al.*, 2013b). The *total dose* from all pathways and sources of radiation was 0.051 mSv (Table 5.1), which was approximately 5 per cent of the dose limit. The representative person was an adult living on a local houseboat. Virtually all of this dose was due to the effects of Sellafield discharges. A similar dose was found in 2014 (0.055 mSv). The decrease observed was due to a decrease in dose rates underlying the houseboat. Source specific assessments for a vegetable consumer and a person living on a local houseboat were also carried out. The doses in 2015 were both less than 6 per cent of the dose limit. As for *total dose*, the Sellafield source dominated the effects at the houseboat. No assessment of seafood consumption was undertaken in 2015 because of the absence of relevant monitoring data. However, the dose from seafood consumption is less important than that from external exposure on a houseboat (EA, FSA, NIEA, NRW and SEPA, 2014).


Dose rates in intertidal areas near Barrow were slightly enhanced above those expected due to natural background (Table 5.3(b)). This enhancement was due to the far-field effects of historical discharges from Sellafield as evidenced by the results of sediment analysis from the local area (Table 5.3(a)). No effects of discharges from Barrow were apparent in the concentrations of radioactivity in vegetables and grass. All but one of such results are reported as less than values (Table 5.3(a)).

### 5.3 Derby, Derbyshire



Rolls-Royce Marine Power Operations Limited (RRMPOL), a subsidiary of Rolls-Royce plc, carries out design, development, testing and manufacture of nuclear-powered submarine fuel at its two adjacent sites in Derby. Small

discharges of liquid effluent are made via the Megaloughton Lane Sewage Treatment Works to the River Derwent and very low concentrations of alpha activity are present in releases to atmosphere. Other wastes are disposed of by transfer to other sites, including the LLWR near Drigg. A habits survey was undertaken at Derby in 2009 (Elliott *et al.*, 2010).



**Figure 5.1.** Trends in liquid discharges of tritium and plutonium-241 from Aldermaston, Berkshire 1990-2015 (including discharges to River Thames at Pangbourne, Silchester sewer and Aldermaston Stream)


## Doses to the public

In 2015, the *total dose* from all pathways and sources of radiation was less than 0.005 mSv (Table 5.1), which is less than 0.5 per cent of the dose limit. Source specific assessments for consumption of vegetables, fish and drinking river water at high-rates, and for a local resident exposed to external and inhalation pathways from gaseous discharges, give exposures that were also less than 0.005 mSv in 2015 (Table 5.1).

Results of the routine monitoring programme at Derby are given in Table 5.3(a). Analysis of uranium activity in samples taken around the site in 2015 found levels broadly consistent with previous years. More detailed analysis in previous years has shown the activity as being consistent with natural sources. Gross alpha and beta activities in water from the River Derwent were less than the investigation levels for drinking water in the European Directive 2013/51, and the dose from using the river as a source of drinking water was much less than 0.005 mSv per year (Table 5.1). Caesium-137 detected in sediments from local water courses is most likely to have been from fallout from overseas sources.

Table 5.3(a) also includes analysis results from a water sample taken from Fritchley Brook, downstream of Hilt's Quarry. RRMPOL formerly used the quarry for the controlled burial of solid low level radioactive waste. Uranium isotopes detected in the sample were similar to those levels observed elsewhere in Derbyshire (Table 8.8).

## 5.4 Devonport, Devon



The Devonport Royal Dockyard consists of two parts and is operated by Her Majesty's Naval Base (MoD) and Devonport Royal Dockyard Limited (owned by Babcock International Group plc). Devonport Royal Dockyard

refits, refuels, repairs and maintains the Royal Navy's nuclear powered submarines and has a permit granted by the Environment Agency to discharge liquid radioactive waste to the Hamoaze, which is part of the Tamar Estuary, and to the local sewer, and gaseous waste to the atmosphere. During June 2011, a habits survey was conducted to determine the consumption and occupancy rates by members of the public (Clyne *et al.*, 2012). The routine monitoring programme in 2015 consisted of measurements of gamma dose rate and analysis of grass, vegetables, fish, shellfish and other marine indicator materials (Tables 5.3(a) and (b)).

### Doses to the public

In 2015, the *total dose* from all pathways and sources of radiation was less than 0.005 mSv (Table 5.1), which was less than 0.5 per cent of the dose limit. The representative person was an adult seafood consumer. Trends in *total doses* in the area of the south coast (and the Severn Estuary) are shown in Figure 6.1.

Source specific assessments for a high-rate consumer of locally grown food and of fish and shellfish, and for an occupant of a houseboat, give exposures that were also less than 0.005 mSv (Table 5.1) which was less than 0.5 per cent of the dose limit for members of the public of 1 mSv.


### Gaseous discharges and terrestrial monitoring

Gaseous discharges were generally similar to those in 2014. Samples of grass and vegetables were analysed for a number of radionuclides, and concentrations were below the limits of detection in both.

### Liquid waste discharges and marine monitoring

Discharges to the Hamoaze were similar to those reported in 2014. Figure 5.2 shows the discharge history of tritium and cobalt-60 since 1990. The main contributor to the variations in tritium discharges over time has been the re-fitting of Vanguard class submarines. These submarines have a high tritium inventory as they do not routinely discharge primary circuit coolant until they undergo refuelling at Devonport. The underlying reason for the overall decrease in cobalt-60 discharges over this period was the improvement in submarine reactor design so that less cobalt-60 was produced during operation, and therefore less was released during submarine maintenance operations. In marine samples, concentrations of tritium and cobalt-60 were below limits of detection. Trace amounts of caesium-137, likely to originate from Chernobyl and global weapon test fallout, were measured in sediment samples. Carbon-14 concentrations in fish species are similar to those in recent years and are likely to be due to natural sources. The seaweed samples contained very low concentrations of iodine-131 in 2015. These were most likely to have originated from the therapeutic use of this radionuclide in a local hospital. Gamma dose rates in the vicinity of Devonport were similar to 2014. They reflect the local effects of enhanced background radiation from natural sources.

## 5.5 Faslane and Coulport, Argyll and Bute




The HMNB Clyde establishment consists of the naval base at Faslane and the armaments depot at Coulport. Babcock Marine, a subsidiary of Babcock International Group plc, operates HMNB Clyde, Faslane in partnership with the

MoD. However, the MoD remains in control of the undertaking, through the Naval Base Commander, Clyde (NBC Clyde) in relation to radioactive waste disposal. MoD through NBC Clyde also remains in control of the undertaking at Coulport although many of the activities undertaken at Coulport have been outsourced to an industrial alliance comprising of AWE plc, Babcock and Lockheed Martin UK (known as ABL). Discharges of liquid radioactive waste into Gare Loch from Faslane and the discharge of gaseous radioactive waste in the form of tritium to the atmosphere from Coulport are made under letters of agreement between SEPA and the MoD. SEPA reviewed these letters in 2012 and the process of updating the letters continued in 2015. The discharges released during 2015 are shown in Appendix 2. The disposal of solid radioactive waste from each site is also made under letters of agreement between SEPA and the MoD. There were no disposals of solid waste from the sites during 2015.

During August 2011, a habits survey was conducted to determine the consumption and occupancy rates by members of the public (Rumney *et al.*, 2013b). Data for consumption, handling and occupancy rates are summarised in Appendix 1 (Table X2.2).

The *total dose* from all pathways and sources of radiation was less than 0.005 mSv in 2015 (Table 5.1). The representative person was an adult exposed to radioactivity in marine sediments, but as in 2014 the dose was less than 0.5 per cent of the dose limit for members of the public. Source specific assessments for a high-rate consumer of fish and shellfish and a consumer of locally grown food (based on limited data), give exposures that were also less than 0.005 mSv.

The routine marine monitoring programme consisted of the analysis of shellfish, seawater, seaweed and sediment samples, and gamma dose rate measurements. Terrestrial monitoring included beef, honey, water, grass and soil sampling. The results are given in Tables 5.3(a) and (b) and were similar to those in 2014. Radionuclide concentrations were generally below the limits of detection, with caesium-137 concentrations in sediment consistent with



**Figure 5.2.** Trends in liquid discharges of tritium and cobalt-60 from Devonport, Devon 1990-2015

the distant effects of discharges from Sellafield, and with weapon testing and Chernobyl fallout. Tritium, gross alpha and gross beta concentrations in freshwater were much lower than the investigation levels in the European Directive 2013/51. Carbon-14 concentrations in shellfish were similar to natural background levels. Gamma dose rates measured in the surrounding area were difficult to distinguish from natural background.

### 5.6 Holy Loch, Argyll and Bute



A small programme of monitoring at Holy Loch continued during 2015 in order to determine the effects of past discharges from the US submarine support facilities which closed in 1992. Radionuclide concentrations were

low (Table 5.3(a)). Gamma dose rate measurements from intertidal areas (Table 5.3(b)) were similar to those observed in 2014. The external radiation dose to a person spending time on the loch shore was 0.012 mSv in 2015, which was approximately 1 per cent of the dose limit for members of the public of 1 mSv (Table 5.1).

### 5.7 Rosyth, Fife



The site is operated by Babcock Marine, a division of Babcock International Group plc, who are responsible for the management of radioactive waste that was generated when the site supported the nuclear submarine

fleet. Site decommissioning started in April 2006 and has mainly been completed, with the exception of some small areas of the site where facilities continue to be required to manage radioactive wastes. To date, more than 99 per cent of the waste arising as a result of site decommissioning is being recycled.

Radioactive waste produced during decommissioning has been disposed of under an authorisation granted to Rosyth Royal Dockyard Limited (RRDL) in October 2008. Radioactive aqueous and gaseous wastes continue to be discharged in accordance with conditions in the same authorisation.

SEPA has received an application under RSA 93 from RRDL to dispose of solid, liquid and gaseous low level radioactive wastes arising from work to dismantle the seven redundant submarines currently berthed at Rosyth. SEPA has also received an application from the MoD to transfer solid and liquid radioactive waste from the submarines to allow RRDL to carry out the dismantling work. This application will be handled administratively in the form of a Letter of Approval

between SEPA and the MoD rather than under RSA 1993. Currently, SEPA is determining both applications.

SEPA, and other stakeholders, are currently engaging with the MoD Nuclear Legacy Works Team based at Rosyth to identify the optimised arrangements to manage radiologically contaminated ion-exchange resins stored securely in the Active Waste Accumulation Facility on the Rosyth site. This will include trials to assess suitable treatment technologies to manage resin wastes. SEPA is working closely with the ONR and the Environment Agency on resin management as the issue is common to the Rosyth and Devonport naval sites.

The *total dose* from all pathways and sources was 0.006 mSv in 2015 (Table 5.1), which was less than 1 per cent of the dose limit. The representative person was an adult with exposure over marine sediments. The source specific assessments for marine pathways user gave an exposure 0.010 mSv in 2015.

In 2015, authorised gaseous discharges from Rosyth were reported as nil. Liquid wastes are discharged via a dedicated pipeline to the Firth of Forth. In all cases the activities in the liquid discharged were below authorised limits. Discharges of tritium from Rosyth increased in 2015, due to an increase in the numbers of samples of nuclear submarine primary coolant that were disposed of following analysis in the Rosyth Radiochemistry Laboratory.

SEPA's routine monitoring programme included analysis of shellfish, environmental indicator materials and measurements of gamma dose rates in intertidal areas. Results are shown in Tables 5.3(a) and (b). The radioactivity

levels detected were at similar low levels to 2014 and in most part due to the combined effects of Sellafield, weapon testing and Chernobyl. Gamma dose rates were difficult to distinguish from natural background. A new habits survey was undertaken in 2015 and the results were incorporated in the dose assessment given above (SEPA, in press).

## 5.8 Vulcan NRTE, Highland



The Vulcan Naval Reactor Test Establishment is operated by Defence Equipment and Support, part of the Ministry of Defence, and its purpose was to prototype submarine nuclear reactors. It is located adjacent to

the Dounreay site and the impact of its discharges is considered along with those from Dounreay (in Section 3). The site ceased critical reactor operations in July 2015 and will not be required for further prototyping. Since the reactor shutdown for the last time, work has focused on post-operational activity. This will include defueling of the reactor, clearance of fuel from the site and preparations for future decommissioning and disposal of both the reactors from the site and their component parts expected sometime after 2022.

**Table 5.1. Individual doses – defence sites, 2015**

Site	Representative person <sup>a,b</sup>	Exposure mSv, per year					
		Total	Fish and shellfish	Other local food	External radiation from intertidal areas or river banks	Intakes of sediment or water	Gaseous plume related pathways
Aldermaston and Burghfield							
<b>Total dose – all sources</b>	<b>Infant milk consumers</b>	<b>&lt;0.005<sup>c</sup></b>	-	<b>&lt;0.005</b>	-	-	-
Source specific doses	Anglers	<0.005 <sup>c</sup>	<0.005	-	<0.005	-	-
	Infant inhabitants and consumers of locally grown food	<0.005 <sup>c</sup>	-	<0.005	-	-	<0.005
	Workers at Silchester STW	<0.005	-	-	<0.005 <sup>d</sup>	<0.005 <sup>e</sup>	-
Barrow							
<b>Total dose – all sources</b>	<b>Adult occupants on houseboats<sup>g</sup></b>	<b>0.051</b>	-	-	<b>0.051</b>	-	-
Source specific doses	Houseboat occupants	0.050	-	-	0.050	-	-
	Consumers of locally grown food	<0.005	-	<0.005	-	-	-
Derby							
<b>Total dose – all sources</b>	<b>Adult consumers of locally sourced water</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	-	<b>&lt;0.005</b>	<b>&lt;0.005</b>	-
Source specific doses	Anglers consuming fish and drinking water <sup>f</sup>	<0.005	<0.005	-	<0.005	<0.005	-
	Inhabitants and consumers of locally grown food	<0.005 <sup>c</sup>	-	<0.005	-	-	<0.005
Devonport							
<b>Total dose – all sources</b>	<b>Adult fish consumers</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	-	<b>0.000</b>	-	-
Source specific doses	Seafood consumers	<0.005	<0.005	-	<0.005	-	-
	Houseboat occupants	<0.005	-	-	<0.005	-	-
	Prenatal children of inhabitants and consumers of locally grown food	<0.005	-	<0.005	-	-	<0.005
Faslane							
<b>Total dose – all sources</b>	<b>Adult consumers of cattle meat</b>	<b>&lt;0.005</b>	-	<b>&lt;0.005</b>	-	-	-
Source specific doses	Seafood consumers	<0.005	<0.005	-	<0.005	-	-
	Consumers of locally grown food	<0.005	-	<0.005	-	-	-
Holy Loch							
Source specific doses	Anglers	0.012	-	-	0.012	-	-
Rosyth							
<b>Total dose – all sources</b>	<b>Adult occupants over sediment</b>	<b>0.006</b>	<b>&lt;0.005</b>	-	<b>0.006</b>	-	-
Source specific doses	Fishermen and beach users	0.010	<0.005	-	0.007	-	-

- <sup>a</sup> The total dose is the dose which accounts for all sources including gaseous and liquid discharges and direct radiation. The total dose for the representative person with the highest dose presented. Other dose values are presented for specific sources, either liquid discharges or gaseous discharges, and their associated pathways. They serve as a check on the validity of the total dose assessment. The representative person is an adult unless otherwise specified
- <sup>b</sup> None of the people represented in this table were considered to receive direct radiation from the sites listed
- <sup>c</sup> Includes a component due to natural sources of radionuclides
- <sup>d</sup> External radiation from raw sewage and sludge
- <sup>e</sup> Intakes of resuspended raw sewage and sludge
- <sup>f</sup> Water is from rivers and streams and not tap water
- <sup>g</sup> Exposures at Barrow are largely due to discharges from the Sellafield site

**Table 5.2(a). Concentrations of radionuclides in food and the environment near Aldermaston, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			Organic <sup>3</sup> H	<sup>3</sup> H	<sup>131</sup> I	<sup>137</sup> Cs	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U
<b>Freshwater samples</b>									
Flounder	Woolwich Reach	1		<25	<0.26	<0.06			
Signal crayfish	Ufton Bridge – Theale	1	<25	<25	<0.19	<0.04	0.061	0.0015	0.046
Sediment	Pangbourne	4 <sup>E</sup>			5.7	<2.2	14	<0.96	16
Sediment	Mapledurham	4 <sup>E</sup>			1.9	16	9.2	<0.75	10
Sediment	Aldermaston	4 <sup>E</sup>				<3.3	20	<1.2	18
Sediment	Spring Lane	4 <sup>E</sup>				1.7	10	<0.83	9.9
Sediment	Stream draining south	4 <sup>E</sup>				<0.94	25	<2.0	26
Sediment	Reading (Kennet)	4 <sup>E</sup>				<2.7	14	<1.2	14
Gullypot sediment	Falcon Gate	1 <sup>E</sup>		<9.1	<1.6	21	1.5	20	
Gullypot sediment	Main Gate	1 <sup>E</sup>		<13	<1.1	17	1.5	18	
Gullypot sediment	Tadley Entrance	1 <sup>E</sup>		<7.1		13	18	<1.5	19
Gullypot sediment	Burghfield Gate	1 <sup>E</sup>		<13		4.2	19	<1.9	20
Freshwater	Pangbourne	4 <sup>E</sup>		<3.2	<0.23	0.0094	<0.0015	0.0081	
Freshwater	Mapledurham	4 <sup>E</sup>		<4.9	<0.20	0.0095	<0.0010	0.0071	
Freshwater	Aldermaston	4 <sup>E</sup>		<5.0	<0.20	0.0071	<0.0013	0.0051	
Freshwater	Spring Lane	4 <sup>E</sup>		<3.6	<0.19	<0.0027	<0.0012	<0.0026	
Freshwater	Reading (Kennet)	4 <sup>E</sup>		<3.0	<0.20	0.0066	<0.0018	<0.0048	
Crude liquid effluent	Silchester treatment works	4 <sup>E</sup>		<15	<0.22	<0.0044	<0.0020	<0.0049	
Final liquid effluent	Silchester treatment works	4 <sup>E</sup>		<12	<0.24	<0.0037	<0.0012	<0.0039	
Sewage sludge	Silchester treatment works	4 <sup>E</sup>		<12	16	<0.32	1.6	<0.080	1.6

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm+ <sup>244</sup> Cm	Gross alpha	Gross beta
<b>Freshwater samples</b>									
Flounder	Woolwich Reach	1			<0.16				
Signal crayfish	Ufton Bridge – Theale	1	0.000011	0.000046	0.000055 *		*		
Sediment	Pangbourne	4 <sup>E</sup>	<0.37	<0.28	<0.70			310	400
Sediment	Mapledurham	4 <sup>E</sup>	<0.25	<0.33	<0.45			<110	250
Sediment	Aldermaston	4 <sup>E</sup>	<0.28	4.0	<1.3			240	440
Sediment	Spring Lane	4 <sup>E</sup>	<0.29	<0.21	<0.57			<160	390
Sediment	Stream draining south	4 <sup>E</sup>	<0.31	<0.26	<0.83			390	790
Sediment	Reading (Kennet)	4 <sup>E</sup>	<0.42	<0.26	<0.90			140	380
Gullypot sediment	Falcon Gate	1 <sup>E</sup>	<0.27	<0.14	<1.6			350	760
Gullypot sediment	Main Gate	1 <sup>E</sup>	<0.33	<0.17	<1.3			570	650
Gullypot sediment	Tadley Entrance	1 <sup>E</sup>	<0.34	0.76	<1.4			240	710
Gullypot sediment	Burghfield Gate	1 <sup>E</sup>	<0.18	<0.30	<1.5			230	600
Freshwater	Pangbourne	4 <sup>E</sup>	<0.0034	<0.0022	<0.0059			<0.054	0.26
Freshwater	Mapledurham	4 <sup>E</sup>	<0.0037	<0.0025	<0.0076			<0.061	0.25
Freshwater	Aldermaston	4 <sup>E</sup>	<0.0030	<0.0019	<0.0069			<0.042	0.19
Freshwater	Spring Lane	4 <sup>E</sup>	<0.0034	<0.0018	<0.0056			<0.038	0.20
Freshwater	Reading (Kennet)	4 <sup>E</sup>	<0.0030	<0.0015	<0.0049			<0.049	0.12
Crude liquid effluent	Silchester treatment works	4 <sup>E</sup>	<0.0045	<0.0026	<0.27			<0.084	0.71
Final liquid effluent	Silchester treatment works	4 <sup>E</sup>	<0.0027	<0.0015	<0.27			<0.060	0.60
Sewage sludge	Silchester treatment works	4 <sup>E</sup>	<0.051	<0.038	<0.40			<13	34

**Table 5.2(a). continued**

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			<sup>3</sup> H	<sup>137</sup> Cs	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U
<b>Terrestrial samples</b>							
Milk		2	<2.6	<0.06	0.0015	<0.00073	<0.00092
Milk	max		<3.0	<0.07	0.0018	<0.00093	<0.0013
Beetroot		1	<3.2	<0.05	0.018	0.00088	0.018
Wheat		1	<5.8	<0.11	0.0026	0.00023	0.0024
Grass	Location 3	1 <sup>E</sup>	<9.3	<0.95	0.27	<0.11	0.19
Grass	Location 7	1 <sup>E</sup>	<16	<0.52	1.2	<0.17	1.2
Grass	Opposite Gate 26A	1 <sup>E</sup>	<28	<0.77	<0.41	<0.18	<0.30
Grass	Tatley	1 <sup>E</sup>	<22	<0.92	0.23	<0.14	0.35
Soil	Location 3	1 <sup>E</sup>	<11	12	20	<1.6	22
Soil	Location 7	1 <sup>E</sup>	<9.7	11	23	<1.2	24
Soil	Opposite Gate 26A	1 <sup>E</sup>	<48	12	7.1	<0.75	8.1
Soil	Tatley	1 <sup>E</sup>	<9.2	3.2	15	<1.2	17

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>				
			<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am	Gross alpha	Gross beta
<b>Terrestrial samples</b>							
Milk		2	<0.000018	<0.000017	<0.000040		
Milk	max			<0.000018	<0.000045		
Beetroot		1	0.000041	0.00037	0.00040		
Wheat		1	0.000031	0.000034	0.000061		
Grass	Location 3	1 <sup>E</sup>	<0.063	<0.047		4.1	160
Grass	Location 7	1 <sup>E</sup>	<0.032	0.046		13	110
Grass	Opposite Gate 26A	1 <sup>E</sup>	<0.040	<0.026		10	220
Grass	Tatley	1 <sup>E</sup>	<0.045	<0.037		5.1	220
Soil	Location 3	1 <sup>E</sup>	<0.33	0.53		420	460
Soil	Location 7	1 <sup>E</sup>	<0.34	1.7		200	530
Soil	Opposite Gate 26A	1 <sup>E</sup>	<0.31	0.40		<120	290
Soil	Tatley	1 <sup>E</sup>	<0.47	<0.34		230	400

\* Not detected by the method used

<sup>a</sup> Except for milk, sewage effluent and water where units are Bq l<sup>-1</sup>, and for sediment and soil where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max'. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 5.2(b). Monitoring of radiation dose rates near Aldermaston, 2015**

Location	Ground type type	No. of sampling observations	µGy h <sup>-1</sup>
<b>Mean gamma dose rates at 1m over substrate</b>			
Pangbourne, riverbank	Grass and mud	1	0.064
Pangbourne, riverbank	Grass	3	0.070
Mapledurham, riverbank	Grass and mud	1	0.075
Mapledurham, riverbank	Grass	2	0.062
Mapledurham, riverbank	Grass and stones	1	0.068

**Table 5.3(a). Concentrations of radionuclides in food and the environment near defence establishments, 2015**

Material	Location or selection <sup>a</sup>	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>							
			Organic <sup>3</sup> H	<sup>3</sup> H	<sup>14</sup> C	<sup>60</sup> Co	<sup>125</sup> Sb	<sup>131</sup> I	<sup>134</sup> Cs	<sup>137</sup> Cs
<b>Barrow</b>										
Grass	Barrow	1 <sup>F</sup>		<3.3		<0.21		<0.23	0.54	
Potatoes	Barrow	1 <sup>F</sup>		<3.0		<0.08		<0.06	<0.09	
Sediment	Walney Channel	2				<0.41	<1.4	<0.37	70	
	- N of discharge point									
Sediment	Walney Channel	2				<0.71	<1.4	<0.37	88	
	- S of discharge point									
Sediment	Walney Channel	1				<1.5	<5.3	<1.3	99	
	- S of discharge point (2)									
<b>Derby</b>										
Potatoes	Derby	1 <sup>F</sup>				<0.10		<0.07	0.12	
Wheat	Derby	1 <sup>F</sup>				<0.06		<0.06	<0.02	
Sediment	River Derwent, upstream	1				<2.0			4.1	
Sediment	Fritchley Brook	1				<0.42			0.64	
Sediment	River Derwent, downstream	4				<2.2			4.8	
Water	River Derwent, upstream	1				<0.23				
Water <sup>c</sup>	Fritchley Brook	1		<3.2		<0.30			<0.24	
Water	River Derwent, downstream	4				<0.25				
<b>Devonport</b>										
Ballan wrasse	Plymouth Sound	1 <sup>F</sup>	<25	<25	43	<0.04		<0.28	<0.03	0.15
Spiny spider crab	Plymouth Sound	1 <sup>F</sup>			22	<0.04		<0.14	<0.04	<0.03
Shrimp	River Lynher	1 <sup>F</sup>			33	<0.20		*	<0.19	<0.16
Mussels	River Lynher	1 <sup>F</sup>	<25	<25	31	<0.11		<0.58	<0.11	<0.10
Seaweed <sup>d</sup>	Kinterbury	2				<0.73		3.5		
Sediment <sup>e</sup>	Kinterbury	2		<22		<0.63				2.2
Sediment	Torpoint South	2		<43		<1.1				
Sediment	Lopwell	2		<42		<1.8				5.8
Seawater	Torpoint South	2		<3.2	<11	<0.23				
Seawater	Millbrook Lake	2		<3.5	<8.8	<0.28				
Beetroot		1 <sup>F</sup>		<3.5		<0.07		<0.11	<0.09	
Grass		1 <sup>F</sup>		<4.5		<0.12		<0.07	<0.13	
<b>Faslane</b>										
Mussels	Rhu	1			28	<0.10		<0.10	0.19	
Winkles	Rhu	1			35	<0.10		<0.10	0.22	
<i>Fucus vesiculosus</i>	Rhu	1				<0.10		<0.10	0.14	
<i>Fucus vesiculosus</i>	Garelochhead	1				<0.10		<0.10	0.20	
<i>Fucus vesiculosus</i>	Carnban boatyard	1				<0.10		<0.10	0.20	
Sediment	Rhu	1				<0.10		<0.10	5.5	
Sediment	Garelochhead	1				<0.10		<0.10	3.4	
Sediment	Carnban boatyard	1				<0.10		<0.10	2.8	
Seawater	Carnban boatyard	2		<1.0		<0.10		<0.10	<0.10	
Beef muscle	Faslane	1				<0.05		<0.05	0.57	
Honey	Faslane	1				<0.05		<0.05	0.80	
Grass	Auchengaich Reservoir	1		<5.0		<0.05		<0.05	0.24	
Grass	Lochan Ghlas Laoigh	1		<5.0		<0.05		<0.05	7.5	
Soil	Auchengaich	1				<0.05		<0.06	20	
Soil	Lochan Ghlas Laoigh	1				<0.08		<0.11	30	
Freshwater	Helensburgh Reservoir	1		<1.0		<0.01		<0.01	<0.01	
Freshwater	Loch Finlas	1		<1.0					<0.01	
Freshwater	Auchengaich Reservoir	1		<1.0		<0.01		<0.01	<0.01	
Freshwater	Lochan Ghlas Laoigh	1		<1.0		<0.01		<0.01	<0.01	
Freshwater	Loch Eck	1		<1.0		<0.01		<0.01	<0.01	
Freshwater	Loch Lomond	1		<1.0		<0.01		<0.01	<0.01	
<b>Holy Loch</b>										
Sediment	Mid-Loch	1				<0.10		<0.10	4.2	
<b>Rosyth</b>										
Winkles	St David's Bay	1				<0.10		<0.11	0.51	
<i>Fucus vesiculosus</i>	East of dockyard	1				<0.10		<0.10	<0.10	
Sediment	East of dockyard	1				<0.10		<0.10	2.2	
Sediment	Port Edgar	1				<0.10		<0.10	12	
Sediment	West of dockyard	1				<0.10		<0.10	2.8	
Sediment	East Ness Pier	1				<0.10		<0.10	5.8	
Sediment	Blackness Castle	1				<0.10		<0.10	0.82	
Sediment	Charlestown Pier	1				<0.10		<0.10	0.56	
Seawater	East of dockyard	2		<1.0		<0.10		<0.10	<0.10	
Freshwater	Castlehill Reservoir	1		<1.0		<0.01		<0.01	<0.01	
Freshwater	Holl Reservoir	1		<1.0		<0.01		<0.01	<0.01	
Freshwater	Gartmorn Dam	1		<1.0		<0.01		<0.01	<0.01	
Freshwater	Morton No. 2 Reservoir	1		<1.0		<0.01		<0.01	<0.01	

**Table 5.3(a). continued**

Material	Location or selection <sup>a</sup>	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>							
			<sup>154</sup> Eu	<sup>155</sup> Eu	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U	<sup>241</sup> Am	Gross alpha	Gross beta
<b>Barrow</b>										
Grass	Barrow	1 <sup>F</sup>	<0.84	<0.48				<0.34		
Potatoes	Barrow	1 <sup>F</sup>	<0.27	<0.20				<0.12		
Sediment	Walney Channel	2	<0.93	<0.79				160	680	660
	- N of discharge point									
Sediment	Walney Channel	2	<0.88	<0.80				210	470	770
	- S of discharge point									
Sediment	Walney Channel	1	<3.6	<1.8				190	510	820
	- S of discharge point (2)									
<b>Derby</b>										
Potatoes	Derby	1 <sup>F</sup>	<0.28	<0.14	0.0099	0.0011	0.011	<0.14		
Wheat	Derby	1 <sup>F</sup>	<0.17	<0.10	<0.0013	<0.00048	<0.00048	<0.09		
Sediment	River Derwent, upstream	1			36	<1.7	34		440	660
Sediment	Fritchley Brook	1			16	<1.1	17		220	690
Sediment	River Derwent, downstream	4			37	<2.3	39		450	800
Water	River Derwent, upstream	1							<0.092	0.15
Water <sup>c</sup>	Fritchley Brook	1			0.025	<0.0022	0.017		<0.043	0.18
Water	River Derwent, downstream	4							<0.063	0.18
<b>Devonport</b>										
Ballan wrasse	Plymouth Sound	1 <sup>F</sup>	<0.11	<0.06				<0.04		
Spiny spider crab	Plymouth Sound	1 <sup>F</sup>	<0.12	<0.07				<0.04		
Shrimp	River Lynher	1 <sup>F</sup>	<0.56	<0.26				<0.14		
Mussels	River Lynher	1 <sup>F</sup>	<0.31	<0.25				<0.23		
Sediment <sup>e</sup>	Kinterbury	2						<0.25		
Beetroot		1 <sup>F</sup>	<0.21	<0.21				<0.14		
Grass		1 <sup>F</sup>	<0.43	<0.29				<0.21		
<b>Faslane</b>										
Mussels	Rhu	1	<0.11	<0.20				<0.11		
Winkles	Rhu	1	<0.10	<0.15				0.14		
<i>Fucus vesiculosus</i>	Rhu	1	<0.10	<0.15				<0.10		
<i>Fucus vesiculosus</i>	Garelochhead	1	<0.10	<0.10				<0.10		
<i>Fucus vesiculosus</i>	Carnban boatyard	1	<0.10	0.20				<0.11		
Sediment	Rhu	1	<0.15	0.90				<0.33		
Sediment	Garelochhead	1	<0.14	<0.23				<0.27		
Sediment	Carnban boatyard	1	<0.10	0.61				<0.10		
Seawater	Carnban boatyard	2	<0.10	<0.10				<0.10		
Beef muscle	Faslane	1						<0.11		
Honey	Faslane	1						<0.07		
Grass	Auchengaich Reservoir	1						<0.07		
Grass	Lochan Ghlas Laoigh	1						<0.06		
Soil	Auchengaich	1		1.4				0.83		
Soil	Lochan Ghlas Laoigh	1	0.26	2.4				<0.27		
Freshwater	Helensburgh Reservoir	1						<0.01	<0.010	0.017
Freshwater	Loch Finlas	1							<0.010	0.029
Freshwater	Auchengaich Reservoir	1						<0.01	<0.010	0.030
Freshwater	Lochan Ghlas Laoigh	1						<0.02	<0.010	<0.031
Freshwater	Loch Eck	1						<0.01	<0.010	0.034
Freshwater	Loch Lomond	1						<0.01	<0.010	0.033
<b>Holy Loch</b>										
Sediment	Mid-Loch	1	<0.15	0.90				0.55		
<b>Rosyth</b>										
Winkles	St David's Bay	1	<0.13	<0.24				<0.14		
<i>Fucus vesiculosus</i>	East of dockyard	1	<0.10	<0.12				<0.10		
Sediment	East of dockyard	1	<0.11	0.49				<0.25		
Sediment	Port Edgar	1	<0.13	1.7				1.3		
Sediment	West of dockyard	1	<0.12	0.69				<0.22		
Sediment	East Ness Pier	1	<0.14	<0.28				<0.26		
Sediment	Blackness Castle	1	<0.11	<0.21				<0.21		
Sediment	Charlestown Pier	1	<0.11	<0.23				<0.21		
Seawater	East of dockyard	2	<0.10	<0.11				<0.10		
Freshwater	Castlehill Reservoir	1						<0.01	<0.010	<0.013
Freshwater	Holl Reservoir	1						<0.01	<0.010	<0.049
Freshwater	Gartmorn Dam	1						<0.01	<0.010	0.15
Freshwater	Morton No. 2 Reservoir	1						<0.01	<0.010	0.33

\* Not detected by the method used

<sup>a</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima.

If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>b</sup> Except for sediment and sewage pellets where dry concentrations apply, and for water where units are Bq l<sup>-1</sup>

<sup>c</sup> The concentrations of <sup>228</sup>Th, <sup>230</sup>Th and <sup>232</sup>Th were <0.0024, <0.00080 and <0.00080 Bq l<sup>-1</sup> respectively

<sup>d</sup> The concentration of <sup>99</sup>Tc was <0.90 Bq kg<sup>-1</sup>

<sup>e</sup> The concentrations of <sup>238</sup>Pu and <sup>239+240</sup>Pu were <0.31 and 0.52 Bq kg<sup>-1</sup>

<sup>f</sup> Measurements labelled "F" are made on behalf of the Food Standards Agency, all other measurements are made on behalf of the environment agencies

**Table 5.3(b). Monitoring of radiation dose rates near defence establishments, 2015**


Establishment	Location	Ground type	No. of sampling observations	$\mu\text{Gy h}^{-1}$
<b>Mean gamma dose rates at 1m over substrate</b>				
Barrow	Walney Channel, N of discharge point	Mud	2	0.089
Barrow	Walney Channel, S of discharge point	Mud	2	0.093
Devonport	Torpoint South	Mud	1	0.082
Devonport	Torpoint South	Mud and shingle	1	0.10
Devonport	Kinterbury Access Gate	Mud	1	0.090
Devonport	Kinterbury Access Gate	Mud and shingle	1	0.084
Devonport	Lopwell	Mud	1	0.087
Devonport	Lopwell	Mud and shingle	1	0.094
Faslane	Garelochhead	Sand and shingle	1	0.061
Faslane	Garelochhead	Sand	1	0.049
Faslane	Gulley Bridge Pier	Sand and pebbles	1	0.065
Faslane	Gulley Bridge Pier	Sand	1	0.049
Faslane	Rhu	Shingle	2	0.050
Faslane	Helensburgh	Sand	2	0.058
Faslane	Carnban boatyard	Shingle	2	0.069
Faslane	Rahane	Shingle	2	0.066
Faslane	Rosneath Bay	Shingle	2	0.055
Faslane	Auchengaich	Grass	1	0.072
Faslane	Lochan Ghlas	Grass	1	0.073
Holy Loch	Kilmun Pier	Rock / Sediment	1	0.071
Holy Loch	Mid-Loch	Sediment	1	0.064
Holy Loch	North Sandbank	Rock / Sediment	1	0.071
Rosyth	Blackness Castle	Sand	2	0.053
Rosyth	Charlestown Pier	Sand	2	<0.052
Rosyth	East Ness Pier	Sand	2	0.060
Rosyth	East of Dockyard	Sand and shingle	2	<0.047
Rosyth	Port Edgar	Sand	2	0.062
Rosyth	West of Dockyard	Sand	2	0.048

## 6. Radiochemical production

This section considers the results of monitoring by the Environment Agency and FSA at two sites associated with the radiopharmaceutical industry. The sites, at Amersham and Cardiff, are operated by GE Healthcare Limited. This is a health science company functioning in world-wide commercial healthcare and life science markets.

Permits have been issued by the Environment Agency and Natural Resources Wales (NRW) to the Amersham and Cardiff sites, respectively, allowing the discharge of radioactive wastes (Appendix 2). Independent monitoring of the environment around the Amersham and Cardiff sites is conducted by the Environment Agency and FSA. The Environment Agency has an agreement with NRW to carry out monitoring on its behalf in Wales. The medium-term trends in discharges, environmental concentrations and dose at Amersham and Cardiff have been considered in a summary report (Environment Agency, FSA, NIEA and SEPA, 2010b).

### 6.1 Grove Centre, Amersham, Buckinghamshire



GE Healthcare Limited's principal establishment is located in Amersham, Buckinghamshire. It consists of a range of plants for manufacturing diagnostic imaging products for use in medicine and research. The

monitoring programme consists of analysis of fish, crops, water, sediments and environmental materials, and measurements of gamma dose rates. The monitoring locations are shown in Figure 3.4. The most recent habits survey was undertaken in 2009 (Clyne *et al.*, 2010b).

#### Doses to the public

The *total dose* from all pathways and sources of radiation was 0.14 mSv in 2015 (Table 6.1) or 14 per cent of the dose limit, and unchanged from 2014. The dominant contribution to *total dose* was from direct radiation. The representative person was an adult living in the vicinity of the site (and a change from a prenatal child in 2014). Exposure from direct radiation varies around the boundary

#### Key points

##### GE Healthcare Limited, Grove Centre, Amersham, Buckinghamshire

- *Total dose* for the representative person was unchanged in 2015. The highest dose was due to direct radiation from the site
- Gaseous discharges of "Alpha" decreased in 2015
- Concentrations of radioactivity in terrestrial and aquatic samples, and gamma dose rates, were low and generally similar to those in 2014

##### GE Healthcare Limited, Maynard Centre, Cardiff, South Glamorgan

- *Total dose* for the representative person was low and unchanged in 2015
- A revised permit for GE Healthcare Limited was granted following its application to partially surrender the environmental permit for the Maynard Centre site and liquid discharges have been removed from the permit
- Gaseous and liquid discharges of tritium and carbon-14 remained low in 2015; gaseous discharges of tritium were increased and liquid discharges of tritium were reduced to nil
- Tritium concentrations in fish species continued their long-term decline; the level in flounder in 2015 was the lowest ever reported

of the Grove Centre and therefore the *total dose* is determined as a cautious upper value. The trend in *total dose* over the period 2004 – 2015 is given in Figure 1.2. *Total doses* remained broadly similar with time (up until 2013) and were dominated by direct radiation. The lower value in 2014 (and subsequently in 2015) was due to changes in working practices (for distribution activities, products spend less time in the dispatch yard) and the construction of a shield wall on the western side of a building that contains legacy radioactive wastes.

Source specific assessments for a high-rate consumer of locally grown foods, for an angler and for a worker at Maple Lodge STW, which serves the sewers to which permitted discharges are made, give exposures that were less than the *total dose* in 2015 (Table 6.1). The dose for a high-rate consumer of locally grown foods (which included a contribution from the gaseous plume related pathways)

was 0.010 mSv, or 1 per cent of the dose limit to members of the public of 1 mSv, and similar to that in 2014 (0.009 mSv). As in previous years, atmospheric discharges of radon-222 remain the dominant contributor in 2015. It should be noted that the current assessment methodology uses a conservative dose factor based on this nuclide being in equilibrium with its daughter products. The dose to a local angler in 2015 was 0.006 mSv. The small increase in dose (from less than 0.005 mSv in 2014) was mostly attributed to the relatively higher gamma dose rate over grass of the Grand Union Canal (given in footnote, Table 6.2) in 2015.

The 2009 habits survey at Amersham did not directly identify any consumers of fish, shellfish or freshwater plants. As in previous surveys, however, there was anecdotal evidence of fish consumption, albeit occasional and at low rates. To allow for this, a consumption rate of 1 kg per year for fish has been included in the dose assessment for an angler.

The Grove Centre discharges liquid waste to Maple Lodge STW, and the proximity to raw sewage and sludge experienced by sewage treatment workers is a likely exposure pathway (National Dose Assessment Working Group, 2004). The dose received by one of these workers in 2015 was modelled using the methods described in Appendix 1. The dose from a combination of external exposure to contaminated raw sewage and sludge and the inadvertent ingestion and inhalation of re-suspended radionuclides was less than 0.005 mSv.

## Gaseous discharges and terrestrial monitoring


The Amersham facility is permitted to discharge gaseous radioactive wastes via stacks on the site. In 2015, gaseous discharges of "Alpha" decreased in comparison with releases in 2014. The results for the terrestrial monitoring are given in Table 6.2. Sulphur-35 was positively detected at a low concentration (just above the LoD) in one crop sample (wheat) in 2015. As in previous years, caesium-137 activities were detected in soil near the site, and this is likely to be due to global fallout from testing of weapons or from the Chernobyl accident.

## Liquid waste discharges and aquatic monitoring

Radioactive liquid wastes are discharged to sewers serving the Maple Lodge STW; treated effluent subsequently enters the Grand Union Canal and the River Colne. In 2015, liquid discharges were generally similar to those in recent years. The results of the aquatic monitoring programme are given in Table 6.2. Activity concentrations in freshwater, and effluent and sludge from Maple Lodge STW, are mostly reported as less than values in 2015. The sludge samples contained very low concentrations of

iodine-131 (just above the less than value, as in 2014), which were most likely to have originated from the therapeutic use of this radionuclide in a local hospital. The caesium-137 detected in sediments upstream of the STW outfall is likely to be derived from weapons test fallout or the Chernobyl accident. Tritium, gross alpha and gross beta concentrations in water (excluding the less than value reported for gross beta in the River Chess) were below the investigation levels for drinking water in the European Directive 2013/51. Gamma dose rates (given in footnote, Table 6.2) above the banks of the Grand Union Canal remained low in 2015, although the dose rates increased by small amounts in comparison to those in 2014.

## 6.2 Maynard Centre, Cardiff



GE Healthcare Limited operates a second establishment, on the Forest Farm industrial estate near Whitchurch, Cardiff. GE Healthcare Limited ceased manufacturing a range of radio-labelled products

containing tritium in 2009 and products containing carbon-14 in 2010.

On 1 June 2015, NRW issued a revised permit for GE Healthcare Limited following its application to partially surrender the environmental permit for the Maynard Centre site. This was concurrently undertaken with a parallel application to ONR to de-license around 90 per cent of the footprint of the site, following decommissioning and clean-up of the wider Maynard Centre, and to re-license the remaining 10 per cent of the site as a stand-alone nuclear licensed site. The area of the site covered by the new nuclear site licence and EPR permit is known as the Cardiff Nuclear Licensed Site (CNLS) and continues to be operated by GE Healthcare Limited. In addition to this change, liquid discharges have been removed from the permit, since the site no longer discharges radioactive liquid waste. Furthermore, gaseous carbon-14 and tritium discharges permitted under the EPR permit have been reduced significantly and gaseous discharge limits for iodine-125 and phosphorus-32/33 have been removed from the permit. The current activities at CNLS relate to the storage and repackaging of legacy ILW for off-site disposal and is located entirely within the confines of the previously licenced site (and its security boundary). Gaseous discharges from the Maynard Centre are now the result of out-gassing of tritium and carbon-14 from stored wastes with only small amounts originating from decommissioning.

GE Healthcare Limited's custom radio-labelling division was acquired by Quotient Bioresearch, (a division of Quotient Bioscience) which operates from different premises in Cardiff (a purpose-built laboratory at Trident Park). This non-nuclear facility also discharges carbon-14 and tritium to atmosphere and in liquid wastes. These are at much reduced levels in comparison to when the Maynard Centre was manufacturing radio-labelled products. The effluents discharged from the site are also treated to ensure that organic matter present is destroyed prior to discharge. The facility has an environmental permit issued and regulated by NRW.

The Environment Agency and FSA conduct a routine monitoring programme on behalf of NRW and the Welsh Government. This includes sampling of locally produced food, fish and shellfish, and external dose rate measurements over muddy, intertidal areas. Environmental materials including seawater, intertidal sediment, freshwater, seaweed, and grass provide additional information. The most recent habits survey was undertaken in 2003 (McTaggart *et al.*, 2004a).

Previous monitoring data from Cardiff has been reviewed in order to compare the apparent enhancement of tritium concentrations on uptake by marine biota with bioaccumulation at other UK sites (Hunt *et al.*, 2010). The observed enhancement factor at Cardiff remains at least an order of magnitude greater than at the other sites studied, although the organically bound fractions were uniformly high. Various earlier monitoring and research efforts have targeted OBT in foodstuffs (FSA, 2001b; Swift, 2001; Williams *et al.*, 2001; Leonard *et al.*, 2001 and McCubbin *et al.*, 2001).

## Doses to the public

The *total dose* from all pathways and sources was less than 0.005 mSv (Table 6.1) in 2015, or less than 0.5 per cent of the dose limit, and unchanged from 2014. This dose estimates take into account the increased dose coefficients for OBT derived for discharges from the Maynard Centre and includes consideration of prenatal children. The representative person was a prenatal child of occupants over sediment (and a change from an infant consuming milk at high-rates in 2014). Trends in *total doses* over time (2004 – 2015) in the Severn Estuary (and areas of the south coast) are shown in Figure 6.1. At Cardiff, the most significant reductions in the *total dose*, prior to 2007, were largely due to lower concentrations of tritium and carbon-14 in seafood. Since 2007, the *total doses* have generally continued to decrease over time and were low (and very recently, less than 0.005 mSv). The increase in *total dose* in 2013 was attributed to higher carbon-14 concentrations in milk.

Source specific assessments for a high-rate consumer of locally grown foods, a recreational user of the River Taff, and for a worker at Cardiff East Waste Water Treatment Works (WWTW), give doses that were also less than

0.005 mSv in 2015 (Table 6.1). The dose to a high-rate consumer of seafood was 0.006 mSv and unchanged from 2014. An assessment of exposure was estimated (modelled using the methods described in Appendix 1) for a worker at WWTW; the dose was much less than 0.005 mSv.

The dose coefficients for OBT differ from those for tritiated water (see Appendix 1, A3.4) and the estimates of dose to members of the public account for this. For ingestion of seafood caught near Cardiff, an area taken to be equivalent to the Bristol Channel, a dose coefficient based on a site-specific study of the consumption of fish caught in Cardiff Bay is used. An experimental study by Hunt *et al.*, (2009) suggests that this raised dose coefficient is conservative, but it is retained for 2015 dose assessments on the advice of PHE. For ingestion of other food, the ICRP dose coefficient for OBT is applied.


The monitoring locations for seafood, water, environmental materials and dose rates near the Cardiff site are shown in Figure 6.2.

## Gaseous discharges and terrestrial monitoring

The Maynard Centre discharges radioactivity to the atmosphere via stacks on the site. As a result of reduced commercial operations, in relation to the site's planned shutdown, discharges of tritium continued to be low in 2015. Carbon-14 discharges also continued to be low, although slightly higher in comparison to releases in 2014 (releases in 2014 were the lowest in recent years). There were no discharges of radionuclides other than those of tritium and carbon-14.

The focus of the terrestrial sampling was for the content of tritium, carbon-14 and sulphur-35 in milk, crops, freshwater and grass. In recent years, the Environment Agency have also analysed additional samples of sewage products from the Cardiff East WWTW. This additional monitoring ceased in 2013 and monitoring returned to normal frequencies in 2014. The constraints of the Sludge (Use in Agriculture) Regulations (United Kingdom – Parliament, 1989) (commonly referred to as the Safe Sludge Matrix) require that crops cannot be harvested within 10 months of the application of sludge pellets. An FSA research project investigated the transfer of tritium from treated soil to crops, under the Safe Sludge Matrix conditions, and concluded that the transfer of tritium to each of the crops considered was small (Ham *et al.*, 2007).

Tritium concentrations in terrestrial food samples are reported as less than values in 2015 (Table 6.3(a)). These values were similar in comparison to those in previous years and are consistent with progressive discharge reductions in recent years. Carbon-14 was detected in locally produced foods at concentrations close to background values. Carbon-14 concentrations in foodstuffs (milk) decreased



**Figure 6.1.** Total dose for major sites in the Severn Estuary and south coast, 2004-2015 (Note small doses, less than or equal to 0.005 mSv, are recorded as being 0.005 mSv)

by small amounts in comparison to those in 2014. Low concentrations of sulphur-35, which is not discharged by the site, were detected in food (potatoes) and grass samples (and were generally similar to those in 2014). Phosphorus-32 and iodine-125 concentrations are reported as less than values in all terrestrial samples. Samples of raw effluent, treated effluent and sludge pellets (analysed in previous years) from Cardiff East WWTW were not sampled in 2015. The most recent results (in 2013) show that all activity concentrations in effluent are reported as less than values (Environment Agency, FSA, NIEA, NRW and SEPA, 2014).


Unlike in previous years, there was no evidence of tritium being detected in sediment and freshwater from the Glamorganshire Canal in 2015; however, this is not used as a source of water for the public water supply. Tritium concentrations in freshwater, downstream (and upstream) from the outfall into the River Taff (potentially containing site run-off water) are reported as less than values in 2015. Freshwater samples from the outfall were not collected in 2015 (as run-off water originating from the site is not continuous); no flow of water occurred during the planned bi-annual sampling visits. The trend of discharges, with tritium concentrations in sediment from the marine and freshwater environments, over time (2004 – 2015) are shown in Figure 6.3. The overall decline in activity concentrations generally replicates that

of the tritium discharges, although the decrease in marine levels (east/west of the pipeline) is less pronounced than that in the canal sediments over the whole time period. The apparent increase in tritium concentrations, in the canal and west of pipeline samples, is because values are reported as less than values in 2015.

### Liquid waste discharges and aquatic monitoring

Up to 2015, Maynard Centre discharged liquid wastes into the Ystradyfodwg and Pontypridd (YP) public sewer. This joins the Cardiff East sewer, which after passing through a WWTW discharges into the Severn Estuary near Orchard Ledges. During periods of high rainfall, effluent from the YP sewer has been known to overflow into the River Taff. In addition, there is run-off from the site into the river via surface water drains.

The bulk of the radioactivity previously discharged to the YP sewer was tritium and carbon-14. Discharges of tritium released to the sewer reduced to nil in 2015 and discharges of carbon-14 were very low (as in recent years). Over the longer term both discharge rates have decreased substantially (Figures 6.4 and 6.5). Permission to discharge radioactive liquid waste to the sewer ceased on 1 June 2015, as a consequence of the partial surrender of




**Figure 6.2.** Monitoring locations at Cardiff, 2015 (not including farms)

the permit. The new permit does not permit discharges of radioactive waste to the sewer.


Marine sampling included locally caught seafood and indicator materials (e.g. seaweed). These were supported by external dose rate measurements over intertidal areas. The results of routine monitoring in 2015 are given in Tables 6.3(a) and (b). Tritium and carbon-14 concentrations were enhanced by small amounts in fish samples (as in 2014). Further analysis of these samples showed that a high proportion of the tritium was still associated with organic matter, a situation that has been observed since the late 1990s (McCubbin *et al.*, 2001; Leonard *et al.*, 2001; Williams *et al.*, 2001). The tritium is strongly bound to organic matter and has the potential to transfer through the marine food chain from small organisms to accumulate in fish. In 2015, the reported tritium (OBT) concentration in flounder decreased (compared with that in 2014); being the lowest value in recent years. The continued overall decline in tritium concentrations in fish from the Cardiff area is likely to be a direct response to the decreasing inputs from the Maynard Centre, as well as a shift in the composition of this discharge away from organically bound compounds. However, the annual uncertainty and

variation in certain species in recent years suggests that complex indirect uptake mechanisms continue to affect tritium concentrations in the region.


Figure 6.4 indicates that the overall tritium concentrations in fish and mollusc samples have decreased significantly over a period of time. Tritium concentrations in marine sediment samples (where comparisons can be made) were measured at similar levels to those in recent years. The mean concentrations of carbon-14 in fish and molluscs in 2015 were generally similar to those in 2014. The longer term trend in concentrations and the relationship to discharges is shown in Figure 6.5 (overall, concentrations in both species declining). Concentrations of caesium-137 in marine samples remain low and can largely be explained by other sources such as Chernobyl, weapon test fallout and discharges from other establishments such as the Hinkley Point, Berkeley and Oldbury nuclear licensed sites. Overall, gamma dose rates over sediment in 2015 (Table 6.3(b)) were generally similar to those in 2014, although the dose rates over mud at Peterstone Wentlooge decreased by a small amount (most likely due to natural variation). These rates are not (in the main) attributable to discharges from the Maynard Centre or Quotient Biosciences at Trident Park.



**Figure 6.3.** Tritium liquid discharge from Cardiff and mean concentrations in sediment near Cardiff, 2004-2015



**Figure 6.4.** Tritium liquid discharge from Cardiff and mean concentrations in fish and molluscs near Cardiff, 1997-2015 (species include all those reported in RIFE for the given year)



**Figure 6.5.** Carbon-14 liquid discharge from Cardiff and mean concentrations in fish and molluscs near Cardiff, 1997-2015 (species include all those reported in RIFE for the given year)

**Table 6.1. Individual doses – radiochemical sites, 2015**

Site	Representative person <sup>a</sup>	Exposure, mSv per year						
		Total	Fish and shellfish	Other local food	External radiation from intertidal areas or river banks	Intakes of sediment or water	Gaseous plume related pathways	Direct radiation from site
Amersham								
<b>Total dose – all sources</b>	<b>Local adult inhabitants (0 - 0.25km)</b>	<b>0.14<sup>d</sup></b>	-	<b>&lt;0.005</b>	<b>&lt;0.005</b>	-	<b>&lt;0.005</b>	<b>0.14</b>
Source	Anglers	0.006	<0.005	-	0.006	-	-	-
specific doses	Infant inhabitants and consumers of locally grown food	0.010 <sup>d</sup>	-	<0.005	-	-	0.009	-
	Workers at Maple Lodge STW	<0.005	-	-	<0.005 <sup>b</sup>	<0.005 <sup>c</sup>	-	-
Cardiff								
<b>Total dose – all sources</b>	<b>Prenatal children of occupants over sediment</b>	<b>&lt;0.005</b>	<b>&lt;0.005</b>	-	<b>&lt;0.005</b>	-	-	-
Source	Prenatal children of seafood consumers	0.006	<0.005	-	0.006	-	-	-
specific doses	Recreational users of River Taff	<0.005	-	-	<0.005	<0.005	-	-
	Infant inhabitants and consumers of locally grown food	<0.005	-	<0.005	-	-	<0.005	-
	Workers at Cardiff East WWTW	<0.005	-	-	<0.005 <sup>b</sup>	<0.005 <sup>c</sup>	-	-

- <sup>a</sup> The total dose is the dose which accounts for all sources including gaseous and liquid discharges and direct radiation. The total dose for the representative person with the highest dose presented. Other dose values are presented for specific sources, either liquid discharges or gaseous discharges, and their associated pathways. They serve as a check on the validity of the total dose assessment. The representative person is an adult unless otherwise stated
- <sup>b</sup> External radiation from raw sewage and sludge
- <sup>c</sup> Intakes of resuspended raw sewage and sludge
- <sup>d</sup> Includes a component due to natural sources of radionuclides

**Table 6.2. Concentrations of radionuclides in food and the environment near Amersham, 2015<sup>g</sup>**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						
			<sup>3</sup> H	<sup>32</sup> P	<sup>35</sup> S	<sup>131</sup> I	<sup>137</sup> Cs	Gross alpha	Gross beta
<b>Freshwater samples</b>									
Flounder	Woolwich Reach	1	<25			<0.26	<0.06		
Sediment	River Colne (Grand Union Canal)	2 <sup>E</sup>				<1.4	<1.6	160	380
Sediment	Upstream of outfall (Grand Union Canal)	2 <sup>E</sup>				<1.8	4.6	200	320
Freshwater	Maple Cross	2 <sup>E</sup>	<3.1			<0.20	<0.22	<0.080	0.57
Freshwater	Upstream of outfall (Grand Union Canal)	2 <sup>E</sup>	<3.0			<0.21	<0.22	<0.056	0.11
Freshwater	River Chess	1 <sup>E</sup>	<2.8			<0.19	<0.21	<0.11	0.085
Freshwater	River Misbourne – upstream	1 <sup>E</sup>	<3.2			<0.20	<0.20	<0.080	0.089
Freshwater	River Misbourne – downstream	1 <sup>E</sup>	<2.9			<0.20	<0.20	<0.060	<0.089
Crude effluent <sup>d</sup>	Maple Lodge Sewage Treatment Works	4 <sup>E</sup>	<9.8	<0.55	<0.41		<0.22	<0.089	0.56
Digested sludge <sup>e</sup>	Maple Lodge Sewage Treatment Works	4 <sup>E</sup>	<17	<1.8	<0.51	0.60	<0.20	<2.7	3.9
Final effluent <sup>f</sup>	Maple Lodge Sewage Treatment Works	4 <sup>E</sup>	<18	<1.2	<0.43		<0.19	<0.076	0.77
<b>Terrestrial samples</b>									
Milk		1	<3.0	<0.18	<0.0055	<0.0022	<0.06		
Apples		1	<2.9	<0.10	<0.057		<0.06		
Wheat		1	<5.5	0.60	<0.056		<0.11		
Grass	Next to site	1 <sup>E</sup>		<2.5		<1.1	<1.3	<0.38	54
Grass	Orchard next to site	1 <sup>E</sup>		<2.1		<0.88	<1.1	<1.3	140
Grass	Water Meadows (River Chess)	1 <sup>E</sup>		<2.2		<1.2	<1.5	<1.4	140
Soil	Next to site	1 <sup>E</sup>				<1.4	12	220	370
Soil	Orchard next to site	1 <sup>E</sup>				<0.50	5.0	330	660
Soil	Water Meadows (River Chess)	1 <sup>E</sup>				<1.7	15	270	360

<sup>a</sup> Except for milk, water and effluent where units are Bq l<sup>-1</sup> and for sediment and soil where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>d</sup> The concentration of <sup>3</sup>H as tritiated water was <3.6 Bq l<sup>-1</sup>

<sup>e</sup> The concentration of <sup>3</sup>H as tritiated water was <6.4 Bq l<sup>-1</sup>

<sup>f</sup> The concentration of <sup>3</sup>H as tritiated water was <3.9 Bq l<sup>-1</sup>

<sup>g</sup> The gamma dose rates in air at 1m over grass and mud, and grass on the bank of the Grand Union Canal were 0.066 and 0.066 mGy h<sup>-1</sup> respectively

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 6.3(a). Concentrations of radionuclides in food and the environment near Cardiff, 2015**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>					
			Organic <sup>3</sup> H <sup>e</sup>	<sup>3</sup> H	<sup>3</sup> H <sup>f</sup>	<sup>14</sup> C	<sup>125</sup> I	<sup>137</sup> Cs
<b>Marine samples</b>								
Flounder	East of new pipeline	2	33	<31		31		0.19
Lesser spotted dogfish	Off Orchard Ledges	1	<25	<25		28		0.35
Limpets	Lavernock Point	1	<25	<25		27		0.30
Seaweed	Orchard Ledges	2 <sup>E</sup>		<11	9.3	<11	<0.67	<0.59
Sediment	East of new pipeline	2 <sup>E</sup>		32		21	<1.3	3.8
Sediment	West of new pipeline	2 <sup>E</sup>		<150	110	<16	<2.4	18
Seawater	Orchard Ledges	2 <sup>E</sup>		<9.6	<4.1	<5.4	<0.32	<0.24

Material	Location or selection <sup>b</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>							
			Organic <sup>3</sup> H <sup>e</sup>	<sup>3</sup> H	<sup>3</sup> H <sup>f</sup>	<sup>14</sup> C	<sup>35</sup> S	<sup>125</sup> I	<sup>137</sup> Cs	Gross alpha
<b>Terrestrial samples</b>										
Milk <sup>g</sup>		2	<2.5	<2.5		18	<0.40	<0.0061	<0.07	
Milk <sup>g</sup>	max		<2.6	<2.6		19	<0.48	<0.0068		
Potatoes		1	<3.0	<3.0		21	0.80	<0.078	<0.07	
Grass		1	<7.0	<7.0		11	1.9	<0.12	0.09	
Sediment	Canal	2 <sup>E</sup>		<97		<45	<2.8	8.3		
Freshwater	River Taff upstream	2 <sup>E</sup>		<11	<3.1	<5.4	<0.18	<0.23	<0.057	0.21
Freshwater	River Taff downstream	2 <sup>E</sup>		<11	<2.9	<4.5	<0.16	<0.21	<0.046	0.17
Freshwater	Canal	2 <sup>E</sup>		<13	<4.5	<4.4	<0.16	<0.22	<0.067	<0.066

<sup>a</sup> Except for milk, water and effluent where units are Bq l<sup>-1</sup> and for sediment and sludge pellets where dry concentrations apply

<sup>b</sup> Data are arithmetic means unless stated as 'max' in this column. 'Max' data are selected to be maxima. If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>d</sup> The concentration of <sup>99</sup>Tc was <1.2 Bq kg<sup>-1</sup>

<sup>e</sup> The organic fraction may be higher than the total tritium value for some analyses due to uncertainties in the analytical methods for tritium. For dose assessments in this report, the higher of the two values has been used

<sup>f</sup> As tritiated water

<sup>g</sup> The concentration of <sup>32</sup>P was <0.20 (max <0.21) Bq l<sup>-1</sup>

<sup>E</sup> Measurements labelled "E" are made on behalf of the Environment Agency, all other measurements are made on behalf of the Food Standards Agency

**Table 6.3(b). Monitoring of radiation dose rates near Cardiff, 2015**

Location	Ground type type	No. of sampling observations	µGy h <sup>-1</sup>
<b>Mean gamma dose rates at 1m over substrate</b>			
East of Pipeline	Mud	1	0.075
East of Pipeline	Mud and sand	1	0.080
West of Pipeline	Mud and stones	1	0.10
West of Pipeline	Rock and shingle	1	0.085
Peterstone Wentlooge	Mud	2	0.075



## 7. Industrial and landfill sites

This section considers the effects of (i) the main disposal site on land for solid radioactive wastes in the UK, at the LLWR near Drigg in Cumbria, as well as other landfill sites which have received small quantities of solid wastes and (ii) other sites where industries or incidents may have introduced radioactivity into the environment.

### 7.1 Low Level Waste Repository near Drigg, Cumbria

The LLWR is the UK's national facility for the disposal of lower activity waste and is located on the west Cumbrian coast, approximately 7 km south east of Sellafield. The main function of the LLWR is to receive solid radioactive wastes from all UK nuclear licensed sites (except Dounreay) and many non-nuclear sites. Where possible the waste is compacted, and then most waste is grouted within containers before disposal. Wastes may be disposed of in engineered concrete vaults on land, whereas prior to the early 1990s waste was disposed of in open clay lined trenches. The site is owned by the NDA and operated on their behalf by LLWR Limited on behalf of NDA. From April 2008, a consortium, UK Nuclear Waste Management Limited (UKNWML), took over as the PBO for LLWR Limited. A plan setting out the long term future of the site through to its final closure, currently planned for 2079, has been published (LLWR Repository Ltd, 2015). In October 2013 LLWR Limited applied to vary its environmental permit to continue to dispose of radioactive waste at the site. The Environment Agency consulted on its draft decision in May 2015. It carefully considered all of the comments made during consultation and granted a new permit. The permit covers disposal of gaseous, liquid and solid wastes and took effect from 1st November 2015.

The disposal permit allows for the discharge of leachate from the site through a marine pipeline. These discharges are small compared with those discharged from the nearby Sellafield site (Appendix 2). Marine monitoring of the LLWR is therefore subsumed within the Sellafield programme, described in Section 2. The contribution to exposures due to LLWR discharges is negligible compared with that attributable to Sellafield and any effects of LLWR discharges in the marine environment could not, in 2014, be distinguished from those due to Sellafield. The most recent habits survey was published in 2013 and the results have been included in the dose assessments for the site (Clyne *et al.*, 2013c).

Site operators were granted a new permit to allow for continued waste disposal at the site, effective from 1st November 2015. No disposals of solid radioactive

### Key points

#### LLWR, near Drigg

- Site operators were granted a new permit (effective 1 November 2015) to allow for continued waste disposal at the site. Planning permission was granted by Cumbria County Council in 2016
- Concentrations and dose rates at the LLWR were similar to those in 2014
- Doses were dominated by the effects of the legacy of discharges into the sea at Sellafield and Whitehaven

#### Other sites

- Tritium was found in leachate from other landfill sites. Probably due to disposal of Gaseous Tritium Light Devices. Doses were less than 0.5 per cent of dose limit
- Monitoring relating to the disposal of LLW at East Northants Resource Management Facility (ENRMF) near Kings Cliffe showed similar results to other landfill sites
- Very small discharges from the Studsvik Metals Recycling Facility (MRF) were made in 2015
- Enhancement in natural radionuclides at Whitehaven from historical phosphate processing is now very difficult to distinguish within the expected range due to natural sources. However, the radiation dose from the enhancement (compared to expected values), taken with effects of disposal of other local wastes, was estimated to be 42 per cent of the dose limit
- The investigation into the radium-226 contamination at Dalgety Bay, Fife continued in 2015
- Discharges from other non-nuclear sites (hospitals, universities etc.) were all within limits set in regulations. Limited monitoring of such sites was undertaken and no significant effects were found

waste were made at the LLWR in 2015, with waste accepted for storage since 2006. Planning permission enabling waste disposal into the future was granted by Cumbria County Council in July 2016.

Although the permit for disposal to the Drigg Stream has been revoked, reassurance monitoring of samples of

water and sediment has continued. The results are given in Table 7.2. The tritium, gross alpha and gross beta concentrations in the stream were below the investigation levels for drinking water in the European Directive 2013/51. Although the stream is not known to be used as a source of drinking water, it is possible that occasional use could occur, for example by campers. If the stream was used as a drinking water supply for three weeks, the dose would be less than 0.005 mSv. Concentrations of radionuclides in sediment from the Drigg stream were similar to those for 2014. They reflect the legacy of direct discharges of leachate from the disposal site into the stream (BNFL, 2002). This practice stopped in 1991.

In the past, groundwater from some of the trenches on the LLWR site moved eastwards towards a railway drain along the perimeter of the site. Radioactivity from the LLWR was detected in the drain water. The previous operators of the site (BNFL) took steps in the early 1990s to reduce ingress of water from the trenches by building a “cut-off wall” to reduce lateral migration of leachate. The results of monitoring in the drain in 2015 have shown that the activity concentrations are now very low and have reduced significantly since the “cut-off wall” was constructed. Tritium, gross alpha and gross beta concentrations were below or just above the investigation levels for drinking water in the European Directive 2013/51.

The monitoring programme of terrestrial foodstuffs at the site was primarily directed at the potential migration of radionuclides from the waste burial site via groundwater since the disposals of gaseous wastes are very small. Results for 2015 are given in Table 7.2. Evidence in support of the proposition that radioactivity in leachate from the LLWR might be transferring to foods was very limited in 2015, as it was in 2014. In general, concentrations of radionuclides detected were similar to or lower than those found near Sellafield (Section 2). The *total dose* from all pathways and sources, including a component due to Chernobyl and weapon test fallout, was 0.42 mSv, or 42 per cent of the dose limit for members of the public of 1 mSv (Table 7.1). This was dominated by the effects of the legacy of discharges into the sea at Sellafield and Whitehaven, which are near to the LLWR site. If these effects were to be excluded, and the sources of exposure from the LLWR are considered, the representative person was a 10 y old child spending time near the site. Their *total dose* in 2015 was 0.041 mSv (Table 1.2), mostly due to direct radiation. Source specific assessments of exposures for consumers of water from Drigg stream and of locally grown terrestrial food were less than 0.005 mSv.

## 7.2 Other landfill sites

Some organisations are granted authorisations or permits by SEPA (in Scotland) or the Environment Agency (in England and Wales) respectively to dispose of solid wastes containing low levels of radioactivity to approved landfill sites. In Northern Ireland, this type of waste is transferred




Figure 7.1. Landfill sites monitored in 2015

to Great Britain for incineration. Waste with very low levels of radioactivity can also be disposed of in general refuse. Radioactivity in wastes can migrate into leachate and in some cases can enter the groundwater. SEPA and the Environment Agency carry out monitoring of leachates. The distribution of landfill sites considered in 2015 is shown in Figure 7.1 and the results are presented in Tables 7.3 and 7.4.

The results, in common with previous years, showed evidence for migration of tritium from some of the disposal sites. The reported tritium concentrations vary from year to year. The variation is thought to be related to changes in rainfall quantity and resulting leachate production and the use of different boreholes for sampling. A possible source of the tritium is thought to be due to disposal of Gaseous Tritium Light Devices (Mobbs *et al.*, 1998). Inadvertent ingestion of leachate (2.5 l per year) from the site with the highest observed concentration of tritium would result in a dose of less than 0.005 mSv or less than 0.5 per cent of the dose limit for members of the public of 1 mSv (Table 7.1). Similarly, the dose from ingestion of uranium isotopes in leachate from Clifton Marsh was also less than 0.005 mSv.

In March 2007, the UK Government introduced a more flexible framework for the disposal of certain categories of LLW to landfill. Further details and information are provided on the GOV.UK website: <https://www.gov.uk/government/policies/managing-the-use-and-disposal-of-radioactive-and-nuclear-substances-and-waste/supporting->

pages/providing-policy-for-the-safe-and-secure-disposal-of-radioactive-waste.

In England and Wales, disposal of LLW at landfill sites requires both landfill companies and nuclear operators to hold permits to dispose of LLW. The 2007 Government policy led to applications from landfill operators for permits to dispose of LLW at their sites. The landfill sites were:

- Waste Recycling Group (WRG) Limited at the Lillyhall Landfill Site in Cumbria. Their permit, issued in 2011, allows them to dispose of VLLW
- Augéan at the East Northants Resource Management Facility (ENRMF), near Kings Cliffe, Northamptonshire. Their permit, issued in 2011, allows them to dispose of low activity LLW and VLLW. The Environment Agency is currently considering an application by Augéan to include an additional landfill area in the west of the site and to clarify the approaches necessary to take into account the distribution of activity which is likely in the type of wastes received. An environmental safety case has been prepared (Mitchell, N., 2015).
- SITA (Lancashire) Limited at Clifton Marsh in Lancashire. They received a permit to dispose of LLW in September 2012. This permit replaced previous arrangements authorising disposals at Clifton Marsh by operators at the Springfields and Capenhurst nuclear licensed sites, whose permits used to allow disposal of solid LLW at Clifton Marsh in their own right. The varied permits now allow those operators to transfer LLW to landfill operators who hold an appropriate EPR 10 permit.

Disposals of LLW at Clifton Marsh have continued under the new permitting arrangements.

Disposals of LLW at the ENRMF, near Kings Cliffe, site began in December 2011 and were from non-nuclear site remediation works. The first consignment from a nuclear licensed site was from Harwell in March 2012 this comprised soil, concrete, rubble and clay pipes from the drains on the Harwell site. In parallel, the Environment Agency began a programme of monitoring within and around the ENRMF landfill site, near Kings Cliffe, in order to provide a baseline and allow any future changes to be detected. In 2015, samples were taken and analysed for radiological composition from both upstream and downstream groundwater boreholes, off-site watercourses, on-site surface water, on-site leachate and on-site soil. Samples of leachate, borehole water and surface water were taken and filtered. Both the filtrate and the particulate were analysed for their radioactivity content, along with some bulk water samples. The results are given in Table 7.5. The results for man-made radionuclides were generally at limits of detection or at low levels expected due to UK-wide fallout from weapon testing and overseas accidents. A low level of tritium was detected in one borehole sample. It was below the investigation level of 100 Bq l<sup>-1</sup> for drinking water in the European Directive 2013/51. Naturally occurring radionuclides were present at levels expected due to natural sources. Gross alpha

and gross beta concentrations in water were below the investigation levels for drinking water in the European Directive 2013/51 of 0.1 and 1.0 Bq l<sup>-1</sup> respectively. No use of water for drinking has been observed. Where sampling was repeated, the results were similar to those in previous years. In 2014, leachate samples were analysed and were found to contain tritium levels in excess of 100 Bq l<sup>-1</sup> (Environment Agency, FSA, FSS, NIEA, NRW and SEPA, 2015). This likely to be due to the disposal of Gaseous Tritium Light Devices. An assessment of exposure based on inadvertent ingestion of borehole or surface water at concentrations presented in Table 7.5 gives a dose of less than 0.005 mSv or less than 0.5 per cent of the dose limit for members of the public of 1 mSv (Table 7.1). The assessment excludes potassium-40 because its presence in the body is homeostatically controlled.

SEPA continued its programme of monitoring at the Stoneyhill Landfill Site in Aberdeenshire which is authorised to dispose of conditioned NORM waste. This programme is complementary to, but independent of, the authorisation holder's monitoring programme.

NORM is found within oil and gas reserves and is consequently extracted along with the oil and gas. The NORM can precipitate onto oil and gas industry equipment creating an insoluble scale (NORM scale). The presence of this scale reduces the efficiency of the equipment and must be removed. SITA UK Limited, who operates Stoneyhill Landfill site, has constructed a descaling facility adjacent to the landfill in partnership with Nuvia Limited. This facility descales oil and gas industry equipment (such as pipes) using pressurised water. The solid scale removed from the equipment is then grouted into drums and can be consigned to Stoneyhill Landfill site in accordance with their authorisation granted in May 2012.

The SEPA monitoring programme involves the collection and analysis of landfill leachate, groundwater and surface water on a quarterly basis and analysing for radium-226 and radium-228, with results reported as less than values in 2015 (Table 7.6).

### 7.3 Metals Recycling Facility, Lillyhall, Cumbria

The Metals Recycling Facility (MRF), operated by Studsvik UK Limited, first commenced operations in September 2009. The facility is located on the north-eastern edge of the Lillyhall Industrial Estate, about 4 km south-east of Workington. The main function of the MRF is to receive, sort, segregate, monitor and size reduce metallic low level radioactive waste (LLW) before either treating it on site by surface decontamination, or sending the metal to a sister plant in Sweden for melting. The intent of the process is, as far as possible, to decontaminate the metal, such that it can be returned to the open market as exempt from control as radioactive waste, for recycling. Secondary wastes from the metal treatment containing radioactivity,

as either LLW or very low level waste (VLLW), are disposed of to the LLWR or to landfills.

A permit for disposal of radioactive waste from the site was issued by the Environment Agency in March 2008, although no radioactive waste disposals were made until September 2009. The permit allows discharges of gaseous waste to the environment via a main stack and aqueous waste to the sewer. Low discharge limits are set for both aqueous and gaseous discharges. Very small discharges were made during 2015 (Appendix 2). The permit includes conditions requiring Studsvik UK Limited to monitor discharges and undertake environmental monitoring.

## 7.4 Phosphate processing, Whitehaven, Cumbria



An important historic man-made source of naturally occurring radionuclides in the marine environment has been the chemical plant at Whitehaven in Cumbria, which used to manufacture phosphoric acid

from imported phosphate ore (Rollo *et al.*, 1992). Phosphogypsum, containing thorium, uranium and their daughter products, was discharged as a liquid slurry by pipeline to Saltom Bay. Processing of phosphate ore ceased in 1992 and processing of phosphoric acid at the plant ceased at the end of 2001. However, an environmental legacy has occurred from these past operations. Such sources are said to give rise to Technologically enhanced Naturally Occurring Radioactive Material (TNORM). Decommissioning of the plant was undertaken in 2002 and released small quantities of uranium to sea, but discharges were very much lower than in previous years. The plant was subsequently demolished in 2004 and the permit to discharge radioactive wastes revoked by the Environment Agency.

The results of routine monitoring for naturally occurring radioactivity near the site in 2015 are shown in Table 7.7. Analytical effort has focused on lead-210 and polonium-210, which concentrate in marine species and are the important radionuclides in terms of potential dose to the public. Concentrations of polonium-210 and other naturally occurring radionuclides were slightly enhanced near Whitehaven but quickly reduce to background levels further away. Figures 7.2 and 7.3 show how concentrations of polonium-210 in winkles and crabs have generally decreased since 1998. Concentrations in the early 1990s were in excess of 100 Bq kg<sup>-1</sup> (fresh weight). There were some variations in concentrations


of polonium-210 in local samples in 2015 compared with 2014. In particular, there were small increases in concentrations in crabs and lobsters. However, polonium-210 concentrations in these samples continued to be within or close to the expected range due to natural sources in 2015. For these and other seafood samples, it is now difficult to distinguish between the measured radionuclide concentrations and the range of concentrations normally expected from naturally sourced radioactivity. The latter are shown in Figures 7.2 and 7.3 and in Appendix 1 (Annex 4). There were small enhancements for some samples at other locations above the expected natural background median levels for marine species, but the majority were within the ranges observed in the undisturbed marine environment. It is considered prudent to continue to estimate doses at Whitehaven based on the positive difference, if any, between observed concentrations and median levels indicative of natural background. A recent analysis has confirmed that this approach is unlikely to underestimate doses (Dewar *et al.*, 2014).

The critical radiation exposure pathway considered for the assessment at Whitehaven was internal


irradiation, due to the ingestion of naturally occurring radioactivity in local fish and shellfish. The representative person was a consumer who, centred on the Sellafield site to the south of Whitehaven, obtained their sources of seafood from locations such as Whitehaven, Saltom Bay and Parton. This consumer is also considered in the assessment of the marine impacts of the Sellafield and LLWR (near Drigg) sites (Sections 2 and 7). An additional, smaller area limited to Saltom Bay is no longer assessed separately because the larger area provides adequate protection and a more robust assessment. The estimated contribution due to background median concentrations of naturally occurring radionuclides is subtracted. Consumption rates for people who eat at high-rates were reviewed and revised in 2015. The dose coefficient for polonium-210 is based on a value of the gut transfer factor of 0.5 for all foods.

The *total dose* to a local high rate consumer of seafood was 0.42 mSv in 2015 (Table 7.1), below the dose limit for members of the public of 1 mSv. The value for 2014 was 0.22 mSv. The dose includes the effects of all sources near the site: enhanced naturally occurring radionuclides from the non-nuclear industrial activity (i.e. TNORM) and Sellafield operations. The source specific assessment of dose, targeted directly at a high-rate seafood consumer confirms the *total dose* assessment and gives a similar result in 2015, 0.37 mSv.

The contribution to the *total dose* from enhanced natural radionuclides was 0.35 mSv in 2015, compared with 0.15 mSv in 2014. This increase in dose was largely due to small increases in the gross concentrations of polonium-210 in locally caught crabs and lobsters. Polonium-210 is an important radionuclide in that small



**Figure 7.2.** Polonium-210 discharge from Whitehaven and concentration in winkles at Parton, 1990-2015




**Figure 7.3.** Polonium-210 discharge from Whitehaven and concentration in crabs at Parton, 1990-2015

increases in net levels (after the natural background is subtracted) significantly influences the dose contribution from this radionuclide and similarly the increase in value of the estimated dose. The largest contribution to dose to a seafood consumer near Whitehaven and Sellafield is now from historic discharges from Whitehaven. The longer term trend in *total dose* over the period 2004 – 2015 is shown in Figure 7.4. The overall reduction in *total dose* (with some variability from year to year), up to 2010, reflects changes in both polonium-210 concentrations and consumption rates, primarily of lobsters and molluscs. Thereafter, variations in *total dose* over the period 2011 – 2014 reflect changes in polonium-210 concentrations, consumption rates and the range of seafood species

consumed by individuals at high-rates, including that of lobsters and fish.

## 7.5 Aberdeen

Scotoil operates a cleaning facility for equipment from the oil and gas industry contaminated with enhanced concentrations of radionuclides of natural origin. They are authorised to discharge liquid effluent to sea with the primary discharge being radium-226 and radium-228, with lead-210 and polonium-210 in smaller quantities. The authorisation includes conditions requiring Scotoil to undertake environmental monitoring. Prior to their



**Figure 7.4.** Trend in total dose to seafood consumers from naturally-occurring radionuclides near Whitehaven, 2004-2015

operations, a fertiliser manufacturing process was operated on the site, which made discharges to sea.

Seaweed (*Fucus vesiculosus*) from Aberdeen Harbour was monitored in 2015. Technetium-99 was detected in seaweed (9.2 Bq kg<sup>-1</sup>, fresh), in line with the expected effect from Sellafield discharges (as the releases become diluted or mixed in moving further afield). Gamma-emitting radionuclides were all below the LoD. In 2015, the dose rate on sediment was 0.087 µGy h<sup>-1</sup> and similar to background. The dose rate was lower than the results in earlier years when discharges were higher.

## 7.6 Dalgety Bay, Fife

Radioactive items containing radium-226 and associated daughter products have been detected at Dalgety Bay in Fife since at least 1990. The contamination is associated with historical disposals of waste from past military operations at the Royal Naval Air Station (RNAS) Donibristle, which closed in 1959 and upon which large areas of the town of Dalgety Bay have been built. The air station played a role as an aircraft repair, refitting and salvage yard. It is believed that waste was incinerated and the resultant ash and clinker was disposed of by reclaiming land from the sea. Following years of erosion at the site the contamination is being exposed on and adjacent to the foreshore. Some of the incinerated material contained items such as dials and levers which had been painted with luminous paint containing radium-226.

In June 1990, environmental monitoring showed elevated radiation levels in the Dalgety Bay area. The monitoring was undertaken as part of the routine environmental monitoring programme for Rosyth Royal Dockyard conducted in accordance with the dockyard's authorisation to dispose of liquid radioactive effluent to the Firth of Forth. Some material was removed for analysis, which indicated the presence of radium-226. Further investigation confirmed that the contamination could not

have originated from the dockyard and was most likely to be associated with past practices related to the nearby former RNAS Donibristle/HMS Merlin military airfield. Since this initial discovery, there have been several monitoring exercises to determine the extent of this contamination.

Following the increased number of particle finds and the discovery of the high activity particles in 2011, additional public protection measures were established and these were maintained during 2015 and into 2016. A monthly beach monitoring and particle recovery programme was adopted in 2012 by a contractor working on behalf of the MoD and this remains in place. The fence demarcating the area where the highest activity particles were discovered remains in place, as well as the information signs advising the public of the contamination and precautions to be taken. In addition, the FEPA Order issued by the Food Standards Agency in Scotland prohibiting the collection of seafood from the Dalgety Bay area remains in force. SEPA undertook a programme of shellfish monitoring between February 2012 and February 2013 during which no particles were detected in the shellfish. All shellfish samples collected were analysed for the presence of radium-226 and all were found to be less than the LoD. During routine monitoring of mussel beds in November 2015 a particle was detected in this area (for the first time since 2011) and retrieved, indicating that the continuation of these protection measures is reducing the risks to members of the public whilst further work continues to address the contamination.

Following the publication of the risk assessment together with the appropriate persons report in 2013, COMARE recommended at its meeting in July 2013 that effective remediation of the affected area is undertaken as soon as is possible. This recommendation, amongst others, was subsequently published in May 2014 in COMARE's 15<sup>th</sup> report. The MoD has progressed with addressing the contamination by initially publishing its Outline Management Options Appraisal Report in January 2014 followed by the publication in July 2014 of its broad

management strategy and timescale for implementation of its preferred management option. Copies of these reports are available on the UK Government website.

Work continues towards the implementation of the preferred management option with the convening of the Dalgety Bay Implementation Group. The Dalgety Bay Permitting Authorities Group has also been convened to ensure that any permits or licences required to proceed with the management option can be in place to allow the addressing of the contamination.

For further information on the work at Dalgety Bay please visit the Radioactive Substances pages on SEPA's website ([www.sepa.org.uk](http://www.sepa.org.uk)).

## 7.7 Former RAF Kinloss, Moray

Radioactive items containing radium-226 and associated daughter products have been detected on an area of land which used to form part of the former RAF Kinloss, now Kinloss Barracks. The contamination is associated with historic disposals of waste from past military operations at the site resulting from the dismantling of aircraft no longer required by the RAF following World War II. During the late 1940s the aircraft were stripped for their scrap metal, with the remains being burnt and/or buried at the site. The source of the radium-226 and associated daughter products are the various pieces of aircraft instrumentation which were luminised with radium paint.

SEPA has undertaken monitoring surveys at the site which positively identified the presence of radium-226 and has published an assessment of the risks posed to the public. Currently, the site is largely undeveloped open land covered in gorse with a number of wind turbines and access tracks. The area is used by visitors and for dog-walking with a number of informal paths crossing the land. The contamination detected at the site is all currently buried at depth. Current uses of the site do not involve intrusion into the ground to any significant depth; thus there is no current pathway for exposure via skin contact, ingestion or inhalation. Exposure via external gamma irradiation is possible but is significantly below the relevant dose criteria detailed in the Radioactive Contaminated Land (RCL) Statutory Guidance.

The risk assessment concluded that under its current use there are no viable or credible exposure pathways for the public to be exposed to the contamination and that this site does not currently meet the definition of radioactive contaminated land (Natural Scotland and SEPA, 2016). However, SEPA will keep this site under review, as a change in land use on the site may alter the potential exposure pathways. To access the full risk assessment report please visit the Radioactive Substances pages on SEPA's website ([www.sepa.org.uk](http://www.sepa.org.uk)).

## 7.8 Other non-nuclear sites

Routine discharges of small quantities of radioactive wastes to air and water are made from a wide range of other non-nuclear sites in the UK on land, and from offshore oil and gas installations.

A summary of the most recent data for the quantities discharged under regulation is given in Tables 7.8 and 7.9. The data are grouped according to the main industries giving rise to such wastes in the UK and exclude information for other industries considered in other sections of this report, principally the nuclear sector. The main industries are:

- Oil and gas (onshore)
- Education (Universities and Colleges)
- Hospitals
- Other (research, manufacturing and public sector)

Discharges may also occur without an authorisation or permit when the quantities are considered to be below the need for specific regulatory control. For example, discharges of natural radionuclides are made from coal-fired power stations because of the presence of trace quantities of uranium and thorium and their decay products in coal.

As indicated in Section 1, general monitoring of the British Isles as reported elsewhere in this report has not detected any gross effects from non-nuclear sources. Occasionally, routine programmes directed at nuclear licensed site operations detect the effects of discharges from the non-nuclear sector and, when this occurs, a comment is made in the relevant nuclear licensed site text. The radiological impact of the radioactivity from the non-nuclear sector detected inadvertently in this way is very low.

Monitoring of the effects of the non-nuclear sector is limited because of the relatively low impact of the discharges. However, programmes are carried out to confirm that impacts are low and, when these occur, they are described in this report.

In 2015, SEPA undertook a small-scale survey (as part of the annual programme) of the effects of discharges from non-nuclear operators by taking and analysing samples of mussels and other materials from the River Clyde, the Firth of Forth and sludge pellets from a STW. The results are given in Table 7.10. They show the expected effects of Sellafield discharges at this distance and the presence of iodine-131 in sludge pellets, probably from a hospital source. The results were generally similar to those in 2014. An assessment of the dose to a representative high-rate mollusc consumer was undertaken. The dose was less than 0.005 mSv or less than 0.5 per cent of the dose limit.

**Table 7.1. Individual doses – industrial and landfill sites, 2015**

Site	Representative person <sup>a,b</sup>	Exposure, mSv per year					
		Total	Seafood (nuclear industry discharges)	Seafood (other discharges)	Other local food	External radiation from intertidal areas	Intakes of sediment and water
<b>Total dose – all sources</b>							
Whitehaven and LLWR near Drigg	<b>Adult mollusc consumers</b>	<b>0.42<sup>d</sup></b>	<b>0.061</b>	<b>0.35</b>	–	<b>0.017</b>	–
<b>Source specific doses</b>							
LLWR near Drigg	Infant consumers of locally grown food	<0.005	–	–	<0.005	–	–
	Consumers of water from Drigg stream	<0.005 <sup>c</sup>	–	–	–	–	<0.005
Landfill sites for low-level radioactive wastes	Inadvertent leachate consumers (infants)	<0.005	–	–	–	–	<0.005
Whitehaven (habits averaged 2011–15)	Seafood consumers	0.37 <sup>d</sup>	0.051	0.29	–	0.033	–

- <sup>a</sup> The total dose is the dose which accounts for all sources including gaseous and liquid discharges and direct radiation. The total dose for the representative person with the highest dose presented. Other dose values are presented for specific sources, either liquid discharges or gaseous discharges, and their associated pathways. They serve as a check on the validity of the total dose assessment. The representative person is an adult unless otherwise stated
- <sup>b</sup> None of the people represented in this table were considered to receive direct radiation from the sites listed
- <sup>c</sup> Includes a component due to natural sources of radionuclides
- <sup>d</sup> Includes the effects of discharges from the adjacent Sellafield site

**Table 7.2. Concentrations of radionuclides in terrestrial food and the environment near Drigg, 2015**

Material	Location or selection <sup>a</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>								
			<sup>3</sup> H	<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>95</sup> Nb	<sup>95</sup> Zr	<sup>99</sup> Tc	<sup>106</sup> Ru	<sup>125</sup> Sb
Milk		1	<2.7	16	<0.06	0.037	<0.07	<0.09	<0.041	<0.48	<0.14
Deer muscle		1	<4.4	34	<0.09	<0.047	<0.11	<0.12	<0.22	<0.47	<0.15
Eggs		1	<2.0	40	<0.06	<0.045	<0.05	<0.15		<0.70	<0.20
Potatoes		1	<3.4	20	<0.06	<0.072	<0.09	<0.13	<0.22	<0.43	<0.11
Sheep muscle		1	<4.8	36	<0.03	<0.043	<0.06	<0.15	<0.21	<0.64	<0.10
Sheep offal		1	<22	30	<0.03	<0.043	<0.05	<0.07	<0.28	<0.38	<0.04
Grass		1	<6.9	14	<0.06	0.38	<0.09	<0.11	<0.22	<0.42	<0.12
Sediment	Drigg Stream	4 <sup>E</sup>			<0.42	<3.5	<0.29	<0.91		<3.2	<1.7
Freshwater	Drigg Stream	4 <sup>E</sup>			<0.30	0.12					
Freshwater	Railway drain	1 <sup>E</sup>			<0.33	0.76					

Material	Location or selection <sup>a</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>								
			<sup>129</sup> I	<sup>134</sup> Cs	<sup>137</sup> Cs	Total Cs	<sup>144</sup> Ce	<sup>210</sup> Po	<sup>228</sup> Th	<sup>230</sup> Th	<sup>232</sup> Th
Milk		1	<0.0033	<0.06	<0.13		<0.35				
Deer muscle		1	<0.014		2.8	3.0	<0.36				
Eggs		1	<0.032		<0.09	0.10	<0.46				
Potatoes		1	<0.020	<0.06	<0.04	0.038	<0.35				
Sheep muscle		1	<0.013		0.46	0.49	<0.35				
Sheep offal		1	<0.015		0.28	0.52	<0.32				
Grass		1	<0.022	<0.05	0.39	0.27	<0.31				
Sediment	Drigg Stream	4 <sup>E</sup>		<0.38	120		<1.7	8.6	16	10	12
Freshwater	Drigg Stream	4 <sup>E</sup>		<0.29	<0.24			<0.0020	<0.0034	<0.0015	<0.0011
Freshwater	Railway drain	1 <sup>E</sup>		<0.32	<0.25			<0.0024	<0.0031	<0.0018	<0.0010

Material	Location or selection <sup>a</sup>	No. of sampling observations <sup>c</sup>	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>								
			<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Pu	<sup>241</sup> Am	Gross alpha	Gross beta
Milk		1				<0.000038	<0.000029	<0.12	<0.000027		
Deer muscle		1				<0.00012	<0.000042	<0.20	0.00021		
Eggs		1				<0.000021	0.000035	0.12	0.000059		
Potatoes		1				<0.000060	0.00018	<0.32	0.00010		
Sheep muscle		1				0.000093	0.00050	<0.19	0.00073		
Sheep offal		1				0.00044	0.0029	<0.20	0.0045		
Grass		1				0.0031	0.017	0.17	0.034		
Sediment	Drigg Stream	4 <sup>E</sup>	47	<2.0	41	9.0	62	260	61	<300	490
Freshwater	Drigg Stream	4 <sup>E</sup>	0.0077	<0.0015	0.0076	<0.0032	<0.0019	<0.14	<0.0044	<0.049	0.42
Freshwater	Railway drain	1 <sup>E</sup>	0.022	<0.0016	0.018	<0.0019	<0.00090	<0.10	<0.0025	<0.023	1.8

<sup>a</sup> Data are arithmetic means unless stated as 'max'. 'Max' data are selected to be maxima.

If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>b</sup> Except for milk and freshwater where units are Bq l<sup>-1</sup>, and for sediment where dry concentrations apply

<sup>c</sup> The number of farms from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

<sup>E</sup> Measurements are made on behalf of the Food Standards Agency unless labelled "E".

In that case they are made on behalf of the Environment Agency

**Table 7.3. Concentrations of radionuclides in surface water leachate from landfill sites in Scotland, 2015**

Area	Location	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>			
			<sup>3</sup> H	<sup>14</sup> C	<sup>137</sup> Cs	<sup>241</sup> Am
Aberdeen City	Ness Tip	1	<5.0	<15	<0.05	<0.05
City of Glasgow	Summerston Tip	1	130	<15	<0.05	<0.05
City of Glasgow	Cathkin	1	400	<15	<0.05	<0.05
Clackmannanshire	Black Devon	1	26	<15	<0.05	<0.05
Dunbartonshire	Birdston	1	<5.0	<15	<0.05	<0.05
Dundee City	Riverside	1	11	<15	<0.05	<0.05
Edinburgh	Braehead	1	<5.0	<15	<0.05	<0.05
Fife	Balbarton	1	66	<15	<0.05	<0.05
Fife	Melville Wood	1	130	<15	<0.05	<0.05
Highland	Longman Tip	1	<5.0	<15	<0.05	<0.05
North Lanarkshire	Dalmacoulter	1	300	<15	<0.05	<0.05
North Lanarkshire	Kilgarth	1	<5.0	<15	<0.05	<0.05
Stirling	Lower Polmaise	1	520	<15	0.07	<0.05

**Table 7.4. Concentrations of radionuclides in water from landfill sites in England and Wales, 2015**

Location	Sample source	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>						
			Total <sup>3</sup> H	<sup>3</sup> H <sup>a</sup>	<sup>14</sup> C	<sup>40</sup> K	<sup>60</sup> Co	<sup>137</sup> Cs	<sup>228</sup> Th
<b>Glamorgan</b>									
Trecatti Landfill, Merthyr Tydfil	Raw leachate	2	480	490	<3.1				
Trecatti Landfill, Merthyr Tydfil	Treated leachate	2	630	560	<3.8				
<b>Lancashire</b>									
Clifton Marsh	Borehole 6	2		<3.5		<4.5	<0.23	<0.20	
Clifton Marsh	Borehole 19	2		<4.0		<3.8	<0.22	<0.20	
Clifton Marsh	Borehole 40	2		<3.2		<5.3	<0.27	<0.23	
Clifton Marsh	Borehole 59	2		8.1		<3.6	<0.22	<0.19	
<b>South Glamorgan</b>									
Lamby Way Tip <sup>b</sup>	Borehole 1A	2		10	<3.6	<4.4	<0.23	<0.20	
Location	Sample source	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>						
			<sup>230</sup> Th	<sup>232</sup> Th	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U	Gross alpha	Gross beta
<b>Lancashire</b>									
Clifton Marsh	Borehole 6	2	<0.0018	<0.00070	0.035	<0.0023	<0.031	<0.31	2.0
Clifton Marsh	Borehole 19	2	<0.0015	<0.00085	0.082	<0.0040	0.073	<0.91	3.9
Clifton Marsh	Borehole 40	2	<0.0026	<0.00070	0.0080	<0.0014	<0.0067	<0.10	1.0
Clifton Marsh	Borehole 59	2	<0.0014	<0.00085	<0.0074	<0.0028	<0.0060	<0.24	2.1
<b>South Glamorgan</b>									
Lamby Way Tip <sup>b</sup>	Borehole 1A	2						<0.097	0.66

<sup>a</sup> As tritiated water

<sup>b</sup> The concentrations of <sup>125</sup>I and <sup>131</sup>I were <0.16 and <0.28 Bq l<sup>-1</sup> respectively

**Table 7.5. Concentrations of radionuclides in water near the East Northants Resource Management Facility landfill site, 2015**

Site reference	Mean radioactivity concentration <sup>a</sup> , Bq kg <sup>-1</sup>					
	<sup>3</sup> H	<sup>40</sup> K	<sup>137</sup> Cs	<sup>226</sup> Ra	<sup>228</sup> Th	<sup>230</sup> Th
K13A Groundwater borehole	<3.2	<4.6	<0.19	0.01	<0.0031	<0.0020
K15A Groundwater borehole	<3.0	<4.5	<0.19	0.01	<0.0027	<0.0018
K17 Northern perimeter Groundwater borehole	5.0	<4.6	<0.20	0.02	<0.0024	<0.0025
Horse Water spring		<2.7	<0.18			
Willow brook		<2.8	<0.19			

Site reference	Mean radioactivity concentration <sup>a</sup> , Bq kg <sup>-1</sup>					
	<sup>232</sup> Th	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U	Gross alpha	Gross beta
K13A Groundwater borehole	<0.00080	0.033	<0.0027	0.032	<0.14	0.24
K15A Groundwater borehole	<0.0012	0.016	<0.0018	0.015	<0.12	0.41
K17 Northern perimeter Groundwater borehole	<0.0012	0.038	<0.0020	0.030	<0.38	0.52
Horse Water spring					<0.054	0.31
Willow brook					<0.086	0.51

<sup>a</sup> Except for <sup>3</sup>H where units are Bq l<sup>-1</sup>

**Table 7.6. Concentrations of radionuclides in water and effluents near the Stoneyhill Landfill site, Aberdeenshire, 2015**

Sample location and type	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>	
		<sup>226</sup> Ra <sup>a</sup>	<sup>228</sup> Ra <sup>b</sup>
<b>Stoneyhill Landfill</b>			
Borehole 50 (Groundwater)	4	<0.09	<0.20
Borehole 78 (Groundwater)	4	<0.07	<0.44
Laeca Burn, adjacent to site (Surface water)	4	<0.08	<0.16
Laeca Burn, downstream of site (Surface water)	4	<0.07	<0.15
Laeca Burn, upstream of site (Surface water)	4	<0.10	<0.18
Leachate collection tank (Leachate)	4	<0.08	<0.19

<sup>a</sup> <sup>226</sup>Ra activity based on <sup>214</sup>Pb activity

<sup>b</sup> <sup>228</sup>Ra activity based on <sup>228</sup>Ac activity

**Table 7.7. Concentrations of naturally occurring radionuclides in the environment, 2015<sup>b</sup>**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>							
			<sup>210</sup> Po	<sup>210</sup> Pb	<sup>228</sup> Th	<sup>230</sup> Th	<sup>232</sup> Th	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U
<b>Phosphate processing, Whitehaven</b>										
Winkles	Parton	2	12	1.5						
Winkles	Nethertown	4	13	1.9	0.68	0.71	0.43	0.78	0.024	0.71
Mussels	Whitehaven	2	43	1.8						
Mussels	Ravenglass	2	42	1.5						
Prawns	Seascale	2	5.6	<0.16						
Crabs	Parton	2	24	0.22						
Crabs	Sellafield coastal area	2	21	0.19	0.16	0.013	0.0071	0.10	0.0037	0.087
Lobsters	Parton	2	19	0.16						
Lobsters	Sellafield coastal area	2	22	0.13						
Nephrops	Whitehaven	2	1.4	0.18	0.060	0.032	0.018	0.024	0.00081	0.023
Cod	Parton	2	0.64	0.074						
Cod	Whitehaven	2	1.1	0.16						
Plaice	Whitehaven	2	2.0	0.19	0.063	0.0010	0.00034	0.014	0.00060	0.013
Plaice	Drigg	2	2.1	0.15	0.060	<0.000084	<0.000084	0.016	0.00066	0.014
<b>Other samples</b>										
Winkles	South Gare (Hartlepool)	2	13	1.2						
Winkles	Middletons Sands	2	9.7							
Winkles	Kirkcudbright	1	5.6							
Mussels	Morecambe	2	38							
Mussels	Ribble Estuary	1			0.21	0.15	0.081			
Limpets	Kirkcudbright	1	16							
Crabs	Kirkcudbright	1	8.6							
Lobsters	Kirkcudbright	1	1.6							
Shrimps	Ribble Estuary	1			0.015	0.0072	0.0026			
Wildfowl	Ribble Estuary	1				0.011	0.0022			
Seaweed	Isle of Man	3						2.1	<0.21	1.8
Sediment	Kirkcudbright	2						6.4	<0.42	6.3
Sediment	Balcary Bay	1						7.4	<0.95	7.0

<sup>a</sup> Except for sediment where dry concentrations apply

<sup>b</sup> Data for artificial nuclides for some of these samples may be available in the relevant sections for nuclear sites

**Table 7.8. Discharges of gaseous radioactive wastes from non-nuclear establishments in the United Kingdom, 2015<sup>a</sup>**

	Discharges during 2015, Bq								
	Education (Universities and Colleges)			Hospitals			Other (Research, manufacturing and public sector)		
	England and Wales	Northern Ireland	Scotland	England and Wales	Northern Ireland	Scotland	England and Wales	Northern Ireland	Scotland
<sup>3</sup> H	1.0E+04		2.7E+09				3.1E+13		1.0E+09
<sup>14</sup> C	2.1E+07					5.6E+07	2.4E+13	1.2E+09	1.3E+10
<sup>18</sup> F	1.5E+12					7.1E+10	2.6E+11		
<sup>35</sup> S	1.3E+06						3.8E+08		
<sup>85</sup> Kr							1.3E+07		
<sup>99m</sup> Tc				3.6E+08			7.5E+06		
<sup>125</sup> I	5.5E+05			3.0E+07			3.3E+08		
<sup>129</sup> I							1.9E+04		
<sup>131</sup> I				4.3E+08			4.6E+08		
<sup>131m</sup> Xe				8.6E+07					
<sup>137</sup> Cs							4.9E+08		
<sup>222</sup> Rn							2.0E+09		
Uranium Alpha							2.0E+00		
Plutonium Alpha							3.4E+02		
<sup>241</sup> Am							6.4E+02		
Other Alpha									1.3E+03
Other Alpha particulate				5.8E+06			1.8E+11		
Other Beta/Gamma			9.3E+10		3.7E+11	5.6E+09			2.5E+10
Other Beta/Gamma Particulate	1.1E+12			2.2E+08			7.4E+12		

<sup>a</sup> Excludes nuclear power, defence and radiochemical manufacturing (Amersham and Cardiff) industries. Excludes discharges which are exempt from reporting. England and Wales discharge data refers to 2014

**Table 7.9. Discharges of liquid radioactive waste from non-nuclear establishments in the United Kingdom, 2015<sup>a</sup>**

	Discharges during 2015, Bq										
	Education (Universities and Colleges)			Hospitals			Other (Research, manufacturing and public sector)			Oil and gas (on- shore)	Oil and gas (off- shore)
	England and Wales	Northern Ireland	Scotland	England and Wales	Northern Ireland	Scotland	England and Wales	Northern Ireland	Scotland	Scotland	United Kingdom
<sup>3</sup> H	1.1E+10	3.4E+07	7.9E+09	2.7E+08	4.4E+08		3.2E+11		1.4E+10		
<sup>14</sup> C	1.2E+09	1.4E+04	3.0E+08	1.4E+07		7.3E+07	2.6E+11	1.0E+04	4.8E+10		
<sup>18</sup> F	7.7E+11			3.2E+12	2.8E+11	4.1E+11	2.6E+12		2.0E+03		
<sup>22</sup> Na			1.0E+06								
<sup>32</sup> P	7.0E+09	2.0E+06	1.7E+09	5.2E+09	1.6E+08	6.0E+08	5.7E+09		5.0E+08		
<sup>33</sup> P	5.1E+08		6.5E+09				2.4E+09		1.0E+10		
<sup>35</sup> S	1.1E+10		8.1E+09	1.8E+09			7.9E+09		1.4E+08		
<sup>51</sup> Cr	1.4E+09		2.5E+07	4.6E+10	6.1E+08	1.3E+09	1.6E+09				
<sup>57</sup> Co	2.0E+00				2.1E+04		8.5E+02		6.4E+05		
<sup>58</sup> Co					4.4E+04						
<sup>60</sup> Co	4.5E+01						1.8E+07				
<sup>67</sup> Ga	3.4E+08			8.0E+09		1.4E+08					
<sup>75</sup> Se	4.7E+07			2.9E+09	4.3E+07	5.3E+07	6.4E+07				
<sup>89</sup> Sr				2.6E+09			1.1E+08				
<sup>90</sup> Sr	1.4E+04						2.5E+07				
<sup>90</sup> Y				6.3E+11	1.0E+07	1.4E+09					
<sup>99</sup> Tc	2.0E+03						2.4E+06				
<sup>99m</sup> Tc	5.6E+10			5.5E+13	1.4E+12	4.6E+12	7.0E+11				
<sup>111</sup> In	1.2E+09			4.4E+11	9.4E+09	6.2E+10	6.1E+09				
<sup>125</sup> Sb							1.2E+03				
<sup>123</sup> I	3.0E+05		2.5E+07	1.1E+12	6.4E+10	8.9E+10	4.9E+10				
<sup>125</sup> I	4.8E+09	9.9E+07	1.4E+08	1.4E+09	1.6E+07	1.2E+08	5.3E+10		3.3E+07		
<sup>129</sup> I							1.7E+03				
<sup>131</sup> I	3.0E+05		2.7E+09	9.2E+12	4.9E+10	6.3E+11	2.6E+11				
<sup>134</sup> Cs							1.0E+07				
<sup>137</sup> Cs	1.4E+06		2.0E+04				9.6E+08				
<sup>153</sup> Sm	1.2E+07			1.6E+10	4.6E+09						
<sup>201</sup> Tl				2.9E+10		2.3E+10					
<sup>210</sup> Pb									3.0E+08		
<sup>210</sup> Po									3.0E+08		
<sup>226</sup> Ra									2.1E+09		
<sup>228</sup> Ra									1.5E+09		
<sup>230</sup> Th							6.0E+00				
<sup>232</sup> Th							1.8E+10		1.3E+06		
Uranium Alpha	1.2E+05		1.7E+06				3.3E+10				
<sup>237</sup> Np							1.5E+01				
<sup>241</sup> Pu							4.4E+04				
Plutonium Alpha			2.5E+06				6.8E+03				
<sup>241</sup> Am	5.2E+01						1.7E+04				
<sup>242</sup> Cm							7.0E+00				
Total Alpha	5.7E+06			3.9E+09	3.6E+07		6.1E+10				6.6E+09
Total Beta/Gamma (Excl Tritium)	8.5E+11			6.5E+13			3.8E+12				5.0E+09
Other Alpha									8.7E+03		
Other Alpha particulate				2.1E+09			4.8E+08				
Other Beta/ Gamma <sup>b</sup>	4.5E+10		1.3E+11	2.9E+12	5.1E+05	4.6E+10	1.3E+11		2.5E+07		
Other Beta/ Gamma particulate							5.1E+09				

<sup>a</sup> Excludes nuclear power, defence and radiochemical manufacturing (Amersham and Cardiff) industries. Excludes discharges which are exempt from reporting. England and Wales discharge data refers to 2014

<sup>b</sup> Excluding specific radionuclides

**Table 7.10. Monitoring in the Firth of Forth, River Clyde and Glasgow, 2015<sup>a</sup>**

Location	Material and selection	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>						
			<sup>3</sup> H	<sup>14</sup> C	<sup>32</sup> P	<sup>54</sup> Mn	<sup>90</sup> Sr	<sup>95</sup> Nb	<sup>99</sup> Tc
Between Finlaystone and Woodhall	Mussels	1		24	<6.1	<0.10		<0.38	0.87
Between Finlaystone and Woodhall	<i>Fucus vesiculosus</i>	1			12	<0.10		<0.44	8.6
Dalmuir Clydebank	Sediment	1		<15	<4.3	<0.10		<0.24	
Downstream of Dalmuir	Freshwater	4			<0.29	<0.10		<0.10	
River Clyde	Freshwater	4	<1.1				<0.0050		
Firth of Forth	Freshwater	4	<1.0				<0.017		
Daldowie	Sludge pellets	4			<79	<0.10		<0.84	

Location	Material and selection	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>					
			<sup>125</sup> Sb	<sup>131</sup> I	<sup>137</sup> Cs	<sup>155</sup> Eu	<sup>241</sup> Am	Gross beta
Between Finlaystone and Woodhall	Mussels	1	<0.24	<2.0	0.17	<0.23	<0.17	
Between Finlaystone and Woodhall	<i>Fucus vesiculosus</i>	1	<0.15	<25	0.20	<0.16	<0.12	
Dalmuir Clydebank	Sediment	1	<0.12	<10	6.3	0.89	<0.19	
Downstream of Dalmuir	Freshwater	4	<0.10	<0.93	<0.10	<0.10	<0.10	
River Clyde	Freshwater	4			<0.06			9.4
Firth of Forth	Freshwater	4			<0.10			13
Daldowie	Sludge pellets	4	<0.27	130	4.0	1.6	<0.39	

<sup>a</sup> Results are available for other radionuclides detected by gamma spectrometry, All such results are less than the limit of detection

<sup>b</sup> Except for water where units are Bq l<sup>-1</sup>, and sludge pellets and sediment where dry concentrations apply

## 8. Regional monitoring

Regional monitoring in areas remote from nuclear licensed sites has continued in 2015 (i) to establish long distance transport of radioactivity from UK and other nuclear licensed sites, (ii) to indicate general contamination of the food supply and the environment and (iii) to provide data under UK obligations under Article 36 of the Euratom Treaty and the OSPAR Convention.

The routine component parts of this programme are:

- Channel Islands, the Isle of Man and Northern Ireland
- Overseas sources
- General diet and milk
- Airborne particulates, rain, drinking water and groundwater
- Seawater and sediments

### 8.1 Channel Islands

Samples of marine environmental materials provided by the Channel Island states have been analysed for levels of radioactivity. The programme monitors the effects of radioactive discharges from the French reprocessing plant at La Hague and the power station at Flamanville. It also monitors any effects of historical disposals of radioactive waste in the Hurd Deep, a natural trough in the western English Channel. Fish and shellfish are monitored to determine exposure from the internal radiation pathway; sediment is analysed for external exposures. Seawater and seaweeds are sampled as environmental indicator materials and, in the latter case, because of their use as fertilisers. A review of marine radioactivity in the Channel Islands from 1990 to 2009 has been published (Hughes *et al.*, 2011).

Table 8.1 shows analysis results for 2015. There was evidence of routine releases from the nuclear industry in some samples (cobalt-60, technetium-99 and iodine-129). However, activity concentrations in fish and shellfish were low and similar to those in previous years. It is generally difficult to attribute the results to different sources, including fallout from weapon testing, due to the low levels detected. No evidence for significant releases of activity from the Hurd Deep site was found.

An assessment of the dose to a representative person who consumes large amounts of fish and shellfish was carried out. In 2015, the representative person was estimated to receive less than 0.005 mSv, which is less than 0.5 per cent of the dose limit for members of the public. The assessment included a contribution from external exposure. The concentrations of artificial radionuclides in the marine environment of the Channel Islands and the effects of

### Key points

- Monitoring in areas remote from nuclear licensed sites continued (i) to establish the effect of long distance transport of radioactivity from UK and other nuclear licensed sites, (ii) to detect any general contamination of the food supply and the environment and (iii) to provide data under UK obligations under Article 36 of the Euratom Treaty and the OSPAR Convention
- Sampling of marine life from the Channel Islands continued to monitor possible effects from French nuclear facilities discharging radioactivity into the English Channel. Doses were less than 1 per cent of the limit
- Monitoring in Northern Ireland and the Isle of Man showed low concentrations of man-made radionuclides from Sellafield and other UK nuclear facilities. Doses in Northern Ireland were less than 1 per cent of the dose limit
- Monitoring UK ports of entry continued for food from Japan, following the Fukushima Dai-ichi accident in 2011, and for other non-specific contamination. No significant contamination in food consignments was detected in 2015
- Samples from the UK food supply, air, rain and drinking water were analysed. Natural radionuclides from consumption of general diet and drinking water dominated the doses
- Surveys of seas around the UK supported international assessments for the OSPAR Treaty and showed the extent of tritium and caesium-137 contamination

discharges from local sources, therefore, continued to be of negligible radiological significance.

Milk and crop samples from the Channel ceased in 2014 following the completion of the risk-based review of the FSA's monitoring programme. Results up to 2013 are included in earlier RIFE reports and they show no significant effects of UK or other nuclear installations.

### 8.2 Isle of Man

The Environment Agency carries out a limited programme of radioactivity monitoring of the marine environment on the Isle of Man in collaboration with the Department of

Environment, Food and Agriculture (DEFA) (Table 8.3). The results complement the Isle of Man Government's own independent radiation monitoring programme and provide an indication of the far-field effects of current and historic discharges from Sellafield and other UK nuclear sites. Monitoring of food on the Isle of Man ceased in 2014 following completion of the risk-based review of FSA's programme.

The analyses carried out showed that levels of most radionuclides are reported as less than values of the method used. Positive values for technetium-99 and americium-241 were detected in seaweed but at the low levels consistent with those observed in recent years. The results demonstrate that there was no significant impact on the Isle of Man in 2015 from discharges to sea from mainland nuclear installations.

### 8.3 Northern Ireland

NIEA monitors the far-field effects of liquid discharges from Sellafield into the Irish Sea. The programme involves sampling fish, shellfish and indicator materials from a range of locations along the coastline (Figure 8.1). The external exposure pathway is studied by monitoring gamma dose rates over intertidal areas. The results are presented in Tables 8.4(a) and (b).

In 2015, the main effect of discharges from Sellafield was observed in concentrations of technetium-99 in shellfish and seaweed samples. These were similar to those in 2014 and recent years, reflecting the considerably decreased inputs to the Irish Sea (see also Section 2.3.3). Caesium-137 concentrations were low and similar to 2014 levels, and trace amounts of transuranic nuclides were detected. Observed concentrations were less than those found nearer to Sellafield and continued at the low levels seen in recent years (Figure 8.2). Further information on the trends in radioactivity in the marine environment of Northern Ireland is described in Ly *et al.*, (2015). The radiation dose rates over intertidal areas were similar to those in previous years.

A survey of consumption and occupancy in coastal regions of Northern Ireland (Smith *et al.*, 2002) established habits representative of people consuming large quantities of fish and shellfish. Based on the monitoring results from the marine environment in 2015, the dose to the most exposed person was 0.008 mSv, which is less than 1 per cent of the dose limit for members of the public.

Monitoring results for the terrestrial environment of Northern Ireland are included in the following parts of Section 8.

### 8.4 Overseas sources

Two overseas accidents have had direct implications for the UK: Chernobyl (1986) and Fukushima Dai-ichi (2011). Earlier RIFE reports have provided detailed results of monitoring by the environment agencies and FSA (Environment Agency, FSA, NIEA and SEPA, 2013).


For Chernobyl, the main sustained impact on the UK environment was in upland areas, where heavy rain fell in the days following the accident, but levels have now reduced substantially. Restrictions on the movement, selling and slaughter of sheep and related monitoring ceased in 2012 and sampling for freshwater fish in locations affected by Chernobyl ceased in 2014 following risk-based reviews by FSA (FSA, 2012a; b and 2013). The results of monitoring and estimated doses to consumers are available earlier RIFE reports (e.g. Environment Agency, FSA, NIEA and SEPA, 2013). The long term time-trend of caesium-137 concentrations in freshwater fish from Devoke Water (1986-2013) is given in Figure 8.3.

On 25 March 2011, the EC implemented controls (Regulation EU/297/2011) on the import of food and feed originating in or consigned from Japan following the Fukushima Dai-ichi accident (EC, 2011).

From 27 March 2011, imports of all feed and food originating in or consigned from Japan could only enter the UK through specific ports and airports where official controls will be carried out. Products of animal origin can only enter through Border Inspection Posts (BIPs) and products of non-animal origin can only enter through Designated Points of Entry (DPE).

The legislation has been updated in 2016 (Regulation EU/6/2016 (EC, 2016)). Certain measures apply to some feed and food originating in or consigned from 13 prefectures of Japan. The list of applicable feeds and foods from the prefectures can be found in Annex II to the legislation. Applicable feed and food products from these prefectures intended to be imported to the EU must be tested before leaving Japan and are subject to random testing in the EU. The exceptions are for certain personal consignments of feed and food. The main requirements of the regulation for imports of feed and food destined for the EU are that:

- Each consignment of food or feed listed in Annex II of the legislation, from the 13 prefectures must be accompanied by a declaration – signed by the Japanese authorities, attesting that the product complies with legislation in force in Japan, as regards the maximum levels for the sum of the radionuclides: caesium-134 and caesium-137. The declarations must be accompanied by the results of analysis.
- Importers are required to notify the BIP or DPE at the port of entry two working days before the arrival of each consignment of food and feed from Japan.



**Figure 8.1.** Monitoring locations in Northern Ireland, 2015

- Each consignment should be identified by a code on the declaration, analytical report, common entry documents, the sanitary certificate accompanying the consignment

Identity and physical checks, including laboratory analysis, on at 5 per cent of the consignments of food or feed will be undertaken by port officials where the product originates in or is consigned from Japan. Products that are found to exceed the maximum levels should not be placed on the market and are either safely disposed of, or returned to Japan. Further information is available on the FSA's website: [http://food.gov.uk/business-industry/imports/banned\\_restricted/japan](http://food.gov.uk/business-industry/imports/banned_restricted/japan).

A percentage of Japanese imports into the EU are monitored in the UK and this work continued in 2015. None of the imports to the UK have contained radioactivity exceeding the maximum permissible levels; no results were in excess of  $100 \text{ Bq kg}^{-1}$ . The doses received due to the imports were of negligible radiological significance.

Screening instruments are used at importation points of entry to the UK as a general check on possible

contamination from unknown sources. In 2015, the instruments were not triggered by a food consignment at any point of entry into the UK.

## 8.5 General diet

As part of the UK governments' general responsibility for food safety, concentrations of radioactivity are determined in regional diets. These data (and data on other dietary components in Sections 8.6 and 8.7) form the basis of the UK submission to the EC under Article 36 of the Euratom Treaty to allow comparison with data from other EU member states (for example, Joint Research Centre, 2009). Concentrations of radioactivity in the general diet are reported to the EC by the FSA (for England, Northern Ireland and Wales), and by SEPA (for Scotland).

In 2015, the concentrations found in a survey of radioactivity in canteen meals and mixed diet collected across the UK were very low or typical of natural sources (Table 8.5). Where comparisons can be made, similar values were observed in 2014.




Figure 8.2. Concentrations of americium-241 and caesium-137 in coastal sediments in Northern Ireland, 2002-2015




Figure 8.3. Caesium-137 concentrations in freshwater fish from Devoke Water, Cumbria 1986-2013

## 8.6 Milk

The programme of milk sampling across dairies in the UK continued in 2015. Its aim is to collect and analyse samples on a monthly basis for their radionuclide content. This programme provides useful information with which to compare data from farms close to nuclear licensed sites and other establishments that may enhance concentrations above background levels. Milk data is reported by FSA (for England, Northern Ireland and Wales) and SEPA (for Scotland) as part of the UK submission to the EC under Article 36 of the Euratom Treaty (for example, Joint Research Centre, 2009).

The results are summarised in Table 8.2. The majority of measurements, where comparable, are similar to those in previous years. Carbon-14 concentrations are very close to the expected background concentration in milk (see Appendix 1, Annex 4). Tritium results were again below detection limits. The mean concentration of strontium-90 detected was about 0.04 Bq l<sup>-1</sup>. In the past, the concentrations of radiocaesium in milk were highest from those regions that received the greatest amounts of Chernobyl fallout. However, the concentrations are now very low and it is not possible to distinguish this trend.

Radiation dose from consuming milk at average rates was assessed for various age groups. In 2015, the maximum dose was to a 1 year-old infant. For the range of radionuclides analysed, the dose was less than 0.005 mSv. Previous surveys (for example, FSA and SEPA, 2002) have shown that if a full range of nuclides are analysed and assessed, the dose is dominated by naturally occurring lead-210 and polonium-210, whereas man-made radionuclides contribute less than 10 per cent.

## 8.7 Crops

The programme of monitoring naturally occurring and man-made radionuclides in crops (in England, Wales and the Channel Islands) as a check on general food contamination (remote from nuclear sites) ceased in 2014 following a risk-based review by FSA. In the last year of sampling in 2013, there were no significant trends in concentrations of radionuclides in crops (Environment Agency, FSA, NIEA, NRW and SEPA, 2014). Tritium concentrations are reported as less than values in most samples. Carbon-14 was generally detected at levels close to those expected to occur through natural processes. Levels of other naturally occurring radionuclides varied from region to region and plutonium isotopes and americium-241 were detected at trace levels in some samples.

## 8.8 Airborne particulate, rain, freshwater and groundwater

Radioactivity in rainwater and air was monitored at several UK locations as part of the programme of background sampling managed by the Environment Agency and SEPA. These data are reported on behalf of BEIS, NIEA and the Scottish and Welsh Governments, as part of the UK submission to the EC under Article 36 of the Euratom Treaty (for example, Joint Research Centre, 2009). The results are given in Table 8.6. The routine programme comprised two components (i) regular sampling and analysis on a quarterly basis and (ii) supplementary analysis on an *ad hoc* basis by gamma-ray spectrometry. Caesium-137 concentrations are reported as less than values. These levels in air, typical of recent years, remain less than 0.01 per cent of those observed in 1986, the year of the Chernobyl reactor accident.

Concentrations of beryllium-7, a naturally occurring radionuclide formed by cosmic ray reactions in the upper atmosphere, were detected at similar levels at all sampling locations. Peak air concentrations of this radionuclide tend to occur during spring and early summer as a result of seasonal variations in the mixing of stratospheric and tropospheric air (Environment Agency, 2002a). Tritium concentrations in rainwater were similar to those in recent years. Activity concentrations in air and rainwater were very low and do not currently merit radiological assessment.

Sampling and analysis of freshwater from drinking water sources throughout the UK continued in 2015 (Figure 8.4). These water data are reported by the Environment Agency (for England and Wales), NIEA (for Northern Ireland) and SEPA (for Scotland) as part of the UK submission to the EC under Article 36 of the Euratom Treaty (for example, Joint Research Centre, 2009). Sampling was designed to represent the main drinking water sources, namely reservoirs, rivers and groundwater boreholes. Most of the water samples were representative of natural waters before treatment and supply to the public water system. The results in Tables 8.7, 8.8 and 8.9 show that concentrations of tritium were all substantially below the investigation level for drinking water of 100 Bq l<sup>-1</sup> in the European Directive 2013/51. The highest value in Scotland was found at Gullielands Burn, which is near to the Chapelcross nuclear licensed site. Concentrations of gross alpha and gross beta were generally below the investigation levels for drinking water in the European Directive 2013/51, however the mean gross alpha concentration in Derbyshire at 0.11 Bq l<sup>-1</sup>, although similar to previous years, was just above the corresponding investigation level of 0.1 Bq l<sup>-1</sup>.

The mean annual dose from consuming drinking water in the UK was assessed as 0.033 mSv in 2015 (Table 8.10). The highest annual dose was estimated to be 0.051 mSv for drinking water from Matlock in Derbyshire. The

estimated doses were dominated by naturally occurring radionuclides and are similar to those in recent years. The annual dose from artificial radionuclides in drinking water was less than 0.001 mSv.

Separately, in 2015, SEPA took a series of groundwater samples from across Scotland and the results are displayed in Table 8.11. All samples contained levels below or near the LoD and are generally consistent with those in recent years. A single positive measurement of tritium in groundwater from Annan could conceivably be due to the operation of the nearby Chapelcross nuclear site. Regardless of the source, at the levels detected there are no radiological protection implications.

## 8.9 Seawater surveys

The UK Governments are committed to preventing pollution of the marine environment from ionising radiation, with the ultimate aim of reducing concentrations in the environment to near background values for naturally-occurring radioactive substances, and close to zero for artificial radioactive substances (DECC, Department of the Environment, Northern Ireland, The Scottish Government and Welsh Assembly Government, 2009). Therefore, a programme of surveillance into the distribution of key radionuclides is maintained using research vessels and other means of sampling.

The seawater surveys reported here also support international studies concerned with the quality status of coastal seas (for example, OSPAR, 2010b). A fourth periodic evaluation of progress towards internationally agreed objectives has been published by OSPAR (OSPAR, 2016b). The programme of radiological surveillance work provides the source data and, therefore, the means to monitor and make an assessment of progress in line with the UK's commitments towards OSPAR's 1998 Strategy for Radioactive Substances target for 2020 (part of the North-east Atlantic Environment Strategy adopted by OSPAR for the period 2010-2020). The surveys also provide information that can be used to distinguish different sources of man-made radioactivity (for example, Kershaw and Baxter, 1995) and to derive dispersion factors for nuclear licensed sites (Baxter and Camplin, 1994). In addition, the distribution of radioactivity in seawater around the British Isles is a significant factor in determining the variation in individual exposures at coastal sites, as seafood is a major contribution to food chain doses. Evidence to help gauge progress towards achieving the Government's vision for radionuclides and other hazardous substances is set out in a report (Defra, 2010).

The research vessel programme on radionuclide distribution currently comprises annual surveys of the Bristol Channel/western English Channel and biennial surveys of the Irish Sea and the North Sea. The results obtained in 2015 are given in Figures 8.5 – 8.9.






Figure 8.4. Drinking water sampling locations, 2015

A seawater survey of the Irish Sea was carried out in 2015 (Figure 8.5). As in previous surveys, a band of higher concentrations of caesium-137 was observed along the coast to the north and south of Sellafield, with levels generally decreasing with distance from the coast. Caesium-137 concentrations were reasonably uniform in a large part of the Irish Sea. The 2015 survey recorded concentrations of up to 0.06 Bq l<sup>-1</sup> in the eastern Irish Sea (0.09 Bq l<sup>-1</sup> in 2013); elsewhere concentrations were generally below 0.02 Bq l<sup>-1</sup>. Overall, concentrations were similar to those reported in the previous Irish Sea survey in 2013 (Environment Agency, FSA, NIEA, NRW and SEPA, 2014). Caesium-137 concentrations in the Irish Sea were only a small percentage of those prevailing in the late 1970s (typically up to 30 Bq l<sup>-1</sup>, Baxter *et al.*, 1992), when discharges were substantially higher.


The predominant source of caesium-137 to the Irish Sea is now considered to be remobilisation into the water column from activity associated with seabed sediment. This was re-confirmed in a recent study (Hunt *et al.*, 2013). Discharges from Sellafield have decreased substantially since the commissioning of the SIXEP waste treatment process in the mid-1980s, and this has been reflected in a near exponential decrease in shoreline seawater concentrations at St Bees (Figure 8.10). In more recent years, the rate of decline of caesium-137 concentrations with time has been decreasing at St Bees. Longer time series showing peak concentrations in the Irish Sea and, with an associated time-lag, the North Sea are also shown in Figure 8.10.




**Figure 8.5.** Concentrations (Bq l<sup>-1</sup>) of caesium-137 in filtered surface water from the Irish Sea, September 2015




**Figure 8.7.** Concentrations (Bq l<sup>-1</sup>) of tritium in surface water from the Irish Sea, September 2015



**Figure 8.6.** Concentrations (Bq l<sup>-1</sup>) of caesium-137 in filtered surface water from the English Channel, February-March 2015



**Figure 8.8.** Concentrations (Bq l<sup>-1</sup>) of tritium in surface water from the Bristol Channel, September 2015



**Figure 8.9.** Concentrations ( $\text{Bq l}^{-1}$ ) of tritium in surface water from the English Channel, September 2015

In 2014, very low concentrations of caesium-137 (up to  $0.006 \text{ Bq l}^{-1}$ ) were found throughout most of the North Sea survey area (Environment Agency, FSA, FSS, NIEA, NRW and SEPA, 2015), and these were only slightly above those observed for global fallout levels in surface seawaters ( $0.0001\text{--}0.0028 \text{ Bq l}^{-1}$ , Povinec *et al.*, 2005).

Concentrations of caesium-137 (up to  $0.003 \text{ Bq l}^{-1}$ ) in the western English Channel (Figure 8.6) were not distinguishable from the background levels of global fallout (within experimental error) in 2015. Activity concentrations near the Channel Islands were similar in 2015 (compared to those in 2014), and lower than concentrations in both the Irish and North Seas.

A full assessment of long-term trends of caesium-137 in surface waters of Northern European seas is provided elsewhere (Povinec *et al.*, 2003).

Tritium concentrations in Irish Sea seawater in 2015 are shown in Figure 8.7. As expected, these are higher than those observed in the North Sea in 2014 (Environment Agency, FSA, FSS, NIEA, NRW and SEPA, 2015) due to the influence of discharges from Sellafield and other nuclear licensed sites. Tritium concentrations to the south and west of the Isle of Man, including along the coastline of Ireland, were mostly below (or close to) the LoD.


In the Bristol Channel, the combined effect of tritium discharges from Cardiff, Berkeley, Oldbury and Hinkley

Point is shown in Figure 8.8. Overall, the general level of tritium concentrations in the Bristol Channel was very low in 2015. Tritium concentrations in samples taken close to these installations were generally similar to those in the 2013 survey, but were lower than levels in the North-east Irish Sea (Figure 8.7). Tritium concentrations in the western English Channel were all below the LoD (Figure 8.9).

Technetium-99 concentrations in seawater are now decreasing following the substantial increases observed from 1994 to their most recent peak in 2003. The results of research cruises to study this radionuclide have been published by Leonard *et al.* (1997a, b; 2004) and McCubbin *et al.* (2002; 2008). Trends in plutonium and americium concentrations in seawater of the Irish Sea have been considered by Leonard *et al.* (1999). Full reviews of the quality status of the north Atlantic and a periodic evaluation of progress towards internationally agreed targets have been published by OSPAR (2000b; 2009; 2010b). A research study, commissioned by the FSA, determined the depth distributions of technetium-99 concentrations in sea-bed cores to produce an estimate of the total inventory residing in the sub-tidal sediments of the Irish Sea (Jenkinson *et al.*, 2013). The study concluded that the inventory of technetium-99 was estimated to have been of the order of 30 TBq (or approximately 2 per cent of the total cumulative Sellafield discharge), with approximately 8 TBq present in surface material and thereby potentially most susceptible to re-dissolution or re-suspension.

Shoreline sampling was also carried out around the UK, as part of routine site and regional monitoring programmes. Much of the shoreline sampling was directed at establishing whether the impacts of discharges from individual sites are detectable. Where appropriate, these are reported in the relevant sections of this report, and the analysis results are collated in Table 8.12. Most radionuclides are below limits of detection, and tritium and caesium-137 levels remote from site discharge points are consistent with those in Figures 8.5 – 8.9.

In 2015, SEPA took a series of marine sediment samples from across Scotland and the results are displayed in Table 8.13. Various radionuclides were detected. The results are generally consistent with those to be expected from measurements at nuclear licensed sites in this report (see, for example, Section 2), and are similar to those sampled in 2014. Overall, the results support the concept of a reducing trend in concentration with distance from the Sellafield site, albeit confounded by natural variability due to sediment type.



**Figure 8.10.** Concentration of caesium-137 in the Irish Sea, North Sea and in shoreline seawater close to Sellafield at St. Bees (Note different scales used for activity concentrations)

**Table 8.1. Concentrations of radionuclides in seafood and the environment near the Channel Islands, 2015**

Location	Material	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>							
			Organic <sup>3</sup> H	<sup>3</sup> H	<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>99</sup> Tc	<sup>106</sup> Ru	<sup>137</sup> Cs
<b>Guernsey</b>										
	Mackerel	1				<0.05			<0.45	0.09
	Bass	1				<0.06			<0.44	0.24
	Crabs	1				<0.09			<0.81	<0.08
	Lobsters	1				<0.08			<0.66	<0.07
	Limpets	1				<0.13			<0.99	<0.11
	Ormers	1				<0.13			<1.0	<0.10
	Scallops	1				<0.11			<0.92	<0.09
St. Sampson's	Mud and sand	1				<0.35			<2.8	<0.34
Harbour	Seawater	4								0.002
<b>Jersey</b>										
	Crabs	1				<0.05			<0.39	<0.05
	Spiny spider crabs	1				<0.06			<0.47	<0.05
	Lobsters	1				<0.07		0.33	<0.53	<0.06
La Rocque	Oysters	1				<0.02			<0.17	<0.02
La Rozel	Limpets	1				<0.03			<0.26	<0.03
Plemont Bay	<i>Porphyra</i>	2				<0.08			<0.58	<0.07
La Rozel	<i>Fucus vesiculosus</i>	4				<0.07	<0.062	2.2	<0.50	<0.06
Gorey	<i>Ascophyllum nodosum</i>	4				<0.07			<0.49	<0.06
<b>Alderney<sup>b</sup></b>										
	Crabs	2	<25	<25	47	<0.06		<0.65	<0.46	<0.05
	Spiny spider crabs	1				0.22			<0.48	<0.05
	Lobsters	1				<0.06			<0.49	<0.06
	Toothed winkles	1	<25	<25	33	<0.23	<0.069		<1.8	<0.19
	<i>Fucus vesiculosus</i>	2								
Quenard Point	<i>Fucus serratus</i>	4				<0.09	<0.040	0.62	<0.64	<0.07
Quenard Point	<i>Laminaria digitata</i>	4				<0.08			<0.63	<0.07
Little Crabbe	Sand	1				<0.29			<2.7	1.8
Harbour	Seawater	4		<3.8						0.002

**Table 8.1. continued**

Location	Material	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>a</sup> , Bq kg <sup>-1</sup>						Gross beta
			<sup>155</sup> Eu	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm+ <sup>244</sup> Cm	
<b>Guernsey</b>									
	Mackerel	1	<0.12	0.000045	0.00021	0.00083	*	0.000016	140
	Bass	1	<0.09	0.000011	0.000019	0.000017	*	*	100
	Crabs	1	<0.20						98
	Lobsters	1	<0.18			<0.21			110
	Limpets	1	<0.17			<0.10			93
	Ormers	1	<0.18			<0.09			91
	Scallops	1	<0.22	0.00093	0.0035	0.0012	*	0.00013	62
St. Sampson's Harbour	Mud and sand	1	<0.73	0.027	0.11	0.14	*	0.011	400
<b>Jersey</b>									
	Crabs	1	<0.08	0.00028	0.00077	0.0020	*	0.00019	77
	Spiny spider crabs	1	<0.09			<0.06			87
	Lobsters	1	<0.11	0.00010	0.00052	0.0055	*	0.00046	120
La Rocque	Oysters	1	<0.04	0.0011	0.0034	0.0046	*	0.00031	72
La Rozel	Limpets	1	<0.06	0.0015	0.0041	0.0071	*	0.00037	83
Plemont Bay	<i>Porphyra</i>	2	<0.13			<0.11			120
La Rozel	<i>Fucus vesiculosus</i>	4	<0.13	0.0025	0.0080	0.0055	*	0.00033	170
Gorey	<i>Ascophyllum nodosum</i>	4	<0.14			<0.16			200
<b>Alderney<sup>b</sup></b>									
	Crabs	2	<0.12	0.00029	0.00085	0.0019	*	0.00020	66
	Spiny spider crabs	1	<0.10	0.0012	0.0030	0.0045	0.000014	0.00052	91
	Lobsters	1	<0.11	0.00039	0.0013	0.010	0.000029	0.0010	94
	Toothed winkles	1	<0.36	0.0064	0.022	0.033	0.00079	0.0027	93
	<i>Fucus vesiculosus</i>	2							
Quenard Point	<i>Fucus serratus</i>	4	<0.17	0.0020	0.0073	0.0045	0.000028	0.00042	190
Quenard Point	<i>Laminaria digitata</i>	4	<0.13			<0.11			190
Little Crabbe Harbour	Sand	1	<0.58			0.63			700

\* Not detected by the method used

<sup>a</sup> Except for seawater where units are Bq l<sup>-1</sup>, and for sediment where dry concentrations apply

<sup>b</sup> The concentration of <sup>129</sup>I based on two observations in *Fucus vesiculosus* is 0.93 Bq kg<sup>-1</sup>

**Table 8.2. Concentrations of radionuclides in milk remote from nuclear sites, 2015**

Location	Selection <sup>a</sup>	No. of farms/ dairies <sup>b</sup>	Mean radioactivity concentration, Bq l <sup>-1</sup>			
			<sup>3</sup> H	<sup>14</sup> C	<sup>90</sup> Sr	Total Cs
<b>Milk</b>						
Co. Antrim		1		19	<0.021	<0.07
Co. Armagh		1			<0.027	<0.05
Ceredigion		1			<0.033	<0.05
Cheshire		1		9.2	<0.024	<0.05
Clwyd		1		15	<0.026	<0.05
Cornwall		1		18	<0.024	<0.05
Devon		1		16	<0.026	<0.04
Dorset		1		20	<0.023	<0.04
Co. Down		1			<0.025	<0.05
Dumfriesshire		1	<5.0	<19	<0.10	<0.05
Essex		1		9.5	<0.023	<0.04
Co. Fermanagh		1			<0.023	<0.05
Gloucestershire		1		9.9	<0.025	<0.05
Gwynedd		1		23	<0.031	<0.05
Hampshire		1		6.5	<0.027	<0.05
Humberside		1		17	<0.025	<0.04
Kent		1		20	<0.025	<0.04
Lanarkshire		1	<5.0	<17	0.018	<0.02
Lancashire		1		15	<0.026	<0.05
Leicestershire		1		20	<0.024	<0.04
Middlesex		1		15	<0.021	<0.05
Midlothian		1	<5.0	17	<0.10	<0.05
Nairnshire		1	<5.0	<18	<0.10	<0.05
Norfolk		1		16	<0.022	<0.04
North Yorkshire		1		24	<0.023	<0.05
Renfrewshire		1	<5.0	<16	<0.11	<0.05
Co. Tyrone		2		20	<0.023	<0.05
		max			<0.024	
<b>Mean Values</b>						
England				15	<0.024	<0.05
Northern Ireland				20	<0.024	<0.05
Wales				19	<0.030	<0.05
Scotland			<5.0	<17	<0.086	<0.04
United Kingdom			<5.0	17	<0.036	<0.05

<sup>a</sup> Data are arithmetic means unless stated as 'max'. 'Max' data are selected to be maxima  
If no 'max' value is given the mean value is the most appropriate for dose assessments

<sup>b</sup> The number of farms or dairies from which milk is sampled. The number of analyses is greater than this and depends on the bulking regime

**Table 8.3. Concentrations of radionuclides in food and the environment from the Isle of Man, 2015**

Material	No. of sampling observations	Mean radioactivity concentration (fresh), Bq kg <sup>-1</sup>						
		<sup>60</sup> Co	<sup>95</sup> Nb	<sup>95</sup> Zr	<sup>99</sup> Tc	<sup>106</sup> Ru	<sup>125</sup> Sb	<sup>134</sup> Cs
<b>Aquatic samples</b>								
Seaweed	3	<0.84	<0.59	<1.1	58	<5.0	<2.9	<0.78
<b>Material</b>								
Material	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>						
		<sup>137</sup> Cs	<sup>144</sup> Ce	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U	<sup>241</sup> Am	
Seaweed	3	<0.62	<2.1	<0.21	1.8	<0.68	2.1	

**Table 8.4(a). Concentrations of radionuclides in seafood and the environment in Northern Ireland, 2015<sup>a</sup>**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>					
			<sup>14</sup> C	<sup>60</sup> Co	<sup>99</sup> Tc	<sup>125</sup> Sb	<sup>134</sup> Cs	<sup>137</sup> Cs
Cod	Kilkeel	4	30	<0.05		<0.12	<0.05	0.74
Plaice	Kilkeel	4		<0.05		<0.12	<0.05	0.50
Haddock	Kilkeel	4		<0.06		<0.12	<0.06	0.30
Herring	Ardglass	2		<0.09		<0.20	<0.10	<0.20
Lesser spotted dogfish	North coast	4		<0.15		<0.32	<0.14	0.60
Skates / rays	Kilkeel	4		<0.10		<0.21	<0.10	0.87
Crabs	Kilkeel	4		<0.06		<0.15	<0.07	<0.10
Lobsters	Ballycastle	2		<0.11	2.6	<0.25	<0.11	0.16
Lobsters	Kilkeel	4		<0.05	8.1	<0.12	<0.06	0.15
<i>Nephrops</i>	Kilkeel	4		<0.12	6.0	<0.25	<0.12	0.66
Winkles	Minerstown	4		<0.05		<0.12	<0.05	<0.15
Mussels	Carlingford Lough	2		<0.11	1.8	<0.28	<0.12	<0.34
Scallops	Co. Down	2		<0.07		<0.16	<0.07	0.11
<i>Ascophyllum nodosum</i>	Carlingford Lough	3		<0.09	77	<0.18	<0.09	0.23
<i>Ascophyllum nodosum</i>	Ardglass	3		<0.06	42	<0.12	<0.06	0.18
<i>Fucus</i> spp.	Portrush	4		<0.03		<0.08	<0.04	<0.04
<i>Fucus vesiculosus</i>	Carlingford Lough	1		<0.05		<0.11	<0.06	0.28
<i>Fucus vesiculosus</i>	Ardglass	1		<0.04		<0.09	<0.05	0.43
<i>Rhodomenia</i> spp.	Portaferry	4		<0.07	0.26	<0.15	<0.07	0.41
Mud	Carlingford Lough	2		<0.62		<1.7	<0.81	43
Mud	Dundrum Bay	1		<0.38		<0.85	<0.49	4.1
Mud	Strangford Lough-Nicky's point	2		<0.34		<1.0	<0.47	14
Mud and sand	Ballymacormick	1		<0.39		<0.99	<0.55	7.0
Mud and sand	Dundrum Bay	1		<0.42		<1.1	<0.58	3.5
Mud and sand	Oldmill Bay	1		<0.23		<0.66	<0.32	5.9
Mud and shell	Carrichue	1		<0.23		<0.65	<0.29	1.4
Sand	Ballymacormick	1		<0.21		<0.69	<0.32	4.8
Sand	Oldmill Bay	1		<0.18		<0.65	<0.26	4.6
Sand	Portrush	2		<0.19		<0.56	<0.26	0.48
Shell and sand	Carrichue	1		<0.20		<0.67	<0.30	1.3
Seawater	North of Larne	12			0.0028		*	0.006

**Table 8.4(a). continued**

Material	Location	No. of sampling observations	Mean radioactivity concentration (fresh) <sup>b</sup> , Bq kg <sup>-1</sup>					
			<sup>155</sup> Eu	<sup>238</sup> Pu	<sup>239</sup> Pu + <sup>240</sup> Pu	<sup>241</sup> Am	<sup>242</sup> Cm	<sup>243</sup> Cm + <sup>244</sup> Cm
Cod	Kilkeel	4	<0.11			<0.08		
Plaice	Kilkeel	4	<0.11			<0.13		
Haddock	Kilkeel	4	<0.10			<0.09		
Herring	Ardglass	2	<0.15			<0.09		
Lesser spotted dogfish	North coast	4	<0.23			<0.15		
Skates / rays	Kilkeel	4	<0.17			<0.13		
Crabs	Kilkeel	4	<0.13			<0.15		
Lobsters	Ballycastle	2	<0.16			0.20		
Lobsters	Kilkeel	4	<0.10			<0.09		
<i>Nephrops</i>	Kilkeel	4	<0.18	0.0083	0.051	0.23	0.00023	0.00015
Winkles	Minerstown	4	<0.11	0.022	0.15	0.11	*	*
Mussels	Carlingford Lough	2	<0.23			<0.25		
Scallops	Co. Down	2	<0.15			<0.18		
<i>Ascophyllum nodosum</i>	Carlingford Lough	3	<0.20			<0.24		
<i>Ascophyllum nodosum</i>	Ardglass	3	<0.12			<0.12		
<i>Fucus</i> spp.	Portrush	4	<0.07			<0.08		
<i>Fucus vesiculosus</i>	Carlingford Lough	1	0.31			<0.06		
<i>Fucus vesiculosus</i>	Ardglass	1	<0.08			0.24		
<i>Rhodomenia</i> spp.	Portaferry	4	<0.12	0.064	0.38	0.70	*	*
Mud	Carlingford Lough	2	<1.5	1.8	11	8.7	*	*
Mud	Dundrum Bay	1	<0.84			1.2		
Mud	Strangford Lough-Nicky's point	2	<1.2			4.0		
Mud and sand	Ballymacormick	1	<0.93			8.1		
Mud and sand	Dundrum Bay	1	<1.1			2.0		
Mud and sand	Oldmill Bay	1	<0.65			13		
Mud and shell	Carrichue	1	<0.82	0.080	0.57	1.0	*	*
Sand	Ballymacormick	1	<0.88			1.2		
Sand	Oldmill Bay	1	<0.75			2.8		
Sand	Portrush	2	<0.65			<0.66		
Shell and sand	Carrichue	1	<0.83			<1.0		

\* Not detected by the method used

<sup>a</sup> All measurements are made on behalf of the Northern Ireland Environment Agency

<sup>b</sup> Except for seawater where units are Bq l<sup>-1</sup>, and for sediment where dry concentrations apply

**Table 8.4(b). Monitoring of radiation dose rates in Northern Ireland, 2015<sup>a</sup>**

Location	Ground type	No. of sampling observations	Mean gamma dose rate in air at 1m, $\mu\text{Gy h}^{-1}$
Lisahally	Mud	1	0.071
Donnybrewer	Shingle	1	0.051
Carrichue	Mud	1	0.052
Bellerena	Mud	1	0.055
Benone	Sand	1	0.058
Castlerock	Sand	1	0.060
Portstewart	Sand	1	0.050
Portrush, Blue Pool	Sand	1	0.058
Portrush, White Rocks	Sand	1	0.063
Portballintrae	Sand	1	0.053
Giant's Causeway	Sand	1	0.056
Ballycastle	Sand	1	0.056
Cushendun	Sand	1	0.055
Cushendall	Sand and stones	1	0.058
Red Bay	Sand	1	0.062
Carnlough	Sand	1	0.053
Glenarm	Sand	1	0.055
Half Way House	Sand	1	0.053
Ballygally	Sand	1	0.053
Drains Bay	Sand	1	0.052
Larne	Sand	1	0.057
Whitehead	Sand	1	0.056
Carrickfergus	Sand	1	0.058
Jordanstown	Sand	1	0.060
Helen's Bay	Sand	1	0.059
Groomsport	Sand	1	0.059
Millisle	Sand	1	0.072
Ballywalter	Sand	1	0.061
Ballyhalbert	Sand	1	0.057
Cloghy	Sand	1	0.058
Portaferry	Shingle and stones	1	0.084
Kircubbin	Sand	1	0.063
Greyabbey	Sand	1	0.076
Ards Maltings	Mud	1	0.071
Island Hill	Mud	1	0.070
Nicky's Point	Mud	1	0.086
Strangford	Shingle and stones	1	0.078
Kilclief	Sand	1	0.080
Ardglass	Mud	1	0.068
Killough	Mud	1	0.079
Ringmore Point	Sand	1	0.078
Tyrella	Sand	1	0.070
Dundrum	Sand	1	0.085
Newcastle	Sand	1	0.078
Annalong	Sand	1	0.080
Cranfield Bay	Sand	1	0.070
Mill Bay	Sand	1	0.090
Greencastle	Sand	1	0.080
Rostrevor	Sand	1	0.11
Narrow Water	Mud	1	0.10

<sup>a</sup> All measurements are made on behalf of the Northern Ireland Environment Agency

**Table 8.5. Concentrations of radionuclides in diet, 2015<sup>a</sup>**

Region	No. of sampling observations	Mean radioactivity concentration (fresh), Bq kg <sup>-1</sup>			
		<sup>14</sup> C	<sup>40</sup> K	<sup>90</sup> Sr	<sup>137</sup> Cs
<b>Canteen meals</b>					
England	8		85	<0.028	<0.07
Northern Ireland	5		85	<0.027	<0.05
Scotland	12	49	82	0.024	<0.021
Wales	5		86	<0.030	<0.06
<hr/>					
Region	No. of sampling observations	Mean radioactivity concentration (fresh), Bq kg <sup>-1</sup>			
		<sup>14</sup> C	<sup>40</sup> K	<sup>90</sup> Sr	<sup>137</sup> Cs
<b>Mixed diet in Scotland</b>					
<b>Dumfriesshire</b>					
Dumfries	4		85	<0.10	<0.05
<b>East Lothian</b>					
North Berwick	4		86	<0.10	<0.05
<b>Renfrewshire</b>					
Paisley	4		92	<0.11	<0.04
<b>Ross-shire</b>					
Dingwall	4		86	<0.10	<0.05

<sup>a</sup> Results are available for other artificial nuclides detected by gamma spectrometry  
All such results were less than the limit of detection

**Table 8.6. Concentrations of radionuclides in rainwater and air, 2015**

Location	Sample	Number of sampling observations	Mean radioactivity concentration <sup>a</sup>					
			<sup>3</sup> H	<sup>7</sup> Be	<sup>7</sup> Be <sup>d</sup>	<sup>90</sup> Sr	<sup>137</sup> Cs	<sup>137</sup> Cs <sup>d</sup>
<b>Ceredigion</b>								
Aberporth	Rainwater	4	<1.1	<0.83			<0.012	
	Air	4		0.0032			<5.9 10 <sup>-7</sup>	
<b>Co. Down</b>								
Conlig	Rainwater	4		<0.92			<0.0085	
	Air	4		0.0024			<5.6 10 <sup>-7</sup>	
<b>Dumfries and Galloway</b>								
Eskdalemuir	Rainwater	12	<1.1	<0.72		<0.0081	<0.010	
	Air	12		0.0011			<1.0 10 <sup>-5</sup>	
<b>North Lanarkshire</b>								
Holytown	Rainwater	8	<1.0	<0.15		<0.0099	<0.010	
	Air	12		0.0011			<1.0 10 <sup>-5</sup>	
<b>North Yorkshire</b>								
Dishforth	Rainwater	4		<0.88			<0.0085	
	Air	4		0.0017			<7.2 10 <sup>-7</sup>	
<b>Oxfordshire</b>								
Chilton	Rainwater	4		<0.84	0.98	<0.00084 <sup>b</sup>	<0.016	<0.0020
	Air	12			0.0020			<3.8 10 <sup>-7</sup>
<b>Shetland</b>								
Lerwick	Rainwater	12	<1.0	<0.95		<0.0064	<0.010	
	Air	12		0.0013			<1.0 10 <sup>-5</sup>	
<b>Suffolk</b>								
Orfordness	Rainwater	4	<1.2	<1.1			<0.0094	
	Air	4		0.0029			<5.1 10 <sup>-7</sup>	

Location	Sample	Number of sampling observations	Mean radioactivity concentration <sup>a</sup>				
			<sup>238</sup> Pu <sup>c</sup>	<sup>239</sup> Pu+ <sup>240</sup> Pu <sup>c</sup>	<sup>241</sup> Am <sup>c</sup>	Gross alpha	Gross beta
<b>Ceredigion</b>							
Aberporth	Rainwater	4	<3.7 10 <sup>-6</sup>	4.8 10 <sup>-6</sup>	<1.1 10 <sup>-5</sup>		
	Air	4	<1.1 10 <sup>-9</sup>	1.2 10 <sup>-9</sup>	2.4 10 <sup>-9</sup>		
<b>Dumfries and Galloway</b>							
Eskdalemuir	Air	12					<0.00020
<b>North Lanarkshire</b>							
Holytown	Air	12					<0.00020
<b>Oxfordshire</b>							
Chilton	Rainwater	4				0.026 <sup>d</sup>	0.064 <sup>d</sup>
<b>Shetland</b>							
Lerwick	Air	12					<0.00020

<sup>a</sup> Bq l<sup>-1</sup> for rainwater and Bq kg<sup>-1</sup> for air. 1.2 kg air occupies 1m<sup>3</sup> at standard temperature and pressure

<sup>b</sup> Bulked from 4 quarterly samples

<sup>c</sup> Separate annual sample for rain, annual bulked sample for air

<sup>d</sup> Bulked from 12 monthly samples

**Table 8.7. Concentrations of radionuclides in sources of drinking water in Scotland, 2015**

Area	Location	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>				
			<sup>3</sup> H	<sup>90</sup> Sr	<sup>137</sup> Cs	Gross alpha	Gross beta
Angus	Loch Lee	4	<1.0	<0.016	<0.01	<0.0095	<0.096
Argyll and Bute	Auchengaich	1	<1.0		<0.01	<0.010	0.030
Argyll and Bute	Helensburgh Reservoir	1	<1.0		<0.01	<0.010	0.017
Argyll and Bute	Loch Ascog	1	<1.0		<0.01	<0.010	0.098
Argyll and Bute	Loch Eck	1	<1.0		<0.01	<0.010	0.034
Argyll and Bute	Lochan Ghlas Laoigh	1	<1.0		<0.01	<0.010	<0.031
Argyll and Bute	Loch Finlas	1	<1.0		<0.01	<0.010	0.029
Clackmannanshire	Gartmorn Dam	1	<1.0		<0.01	<0.010	0.15
Dumfries and Galloway	Black Esk	1	<1.0		<0.01	<0.010	0.81
Dumfries and Galloway	Gullielands Burn	1	16		<0.01	<0.010	0.19
Dumfries and Galloway	Purdomstone	1	1.8		<0.01	<0.010	0.040
Dumfries and Galloway	Winterhope	1	1.6		<0.01	0.012	0.044
East Lothian	Hopes Reservoir	1	<1.0		<0.01	<0.010	0.036
East Lothian	Thorters Reservoir	1	<1.0		<0.01	0.010	0.041
East Lothian	Whiteadder	1	<1.0		<0.01	<0.010	0.033
East Lothian	Thornton Loch Burn	1	<1.0		<0.01	0.016	0.10
Fife	Holl Reservoir	1	<1.0		<0.01	<0.010	<0.049
Highland	Loch Baligill	1	<1.0		<0.01	<0.010	0.050
Highland	Loch Calder	1	<1.0		<0.01	<0.010	0.060
Highland	Loch Glass	4	<1.0	<0.0052	<0.01	<0.0088	<0.044
Highland	Loch Shurrerey	1	<1.0		<0.01	0.015	0.070
North Ayrshire	Camphill	1	<1.0		<0.01	<0.010	0.030
North Ayrshire	Knockendon Reservoir	1	<1.0		<0.01	<0.010	0.051
North Ayrshire	Munnoch Reservoir	1	<1.0		<0.01	<0.010	0.052
North Ayrshire	Outerwards	1	<1.0		<0.01	<0.010	0.056
Orkney Islands	Heldale Water	1	<1.0		<0.01	<0.010	0.060
Perth and Kinross	Castlehill Reservoir	1	<1.0		<0.01	<0.010	<0.013
Scottish Borders	Knowesdean	4	<1.0	<0.0050	<0.01	<0.0087	0.029
Stirling	Loch Katrine	12	<1.0	<0.0026	<0.001	<0.0077	0.061
West Dunbartonshire	Loch Lomond (Ross Priory)	1	<1.0		<0.01	<0.010	0.033
West Lothian	Morton No 2 Reservoir	1	<1.0		<0.01	<0.010	0.33

**Table 8.8. Concentrations of radionuclides in sources of drinking water in England and Wales, 2015**

Location	Sample source	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>								
			<sup>3</sup> H	<sup>40</sup> K	<sup>90</sup> Sr	<sup>125</sup> I	<sup>137</sup> Cs	Gross alpha	Gross beta <sup>1</sup>	Gross beta <sup>2</sup>	
<b>England</b>											
Buckinghamshire	Bourne End, Groundwater	1	<4.0	<0.050	<0.0010		<0.0019				
Cambridgeshire	Grafham Water	4	<4.0	0.31	<0.0022		<0.0010	0.033	0.48	0.31	
Cornwall	River Fowey	4	<4.0	0.066	0.0025	<0.0020	<0.0010	<0.027	0.092	0.062	
County Durham	Honey Hill Water Treatment Works, Consett	4	<4.0	0.079	0.0039		<0.0023	0.097	0.23	0.15	
County Durham	River Tees, Darlington	4	<4.0	<0.041	0.0033	<0.0020	<0.0010	<0.019	0.067	0.050	
Cumbria	Ennerdale Lake	4	<4.0	<0.016	0.0041		<0.0011	<0.019	<0.049	<0.050	
Cumbria	Haweswater Reservoir	4	<4.0	<0.011	0.0027		<0.0010	<0.020	<0.050	<0.050	
Derbyshire	Arnfield Water Treatment Plant	4	<4.0	<0.044	0.0024		<0.0012	<0.020	<0.050	<0.050	
Derbyshire	Matlock, Groundwater <sup>a</sup>	4	<4.0	<0.051	<0.0018		<0.0014	0.11	0.12	0.077	
Devon	River Exe, Exeter	2	<4.0	<0.053	0.0034	<0.0054	<0.0019	<0.029	0.092	0.065	
Devon	Roadford Reservoir, Broadwoodwidge	4	<4.0	0.071	0.0025		<0.0010	<0.020	0.086	0.056	
Greater London	River Lee, Chingford	4	<4.0	0.28	<0.0012	<0.0019	<0.0010	<0.021	0.43	0.27	
Hampshire	River Avon, Christchurch	3	<4.0	0.083	<0.0012	<0.0020	<0.0011	<0.022	0.15	0.10	
Humberside	Littlecoates, Groundwater	4	<4.0	0.12	<0.0010		<0.0010	0.021	0.17	0.11	
Kent	Chatham, Deep Groundwater	1	<4.0	0.050	0.0014		<0.0010	0.016	0.058	<0.050	
Kent	Denge, Shallow Groundwater	4	<4.0	0.091	0.0064		<0.0031	<0.024	0.15	0.095	
Lancashire	Corn Close, Groundwater	4	<4.0	0.091	<0.0085		<0.0083	0.021	0.16	0.10	
Norfolk	River Drove, Stoke Ferry	4	<4.0	0.12	<0.0022	<0.0020	<0.0010	0.033	0.19	0.12	
Northumberland	Kielder Reservoir	4	<4.0	<0.10	0.0033		<0.0011	<0.019	<0.19	<0.13	
Oxfordshire	River Thames, Oxford	4	<4.0	0.16	<0.0014	<0.0020	<0.0010	0.029	0.23	0.15	
Somerset	Ashford Reservoir, Bridgwater	4	<4.0	0.076	<0.0014		<0.0010	<0.025	0.11	0.074	
Somerset	Chew Valley Lake Reservoir, Bristol	4	<4.0	0.11	0.0025		<0.0010	0.022	0.19	0.12	
Surrey	River Thames, Chertsey	2	<4.0	0.19	0.0025	<0.0020	<0.0010	0.045	0.30	0.18	
Surrey	River Thames, Walton	4	<4.0	0.24	<0.0013	<0.0020	<0.0010	<0.021	0.32	0.20	
<b>Wales</b>											
Gwynedd	Cwm Ystradlyn Treatment Works	4	<4.0	<0.021	0.0036		<0.0014	<0.019	<0.055	<0.049	
Mid-Glamorgan	Llwyn-on Reservoir	4	<4.0	<0.048	0.0034		<0.0022	0.016	0.050	<0.050	
Powys	Elan Valley Reservoir	4	<4.0	<0.026	0.0037		<0.0010	<0.020	<0.050	<0.050	

<sup>1</sup> Using <sup>137</sup>Cs standard

<sup>2</sup> Using <sup>40</sup>K standard

<sup>a</sup> The concentrations of <sup>210</sup>Po, <sup>226</sup>Ra, <sup>234</sup>U, <sup>235</sup>U and <sup>238</sup>U were <0.020, <0.011, 0.042, <0.010 and 0.023 Bq l<sup>-1</sup> respectively

**Table 8.9. Concentrations of radionuclides in sources of drinking water in Northern Ireland, 2015**

Area	Location	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>								Gross alpha	Gross beta
			<sup>3</sup> H	<sup>90</sup> Sr	<sup>137</sup> Cs	<sup>210</sup> Po	<sup>226</sup> Ra	<sup>234</sup> U	<sup>235</sup> U	<sup>238</sup> U		
Co. Londonderry	R Faughan	4	<1.6	<0.0025	<0.05	<0.010	<0.010	<0.010	<0.010	<0.010	<0.010	0.035
Co. Antrim	Lough Neagh	4	<1.1	<0.0028	<0.05	<0.010	<0.010	<0.010	<0.010	<0.010	<0.010	0.11
Co. Down	Silent Valley	4	<1.1	<0.0015	<0.05	<0.010	<0.010	<0.010	<0.010	<0.010	<0.010	0.036

**Table 8.10. Doses from radionuclides in drinking water, 2015<sup>a</sup>**

Region	Mean exposure, mSv per year			Maximum exposure, mSv per year	
	Man-made radionuclides <sup>b,c</sup>	Naturally occurring radionuclides <sup>b</sup>	All radionuclides	Location	All radionuclides
England	<0.001	0.051	0.051	Matlock, Groundwater, Derbyshire	0.051
Wales <sup>d</sup>	<0.001			Elan Valley Reservoir, Powys	<0.001 <sup>d</sup>
Northern Ireland	<0.001	0.026	0.027	R Faughan, Co. Londonderry	0.027
Scotland <sup>d</sup>	<0.001			Gullielands Burn, Dumfries and Galloway	<0.001 <sup>d</sup>
UK	<0.001	0.033	0.033	Matlock, Groundwater, Derbyshire	0.051

<sup>a</sup> Assessments of dose are based on some concentration results at limits of detection.

Exposures due to potassium-40 content of water are not included here because they do not vary according to the potassium-40 content of water.

Levels of potassium are homeostatically controlled

<sup>b</sup> Average of the doses to the most exposed age group at each location

<sup>c</sup> Including tritium

<sup>d</sup> Analysis of naturally occurring radionuclides was not undertaken

**Table 8.11. Analysis of groundwater – background survey in Scotland, 2015**

Location	Sample source	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>			
			<sup>3</sup> H	<sup>137</sup> Cs	Gross alpha	Gross beta
Scottish Borders	Selkirk	1	1.3	<0.10	<0.010	0.021
East Lothian	Dunbar	1	<1.0	<0.10	<0.010	0.037
West Lothian	Livingston	1	<1.0	<0.10	<0.025	0.72
Fife	Kingsbarns	1	<1.0	<0.10	0.077	0.21
Fife	Falkland	1	<1.0	<0.10	<0.010	0.082
Angus	Brechin	1	<1.0	<0.10	<0.010	0.047
Angus	Montrose	1	<1.0	<0.10	<0.010	0.047
Aberdeenshire	Stonhaven	1	<1.0	<0.10	<0.010	0.053
Moray	Fochabers	1	<1.0	<0.10	<0.010	0.21
Dumfries and Galloway	Stranraer	1	<1.0	<0.10	0.035	0.089
Dumfries and Galloway	Dumfries	1	<1.0	<0.10	1.1	0.14
Dumfries and Galloway	Annan	1	2.4	<0.10	0.025	0.13

**Table 8.12. Concentrations of radionuclides in seawater, 2015**

Location	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>						
		<sup>3</sup> H	<sup>14</sup> C	<sup>60</sup> Co	<sup>90</sup> Sr	<sup>99</sup> Tc	<sup>106</sup> Ru	<sup>110m</sup> Ag
Dounreay (Brims Ness)	2 <sup>S</sup>	<1.0		<0.10				
Rosyth	2 <sup>S</sup>	<1.0		<0.10			<0.31	<0.10
Tornessa	2 <sup>S</sup>	<1.8		<0.10			<0.41	<0.10
Hartlepool (North Gare)	2	<3.6		<0.25			<1.7	<0.29
Sizewell	2	<3.2		<0.31			<2.3	<0.39
Bradwell (Beach pipeline)	2	<3.5		<0.26			<1.9	<0.30
Bradwell (Marina)	2			<0.24			<1.8	<0.28
Bradwell (Steeple)	2			<0.29			<1.9	<0.32
Bradwell (Maylandsea Bay)	2			<0.27			<2.0	<0.32
Bradwell (Blackwater)	2			<0.22			<1.7	<0.26
Bradwell (Osea Causeway)	2			<0.31			<2.4	<0.36
Bradwell (Maldon)	2	<3.3		<0.38			<2.6	<0.42
Bradwell (Heybridge)	2			<0.22			<1.6	<0.26
Bradwell (Strood Channel)	2	<3.2		<0.32			<2.3	<0.36
Bradwell (Tollesbury boatyard)	2			<0.26			<1.9	<0.28
Bradwell (Tollesbury saltwater pool)	2			<0.30			<2.3	<0.36
Bradwell (Pyefleet)	2	<3.3		<0.34			<2.3	<0.39
Bradwell (Rowhedge)	2			<0.27			<1.9	<0.31
Bradwell (Alresford Creek)	2			<0.22			<1.7	<0.26
Bradwell (Brightlingsea Bateman's Tower)	2			<0.30			<2.3	<0.35
Bradwell (Brightlingsea saltwater pool)	2			<0.23			<1.7	<0.27
Bradwell (St Osyth)	2			<0.27			<1.9	<0.30
Dungeness south	2	<3.0		<0.22			<1.8	<0.30
Winfrith (Lulworth Cove)	1			<0.24			<2.0	<0.32
Alderney	5 <sup>F</sup>	<3.8						
Devonport (Millbrook Lake)	2	<3.5	<8.8	<0.28				
Devonport (Tor Point South)	2	<3.2	<11	<0.23				
Hinkley	2			<0.31	<0.040		<2.3	<0.37
Berkeley and Oldbury	2			<0.23			<1.8	<0.27
Cardiff (Orchard Ledges) <sup>b</sup>	2	<9.6	<5.4	<0.29				
Wylfa (Cemaes Bay)	2	<3.3		<0.20			<1.8	<0.27
Wylfa (Cemlyn Bay West)	2			<0.28			<2.1	<0.36
Heysham (inlet)	2	31		<0.28			<2.2	<0.35
Seascale (Particulate)	2			<0.03	<0.0083		<0.20	<0.03
Seascale (Filtrate)	2			<0.21	<0.030	<0.21	<1.6	<0.23
St. Bees (Particulate)	2			<0.03	<0.020		<0.20	<0.03
St. Bees (Filtrate)	2	<7.0		<0.22	<0.018	<0.19	<1.8	<0.26
Seafield	2 <sup>S</sup>	2.7		<0.10			<0.26	<0.10
Southernness	2 <sup>S</sup>	2.5		<0.10			<0.24	<0.10
Auchencairn	2 <sup>S</sup>	<2.4		<0.10			<0.26	<0.10
Port Patrick	2 <sup>S</sup>	<1.0		<0.10			<0.41	<0.10
Hunterston <sup>c</sup>	2 <sup>S</sup>	4.2		<0.10			<0.22	<0.10
North of Larne	12 <sup>N</sup>					0.0028		
Faslane (Carnban)	2 <sup>S</sup>	<1.0		<0.10			<0.18	<0.10

**Table 8.12. continued**

Location	No. of sampling observations	Mean radioactivity concentration, Bq l <sup>-1</sup>					
		<sup>134</sup> Cs	<sup>137</sup> Cs	<sup>144</sup> Ce	<sup>241</sup> Am	Gross alpha	Gross beta
Dounreay (Sandside Bay)	2 <sup>S</sup>	<0.10	<0.10	<0.13	<0.10		
Dounreay (Brims Ness)	2 <sup>S</sup>	<0.10	<0.10	<0.12	<0.10		
Rosyth	2 <sup>S</sup>	<0.10	<0.10	<0.20	<0.10		
Tornessa	2 <sup>S</sup>	<0.10	<0.10	<0.23	<0.10		
Hartlepool (North Gare)	2	<0.24	<0.20	<0.88	<0.28	<4.5	12
Sizewell	2	<0.32	<0.25	<1.1	<0.27	<3.2	13
Bradwell (Beach pipeline)	2	<0.26	<0.21	<0.87	<0.28	<3.9	12
Bradwell (Marina)	2	<0.24	<0.20	<0.89	<0.29	<4.0	13
Bradwell (Steeple)	2	<0.30	<0.23	<0.91	<0.29	<4.2	18
Bradwell (Maylandsea Bay)	2	<0.29	<0.22	<0.91	<0.29	<4.2	14
Bradwell (Blackwater)	2	<0.23	<0.19	<0.84	<0.28	<4.3	14
Bradwell (Osea Causeway)	2	<0.32	<0.27	<0.93	<0.30	<4.1	16
Bradwell (Maldon)	2	<0.37	<0.30	<0.98	<0.31	<1.5	6.0
Bradwell (Heybridge)	2	<0.23	<0.19	<0.84	<0.28	<3.6	15
Bradwell (Strood Channel)	2	<0.31	<0.26	<0.98	<0.30	<3.9	16
Bradwell (Tollesbury boatyard)	2	<0.27	<0.22	<0.86	<0.28	<3.8	15
Bradwell (Tollesbury saltwater pool)	2	<0.31	<0.26	<0.94	<0.31	<4.8	15
Bradwell (Pyefleet)	2	<0.33	<0.26	<0.92	<0.28	<3.6	14
Bradwell (Rowhedge)	2	<0.28	<0.22	<0.88	<0.27	<4.6	11
Bradwell (Alresford Creek)	2	<0.23	<0.19	<0.86	<0.29	<4.1	13
Bradwell (Brightlingsea Bateman's Tower)	2	<0.31	<0.26	<0.99	<0.31	<3.2	11
Bradwell (Brightlingsea saltwater pool)	2	<0.23	<0.19	<0.84	<0.28	<4.1	14
Bradwell (St Osyth)	2	<0.27	<0.22	<0.88	<0.28	<3.7	13
Dungeness south	2	<0.24	<0.20	<0.99	<0.27	<3.3	14
Winfrith (Lulworth Cove)	1	<0.27	<0.20	<1.1	<0.30	<2.5	14
Alderney	5 <sup>F</sup>	*	0.002				
Guernsey	4 <sup>F</sup>	*	0.002				
Hinkley	2	<0.31	<0.24	<0.96	<0.27	<3.1	13
Berkeley and Oldbury	2	<0.23	<0.19	<0.91	<0.28	<1.6	6.5
Cardiff (Orchard Ledges) <sup>b</sup>	2		<0.24				
Wylfa (Cemaes Bay)	2	<0.22	<0.18	<0.91	<0.26	<3.0	9.4
Wylfa (Cemlyn Bay West)	2	<0.29	<0.22	<1.0	<0.28	<3.7	14
Heysham (inlet)	2	<0.28	<0.23	<1.1	<0.29	<2.9	9.4
Seascale (Particulate)	2	<0.03	<0.02	<0.07	0.06	0.14	0.061
Seascale (Filtrate)	2	<0.20	<0.18	<0.79	<0.27	<3.3	9.7
St. Bees (Particulate)	2	<0.03	<0.04	<0.07	<0.09	0.22	0.076
St. Bees (Filtrate)	2	<0.24	<0.20	<0.67	<0.23	<3.6	<12
Seafield	2 <sup>S</sup>	<0.10	<0.10	<0.14	<0.10		
Southernness	2 <sup>S</sup>	<0.10	<0.11	<0.16	<0.10		
Auchencairn	2 <sup>S</sup>	<0.10	<0.10	<0.17	<0.10		
Port Patrick	2 <sup>S</sup>	<0.10	<0.10	<0.23	<0.10		
Hunterston <sup>c</sup>	2 <sup>S</sup>	<0.10	<0.10	<0.15	<0.10		
North of Larne	12 <sup>N</sup>	*	0.006				
Faslane (Carnban)	2 <sup>S</sup>	<0.10	<0.10	<0.12	<0.10		

\* Not detected by the method used

<sup>a</sup> The concentration of <sup>35</sup>S was <0.53 Bq l<sup>-1</sup>

<sup>b</sup> The concentrations of <sup>3</sup>H as tritiated water and <sup>125</sup>I were <4.1 Bq l<sup>-1</sup> and <0.32 Bq l<sup>-1</sup> respectively

<sup>c</sup> The concentration of <sup>35</sup>S was <0.81 Bq l<sup>-1</sup>

Results are made on behalf of the Environment Agency unless indicated otherwise

<sup>F</sup> Measurements labelled "F" are made on behalf of the Food Standards Agency and the Channel Islands States

<sup>N</sup> Measurements labelled "N" are made on behalf of the Northern Ireland Environment Agency

<sup>S</sup> Measurements labelled "S" are made on behalf of the Scottish Environment Protection Agency

**Table 8.13. Concentrations of radionuclides in marine sediments – background survey in Scotland, 2015<sup>a</sup>**

Location	Sample source	No. of sampling observations	Mean radioactivity concentration, Bq kg <sup>-1</sup> (dry) <sup>b</sup>									
			<sup>3</sup> H	<sup>60</sup> Co	<sup>110</sup> Ag	<sup>125</sup> Sb	<sup>137</sup> Cs	<sup>155</sup> Eu	<sup>241</sup> Am	Gross alpha	Gross beta	
<b>Marine Sediments</b>												
Firth of Clyde	NW Cloch Point	1	<5.0	<0.22	<0.37	<0.74	40	<0.72	6.6	130	860	
Firth of Clyde	Lunderston Bay	1	<5.0	<0.22	<0.40	<0.80	52	<0.89	7.9	200	890	
Firth of Clyde	Wemyss Point	1	<5.0	<0.24	<0.40	<0.76	39	1.3	7.2	110	910	
Firth of Clyde	East of Toward	1	<5.0	<0.10	<0.12	<0.23	20	0.67	2.9	100	960	
Firth of Clyde	East of Strone Point	1	<5.0	0.30	<0.59	<1.1	83	<1.1	16	83	730	
Firth of Clyde	East of Brodick	1	<5.0	<0.20	<0.16	<0.63	45	1.2	18	82	830	
Firth of Clyde	7km off Whiting Bay	1	<5.0	<0.24	<0.45	<0.87	73	1.7	26	93	960	
Firth of Clyde	5km SW of Lady Isle	1	<5.0	<0.26	<0.48	<0.92	51	1.8	19	94	1000	
Firth of Clyde	Middle at offshore	1	<5.0	<0.30	<0.55	<1.1	64	2.5	35	89	910	
Firth of Clyde	Middle at offshore south	1	<5.0	<0.23	<0.40	<0.75	48	1.8	21	100	860	
<b>Seawater</b>												
Firth of Clyde	NW Cloch Point	1	1.7	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	
Firth of Clyde	Lunderston Bay	1	1.6	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	
Firth of Clyde	Wemyss Point	1	1.3	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	
Firth of Clyde	East of Toward	1	1.8	<0.10	<0.01	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	
Firth of Clyde	East of Strone Point	1	1.4	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	
Firth of Clyde	East of Brodick	1	1.6	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	
Firth of Clyde	7km off Whiting Bay	1	1.8	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	
Firth of Clyde	5km SW of Lady Isle	1	1.7	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	
Firth of Clyde	Middle at offshore	1	1.3	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	
Firth of Clyde	Middle at offshore south	1	1.6	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	<0.10	

<sup>a</sup> Results are available for other radionuclides detected by gamma spectrometry, All such results are less than the limit of detection

<sup>b</sup> Except for seawater where units are Bq l<sup>-1</sup>



## 9. References

**(Includes references from Appendix 1: CD supplement; sorted in order of first author and then date)**

- Allott, R., 2005. Assessment of compliance with the public dose limit. Principles for the assessment of total retrospective public doses. NDAWG/2/2005. Environment Agency, FSA, HPA, NII, Chilton.
- Baxter, A.J., Camplin, W.C. and Steele, A.K., 1992. Radiocaesium in the seas of northern Europe: 1975 – 79. Fish. Res. Data Rep., MAFF Direct. Fish. Res., Lowestoft, (28): 1 – 166.
- Baxter, A.J. and Camplin, W.C., 1994. The use of caesium-137 to measure dispersion from discharge pipelines at nuclear sites in the UK. Proc. Instn. Civ. Engrs. Wat., Marit. And Energy, (106): 281 – 288.
- Baxter, J.M., Boyd, I.L., Cox, M., Donald, A.E., Malcolm, S.J., Miles, H., Miller, B. and Moffat, C.F., (Editors), 2011. Scotland's Marine Atlas: Information for the national marine plan. Marine Scotland, Edinburgh. pp. 191
- <http://www.scotland.gov.uk/Topics/marine/science/atlas>
- BNFL, 2002. Discharges and monitoring of the environment in the UK. Annual Report 2001. BNFL, Warrington.
- Brenk, H.D., Onishi, Y., Simmonds, J.R. and Subbaratnam, T., (unpublished). A practical methodology for the assessment of individual and collective radiation doses from radionuclides in the environment. International Atomic Energy Authority draft working document no. 1987 – 05 – 06, Vienna.
- Brown, J., Hammond, D., Wilding, D., Wilkins, B.T., and Gow, C., 2009. Transfer of radioactivity from seaweed to terrestrial foods and potential radiation exposures to members of the public: 2009. HPA-RPD-059. HPA, Chilton.
- Brown, J. and Etherington, G., 2011. Health Risks from Radioactive Objects on Beaches in the Vicinity of the Sellafield Site. HPA-CRCE-018, April 2011, HPA, Chilton.
- Byrom, J., Robinson, C.A., Simmonds, J.R., Walters, C.B. and Taylor, R.R., 1995. Food consumption rates for use in generalised radiological dose assessments. J. Rad. Prot., 15(4): 335 – 342.
- Camplin, W.C., Grzechnik, M. and Smedley, C.A., 2005. Methods for assessment of total dose in the Radioactivity in Food and the Environment report. NDAWG/3/2005. Environment Agency, FSA, HPA, NII, Chilton.
- Camplin, W.C. and Jenkinson, S., 2007. Use of measurements in determining retrospective dose assessments in RIFE. (NDAWG/11/03) 2007. Environment Agency, FSA, HPA, NII, Chilton.
- CEC, 1989. Council regulation (Euratom) No 3954/87 laying down the maximum permitted levels of radioactive contamination of foodstuffs and feeding stuffs following a nuclear accident or any other case of radiological emergency. OJ 11(L371), amended by Council Regulation 2218/89 OJ 1(L211).
- CEC, 1992. Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. OJ L206: 7 – 50.
- CEC, 1996. Council Directive 96/29/Euratom of 13 May 1996 laying down basic safety standards for the protection of the health of workers and the general public against the dangers arising from ionising radiation. OJ 39(L159): 1 – 114.
- CEC, 1998. Council Directive 98/83/EC of 3 November 1998 on the quality of water intended for human consumption. OJ L330: 32 – 54.
- CEC, 1999a. Council Directive 1999/2/EC of the European Parliament and of the Council of 22 February 1999 on the approximation of the laws of Member States concerning foods and food ingredients treated with ionising radiation. OJ L 66, 13.3.99, 16.
- CEC, 1999b. Council Directive 1999/3/EC of the European Parliament and of the Council of 22 February 1999 on the establishment of a Community list of foods and food ingredients treated with ionising radiation. OJ L 66, 13.3.99, 24.
- CEC, 2000a. Commission recommendation on the application of Article 36 of the Euratom Treaty concerning the monitoring of the concentrations of radioactivity in the environment for the purpose of assessing the exposure of the population as a whole. OJ 27th July 2000. 2000/473/ Euratom.

- CEC, 2008. Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for Community action in the field of marine environmental policy (Marine Strategy Framework Directive). OJ L164.
- CEC, 2009. Directive 2009/147/EC of the European Parliament and of the Council of 130 November 2009 on the conservation of wild birds. OJ L 20, 26.1.2010, 7 – 25.
- Clyne, F.J., Gough, C., Edgar, A. and Smedley, C.A., 2008. Radiological Habits Survey: Sellafield Beach Occupancy, 2007. Project C3015. RL 02/08. Cefas, Lowestoft
- Clyne, F.J., Gough, C., Edgar, A., Garrod, C.J. and Elliott, J., 2010a. Radiological Habits Survey: Sellafield Beach Occupancy, 2009. Project C3635. RL 01/10. Cefas, Lowestoft
- Clyne, F.J., Garrod, C.J. and Elliott, J., 2010b. Radiological Habits Survey: Amersham 2009. Project C2848. RL 04/10. Cefas, Lowestoft.
- Clyne, F.J., Garrod, C.J., Rumney, P., Hughes, L.M. and Ly, V.E., 2011a. Radiological Habits Survey: Hinkley Point, 2010. Project C2848. RL 01/11. Cefas, Lowestoft.
- Clyne, F.J., Garrod, C.J., Ly, V.E. and Rumney, P., 2011b. Radiological Habits Survey: Dungeness, 2010. Project C2848. RL 11/11. Cefas, Lowestoft.
- Clyne, F.J., Garrod, C.J., Ly, V.E., and Rumney, P., 2012. Radiological Habits Survey: Devonport, 2011. RL 01/12. Cefas, Lowestoft.
- Clyne, F.J., Garrod, C.J., Rumney, P. and Ly, V.E., 2013a. Radiological Habits Survey: Chapelcross, 2010. RL 17/13. Cefas, Lowestoft.
- Clyne, F.J., Garrod, C.J., and Ly, V.E., 2013b. Radiological Habits Survey: Torness, 2011. RL 19/13. Cefas, Lowestoft.
- Clyne, F.J., Garrod, C.J., Ly, V.E., Dewar, A. and Papworth, G.P., 2013c. Radiological Habits Survey: Low Level Waste Repository, 2012. RL 02a/13. Cefas, Lowestoft.
- Clyne, F.J., Garrod, C.J. and Papworth, G.P., 2014. Radiological Habits Survey: Sellafield, 2012. RL 02/14. Cefas, Lowestoft.
- Clyne, F.J., Garrod, C.J. and Papworth, G.P., 2015. Radiological Habits Survey: Berkeley and Oldbury, 2014. RL 02/15. Cefas, Lowestoft. Clyne, F.J., Garrod, C.J. and Ly, V.E. 2016. Radiological Habits Survey: Bradwell, 2015. RL 02/16. Cefas, Lowestoft.
- Clyne, F.J., Garrod, C.J. and Dewar, A. 2016. Radiological Habits Survey: Harwell, 2015. RL 03/16. Cefas, Lowestoft.
- Clyne, F.J. and Garrod, C.J. 2016. Radiological Habits Survey: Sellafield Review, 2015. RL 04/16. Cefas, Lowestoft.
- Codex Alimentarius Commission, 2006. Codex Alimentarius Commission Report, Twenty-ninth session 3-7 July 2006. AL INORM 06/29/41. Codex, Rome.
- CoRWM, 2014. Tenth annual report 2013-14. CoRWM, London.
- Cooper, J. R., 2008. Review of risks from tritium – report of the AGIR – November 2007. Letter dated 17<sup>th</sup> April 2008. HPA, Chilton.
- Copeland Borough Council *et al.*, 1999. The radiological implications of contaminated feral pigeons found at Sellafield and Seascale. CBC, DH, Environment Agency, HSE, MAFF and NRPB, Penrith.
- Dale, P., Robertson, I and Toner, M., 2008. Radioactive particles in dose assessments. J. Environ. Rad., 99: 1589-1595.
- Darby, S, 2016. Personal communication. ONR, Bootle.
- DECC, 2009. Summary of responses to the consultation on the draft UK Strategy for radioactive discharges 2006-2030. DECC, London.
- DECC, Department of the Environment Northern Ireland, the Scottish Government and Welsh Assembly Government, 2009. UK Strategy for Radioactive Discharges. DECC, London. <https://www.gov.uk/government/publications/implementing-geological-disposal>
- DECC and Welsh Assembly Government, 2009. Statutory Guidance to the Environment Agency concerning the regulation of radioactive discharges into the environment. DECC and Welsh Assembly Government, London and Cardiff.
- DECC, 2010. The United Kingdom's fifth national report on compliance with the convention on nuclear safety obligations. DECC, London.
- DECC, 2011. The United Kingdom's fourth national report on compliance with the obligations of the joint convention on the safety of spent fuel management and on the safety of radioactive waste management. DECC, London.
- DECC, 2012. Environmental Protection Act 1990: Part iiA. Contaminated Land. Statutory Guidance. DECC, London.

- DECC, Scottish Government, Welsh Government and the Northern Ireland Department of the Environment, 2012. Strategy for the Management of Solid Low Level Radioactive Waste from the Non-Nuclear Industry; Part 1 – Anthropogenic Radionuclides, DECC, London.
- DECC, 2014a. Implementing Geological Disposal. DECC, London.
- DECC, 2014b. The United Kingdom's fifth national report on compliance with the obligations of the Joint Convention on the safety of spent fuel management and the safety of radioactive waste management. DECC, London.
- DECC, 2016a. Geological Disposal Facility (GDF) for higher-activity radioactive waste. 4 March 2016. DECC, London.
- DECC, 2016b. Appraisal of Sustainability Scoping Report. DECC, London.
- DECC and NDA, 2014. 2013 UK Radioactive Waste Inventory, February 2014, DECC and NDA, London and Moor Row, Cumbria.
- DECC, Scottish Government, Welsh Government and the Northern Ireland Department of the Environment, 2014. Strategy for the management of naturally occurring radioactive material (NORM) waste in the United Kingdom. The Scottish Government, Edinburgh.
- DECC, Scottish Government, Welsh Government and the Department of the Environment Northern Ireland, 2015. UK Strategy for the management of solid low level waste for the nuclear industry. DECC, London.
- Defra, 2002. UK strategy for radioactive discharges 2001 – 2020. Defra, London.
- Defra, Scottish Executive and Welsh Assembly Government, 2002. Safeguarding our seas. A strategy for the conservation and sustainable development of our marine environment. Defra, London.
- Defra, 2004. Contribution of aerial radioactive discharges to radionuclide concentrations in the marine environment. DEFRA/RAS/04.002. Defra, London.
- Defra, 2005a. 1: Marine Environment Quality. Report 1 of 5 contributing to Charting Progress: an Integrated Assessment of the State of UK Seas. Defra, London.
- Defra, 2005b. 5: Integrated Regional Assessment. Report 5 of 5 contributing to Charting Progress: an Integrated Assessment of the State of UK Seas. Defra, London.
- Defra, Department of the Environment Northern Ireland, Scottish Executive, Welsh Assembly Government, 2005. Charting Progress. An Integrated Assessment of the State of UK Seas. Defra, London.
- Defra, 2006. Industry profile. Industrial activities which have used materials containing radioactivity. Defra, London.
- Defra, 2007. Low level radioactive waste policy announced. News release 26 March 2007. Defra, London.
- Defra, Department for Business, Enterprise and Regulatory Reform, National Assembly for Wales and Northern Ireland Assembly, 2008. Managing Radioactive Waste Safely: A framework for Implementing Geological Disposal, 2008. Cm7386. The Stationery Office, London.
- Defra, 2010. Charting Progress 2. Defra, London.
- Defra, 2016. Drinking Water Regulations: Consultation on the draft Water Supply (Water Quality) Regulations 2016. Defra, London.
- Department for Business, Enterprise and Regulatory Reform, 2008. Meeting the energy challenge. A White Paper on Nuclear Power. Cmnd.7296. HMSO, London.
- Department of the Environment Northern Ireland, 2010. A Northern Ireland Marine Bill – policy proposals. Consultation document. DoENI, Belfast.
- Department of the Environment Northern Ireland, 2011. The State of the Seas Report. DoENI, Belfast.
- <https://www.daera-ni.gov.uk/publications/state-seas-report>
- DETR, 2000. Radioactive Substances (Basic Safety Standards) (England and Wales) Direction 2000. DETR, London.
- Dewar, A., Camplin, W., Barry, J. and Kennedy, P., 2014. A statistical approach to investigating enhancement of polonium-210 in the Eastern Irish Sea arising from discharges from a former phosphate processing plant. Journal of Environmental Radioactivity, 138: 289-301.
- Dick, R., 2012. Personal communication. Thames Water Utilities Limited, Reading.
- DPAG, 2008. 4th Report, November 2008. SEPA, Stirling.
- DSTL Radiological Protection Services, 2013. Marine environmental radioactivity surveys at nuclear submarine berths 2011. The Stationery Office, London.
- EC, 2011. Council implementing regulation (EU) No 297/2011 of 25 March 2011 imposing special conditions governing the import of feed and food originating in or consigned from Japan following the accident at the Fukushima nuclear power station. EC, Brussels.

- EC, 2012. ACSUK\_MSFD8bPressures\_20121101, 1 Nov 2012. EC, Brussels.
- EC, 2013. Council Directive 2013/51/EURATOM of 22 October 2013 laying down requirements for the protection of the health of the general public with regard to radioactive substances in water intended for human consumption. OJ L 296/2. EC, Brussels.
- EC, 2014. Council Directive 2013/59/EURATOM laying down basic safety standards for protection against the dangers arising from exposure to ionising radiation. OJ L 13, 17.1.2014:1-73. EC, Brussels.
- EC, 2016. Council implementing regulation (EU) 2016/6 of 5 January 2016 imposing special conditions governing the import of feed and food originating in or consigned from Japan following the accident at the Fukushima nuclear power station and repealing Implementation Regulation (EU) No. 322/2014. EC, Brussels.
- Elliott, J., Clyne F.J. and Garrod C.J. 2010. Radiological Habits Survey: Derby 2009. Project C2848. RL 05/10. Cefas, Lowestoft.
- Environment Agency, 2002a. Radioactivity in the environment. Report for 2001. Environment Agency, Lancaster.
- Environment Agency, 2002b. Radioactive Waste Regulation. Summary of Research 1996 – 2001. R&D Publication 129. Environment Agency, Bristol and London.
- Environment Agency, 2006a. Initial radiological assessment methodology – part 1 user report. SC030162/SR1. Environment Agency, Bristol and London.
- Environment Agency, 2006b. Initial radiological assessment methodology – part 2 methods and input data. SC030162/SR2. Environment Agency, Bristol and London.
- Environment Agency, Environment and Heritage Service, FSA and SEPA, 2007. Radioactivity in Food and the Environment, 2006. RIFE-12. Environment Agency, Environment and Heritage Service, FSA and SEPA. Bristol, Belfast, London and Stirling.
- Environment Agency, 2008a. Radioactive substances regulation. Environmental Principles. Public consultation. Environment Agency, Bristol and London.
- Environment Agency, 2008b. Assessment of Best Available Techniques (BAT). Consultation draft. Environment Agency, Bristol and London.
- Environment Agency, 2008c. Sellafield Radioactive Particles in the Environment – Programme of Work, February 2008. Environment Agency, Bristol and London.
- Environment Agency, 2009a. Habitats assessment for Radioactive Substances. Science report: SC060083/SR1, May 2009. Environment Agency, Bristol.
- Environment Agency, 2009b. Impact of radioactive substances on Ribble and Alt estuarine habitats. Science report: SC060083/SR2. May 2009. Environment Agency, Bristol.
- Environment Agency, 2009c. Dealing with contaminated land in England and Wales. Environment Agency, Bristol and London.
- Environment Agency and NIEA, 2009. Geological disposal facilities on land for solid radioactive wastes: guidance on requirements for authorisation. Environment Agency and NIEA, Bristol, London and Belfast.
- Environment Agency, NIEA and SEPA, 2009. Near-surface disposal facilities on land for solid radioactive wastes: guidance on requirements for authorisation. Environment Agency, NIEA and SEPA, Bristol, London, Belfast and Stirling.
- Environment Agency, FSA and SEPA, 2010. Environmental Radiological Monitoring. Radiological Monitoring Technical Guidance Note 2. Environment Agency, FSA and SEPA, Bristol, London and Stirling.
- Environment Agency, 2010. Briefing Note – Enhanced Beach Monitoring for Radioactive Particles near Sellafield site, March 2010. Environment Agency, Bristol and London.
- Environment Agency, FSA, NIEA and SEPA, 2010a. Radioactivity in Food and the Environment, 2009. RIFE 15. Environment Agency, FSA, NIEA and SEPA, Bristol, London, Belfast and Stirling.
- Environment Agency, FSA, NIEA and SEPA, 2010b. Summary of Radioactivity in Food and the Environment 2004-2008. Environment Agency, FSA, NIEA and SEPA, Bristol, London, Belfast and Stirling.
- Environment Agency, 2011. Briefing Note – Enhanced Beach Monitoring for Radioactive Particles near Sellafield site, April 2011. Environment Agency, Bristol and London.
- Environment Agency, 2012. Radioactive Contaminated Land. Environment Agency, Bristol and London.
- Environment Agency, SEPA, NIEA, HPA and FSA, 2012. Principles for the assessment of prospective public doses arising from authorised discharges of radioactive waste to the environment. Environment Agency, SEPA, NIEA, HPA and FSA. Bristol, Stirling, Belfast, Chilton and London.

- Environment Agency, 2013a. Guidance Note for Developers and Operators of Radioactive Waste Disposal Facilities in England and Wales. Environment Agency, Bristol and London.
- Environment Agency, 2013b. An environmental risk assessment for shale gas exploratory operations in England. Environment Agency, Bristol and London.
- Environment Agency, 2013c. Briefing Note – Enhanced Monitoring for Radioactive Particles, May 2013. Environment Agency, Bristol and London.
- Environment Agency, FSA, NIEA and SEPA, 2013. Radioactivity in Food and the Environment, 2012. RIFE 18. Environment Agency, FSA, NIEA and SEPA, Bristol, London, Belfast and Stirling.
- Environment Agency, FSA, NIEA, NRW and SEPA, 2014. Radioactivity in Food and the Environment, 2013. RIFE 19. Environment Agency, FSA, NIEA, NRW and SEPA, Bristol, London, Belfast, Cardiff and Stirling.
- Environment Agency, FSA, FSS, NIEA, NRW and SEPA, 2015. Radioactivity in Food and the Environment, 2014. RIFE 20. Environment Agency, FSA, FSS, NIEA, NRW and SEPA, Bristol, London, Aberdeen, Belfast, Cardiff and Stirling.
- Environment Agency and ONR, 2016. Generic design assessment. Progress report. Nov. 2015 – Jan. 2016. Environment Agency and ONR, Bristol and Bootle.
- Environment Agency and DECC, 2015. Environment Agency grants shale gas permit for Roseacre Wood. Press Release 6/2/2015. Environment Agency and DECC, Bristol and London.
- Etherington, G., Youngman, M.J., Brown, J. and Oatway, W., 2012. Evaluation of the Groundhog Synergy Beach Monitoring System for Detection of Alpha-rich Objects and Implications for the Health Risks to Beach Users. HPA-CRCE-038, August 2012, HPA, Chilton.
- FSA, 2001a. Consultative Exercise on Dose Assessment, 3 and 4 October 2000. FSA/0022/0501.500. FSA, London.
- FSA, 2001b. Radiological survey of foodstuffs from the Cardiff area. Food Survey Information Sheet 18/01.
- FSA and SEPA, 2002. Radioactivity in Food and the Environment, 2001. RIFE-7. FSA and SEPA, London and Stirling.
- FSA, 2003. Analysis of farmed salmon for technetium-99 and other radionuclides. Food Survey Information Sheet Number 39/03. FSA, London.
- FSA, 2004. Review of FSA research programmes on radiological protection. Research review report for the period 1998 – 2003. FSA, London.
- FSA, 2009. Estimate of the Food Chain Risks to Inform an Assessment of the Need for and Extent of the Food and Environment Protection Act Area at Dounreay. FSA, Aberdeen.
- <http://www.food.gov.uk/multimedia/pdfs/dounreayfepariskassessment.pdf>
- FSA, 2012a. Radioactivity in Food Monitoring Review. FSA, London.
- FSA, 2012b. The removal of post-Chernobyl sheep controls. FSA Board Paper 12/03/06. FSA, London.
- FSA, 2013. Radioactivity in Food Monitoring Review. Summary report of responses to consultation from stakeholders. FOODSA0128. FSA, London.
- Garrod, C.J., Clyne, F.J., Ly, V.E. and Rumney, P., 2012. Radiological Habits Survey: Heysham, 2011. Project C2848. RL 02/12. Cefas, Lowestoft.
- Garrod, C.J., Clyne, F.J., Ly, V.E. and Papworth, G.P., 2013a. Radiological Habits Survey: Dumfries and Galloway Coast, 2012. RL 25/13. Cefas, Lowestoft.
- Garrod, C.J., Clyne, F.J., Ly, V.E., Rumney, P. and Papworth, G.P., 2013b. Radiological Habits Survey: Barrow and the south-west Cumbrian coast, 2012. RL 01/13. Cefas, Lowestoft.
- Garrod, C.J., Clyne, F.J. and Papworth, G.P., 2014. Radiological Habits Survey: Wylfa, 2013. RL 03/14. Cefas, Lowestoft.
- Garrod, C.J., Clyne, F.J. and Papworth, G.P., 2015b. Radiological Habits Survey: Hartlepool, 2014. RL 01/15. Cefas, Lowestoft.
- Garrod, C.J., Clyne, F.J. and Rumney, P. 2016. Radiological Habits Survey: Sizewell, 2015. RL 01/16. Cefas, Lowestoft.
- Graven, H.D. and Gruber, N., 2011. Continental-scale enrichment of atmospheric  $^{14}\text{CO}_2$  from the nuclear power industry: potential impact on the estimation of fossil fuel derived  $\text{CO}_2$ . Atmos. Chem. Phys., 11, 12339-12346.
- Ham, G.J., Wilding, D. and Wilkins, B.T., 2007. Tritium concentrations in crops fertilised with contaminated sewage sludge. R01061. HPA-RPD-038. HPA, Chilton.
- Harrison, J.D. and Phipps, A., 2001. Invited editorial: gut transfer and doses from environmental technetium. J. Radiol. Prot., 21: (9 – 11).

- Harrison, J.D., Khurshed, A and Lambert, B.E., 2002. Uncertainties in dose coefficients for intakes of tritiated water and organically bound forms of tritium by members of the public. *Radiation Protection Dosimetry*, 98, 299 – 311.
- Harvey, M., Smith, J. and Cabianca, T., 2010. Assessment of collective and per caput doses due to discharges of radionuclides from the oil and gas industry into the marine environment. RPD-EA-4-2010. HPA, Chilton.
- HMIP, 1995. Routine measurement of gamma ray air kerma rate in the environment. Technical Guidance Note (Monitoring) M5. HMSO, London.
- Hodgson, A., Scott, J.E., Fell, T.P., and Harrison, J.D., 2005. Doses from the consumption of Cardiff Bay flounder containing organically bound tritium. Project SC020042/SR. Environment Agency, Bristol.
- HPA, 2007. Review of the risks from tritium. Report of the Independent Advisory Group on Ionising Radiation. RCE-4. HPA, London.
- HPA, 2009. Application of the 2007 Recommendations of the ICRP to the UK. Advice from the HPA. HPA, London.
- Hughes, L.M., Runacres, S.M. and Leonard, K.S., 2011. Marine Radioactivity in the Channel Islands, 1990 – 2009. *Environmental Radiochemical Analysis IV*, 170-180.
- Hunt, G.J., Hewitt, C.J. and Shepherd, J.G., 1982. The identification of critical groups and its application to fish and shellfish consumers in the coastal area of the north-east Irish Sea. *Health Physics* 43 (6) 875 – 889.
- Hunt, G.J., 1984. Simple models for prediction of external radiation exposure from aquatic pathways. *Radiat. Prot. Dosim.*, 8: 215 – 224.
- Hunt, G.J., Leonard, D.R.P. and Lovett, M.B., 1986. Transfer of environmental plutonium and americium across the human gut. *Sci. Total Environ.*, 53: 89 – 109.
- Hunt, G.J., 1998. Transfer across the human gut of environmental plutonium, americium, cobalt, caesium and technetium: studies with cockles (*Cerastoderma edule*) from the Irish Sea. *J. Radiol. Prot.*, 18(2): 101 – 109.
- Hunt, G.J., Leonard, D.R.P. and Lovett, M.B., 1990. Transfer of environmental plutonium and americium across the human gut. *Sci. Total Environ.*, 90: 273 – 282.
- Hunt, G.J. and Allington, D.J., 1993. Absorption of environmental polonium-210 by the human gut. *J. Radiol. Prot.*, 13(2):119 – 126.
- Hunt, G.J., Young, A.K. and Bonfield, R.A., 2001. Transfer across the human gut of environmental technetium in lobsters (*Homarus gammarus L.*) from the Irish Sea. *J. Radiol. Prot.*, 21: 21 – 29.
- Hunt, G.J. and Rumney, H.S., 2004. The human gut transfer of environmental polonium-210. *Proc. Int. Conf. on widening the radiation protection world*, 23 – 28 May 2004, Madrid. IRPA, Fontenay-aux-Roses.
- Hunt, G.J. and Rumney, H.S., 2005. The human alimentary tract transfer of environmental polonium-210. *Proceedings of the Seventh International Symposium of the Society for Radiological Protection*, 12th-17th June 2005, Cardiff. SRP, London.
- Hunt, G.J. and Rumney, H.S., 2007. The human alimentary tract transfer and body retention of environmental polonium-210. *J. Radiol. Prot.*, 27(4):405-26.
- Hunt, G.J., Leonard, K.S., Gitzinger, C., Janssens, A., Godeanu-Metz, A. and Henrich, E., 2007. Article-35 of the Euratom Treaty: Overview of national radioactive discharge and environmental monitoring requirements in the European Union. *Extended Synopsis in International Conference on Environmental Radioactivity, "From Measurement and Assessments in Regulations"*, 23-27 April 2007, Vienna.
- Hunt, J., Bailey, T. and Reese, A., 2009. The human body retention time of environmental organically bound tritium. *J. Radiol. Prot.*, 29(1):23-36.
- Hunt, G.J., Bailey, T.A., Jenkinson, S.B. and Leonard, K.S., 2010. Enhancement of tritium on uptake by marine biota: experience from UK coastal waters. *J. Radiol. Prot.*, 30(1):73.
- Hunt, G.J., Leonard, K.S. and Hughes, L.M., 2013. Artificial radionuclides in the Irish Sea from Sellafield: remobilisation revisited. *J. Radiol. Prot.*, 33(2):261.
- IAEA, 1996. International basic safety standards for protection against ionising radiation and for the safety of radiation sources. *Saf. Ser. No. 115*. IAEA Vienna.
- IAEA, 1997. Joint Convention on the Safety of Spent Fuel Management and on the Safety of Radioactive Waste Management. INF CIRC/546. IAEA, Vienna.
- IAEA, 1999. Application of radiological exclusion and exemption principles to sea disposal. IAEA-TECDOC-1068. IAEA, Vienna.
- IAEA, 2003. Determining the suitability of materials for disposal at sea under the London Convention 1972: a radiological assessment procedure. IAEA-TECDOC-1375. IAEA, Vienna.

- IAEA, 2015. Determining the suitability of materials for disposal at sea under the London Convention 1972 and London Protocol 1996: a radiological assessment procedure. IAEA-TECDOC-1759. IAEA, Vienna.
- ICRP, 1991. 1990 Recommendations of the ICRP. Annal. ICRP 21(1 – 3). Pergamon Press, Oxford, 201pp. (ICRP Publ. (60)).
- ICRP, 1994. Age-dependent doses to members of the public from intake of radionuclides: Part 2 Ingestion dose coefficients. Annal ICRP 23(3/4). Pergamon Press, Oxford, 167pp. (ICRP Publ. (67)).
- ICRP, 1996a. Age-dependent doses to members of the public from intake of radionuclides: Part 5 Compilation of ingestion and inhalation dose coefficients. Annal. ICRP 26(1). Elsevier Science, Oxford, 91pp. (ICRP Publ. (72)).
- ICRP, 1996b. Conversion coefficients for use in radiological protection against external radiation. Annal ICRP 26(3/4). Elsevier Science, Oxford (ICRP Publ. (74)).
- ICRP, 2001. Doses to the embryo and fetus from intakes of radionuclides by the mother. Annal ICRP 31(1 – 3). Elsevier Science, Oxford. (ICRP Publ. (88)).
- ICRP, 2007. The 2007 recommendations of the International Commission on Radiological Protection. Annal ICRP 37 (2-4). Elsevier Science, Oxford (ICRP Publ (103)).
- ICRP, 2008. Environmental protection: the concept and use of reference animals and plants. Annal ICRP 38(4-6). Elsevier Science, Oxford, 242pp. (ICRP Publ (108)).
- ICRP, 2014. Protection of the environment under different exposure situations. Annal ICRP 43(1). SAGE, Stanford, 58pp. (ICRP Publ (124)).
- International Organisation for Standardisation, 2005. General requirements for the competence of testing and calibration laboratories. ISO 17025.
- Jenkinson, S.B., McCubbin, D., Kennedy, P.H.W., Dewar, A., Bonfield, R and Leonard, K.S., 2014. An estimate of the inventory of technetium-99 in the sub-tidal sediments in the Irish Sea. J. Environ. Rad., 133, 40-47.
- Jobling, S., Williams, R., Johnson, A., Taylor, A., Gross-Sorokin, M., Nolan, M., Tyler, C., van Aerle, R., Santos, E., and Brighty, G., 2006. Predicted exposures to steroid estrogens in UK rivers correlate with widespread sexual disruption in wild fish populations. Environmental Health Perspective, 114 (S-1), 32 – 39.
- Joint Research Centre of the European Commission, 2009. Environmental Radioactivity in the European Community 2004 – 2006. Radiation Protection No. 161. CEC, Luxembourg.
- Jones, K.A., Smith, J.G., Anderson, T., Harvey, M.P., Brown, I., Field, S.J. and Jones, A.L., 2013. Implied doses to the population of the EU arising from reported discharges from EU nuclear power stations and reprocessing sites in the years 2004 to 2008, EC RP 176 [http://ec.europa.eu/energy/nuclear/radiation\\_protection/doc/publication/176.pdf](http://ec.europa.eu/energy/nuclear/radiation_protection/doc/publication/176.pdf)
- Jones, A.L. and Harvey, M.P., 2014. Radiological consequences resulting from accidents and incidents involving the transport of radioactive materials in the UK – 2012 review. PHE-CRCE-0147, Public Health England, Chilton.
- Kelly, M. and Emptage, M., 1991. Radiological assessment of the estuary at Ravensglass: Distribution of radioactivity and its relationship to sedimentary processes. Department of Environment, London.
- Kershaw, P.J. and Baxter, A.J., 1995. The transfer of reprocessing wastes from north-west Europe to the Arctic. Deep-Sea Res. II, 43(6): 1413 – 1448.
- Kibble, A., Cabianca, T., Daraktchieva, Z., Gooding, T., Smithard, J., Kowalczyk, G., McColl, N.P., Singh, M. Mitchem, L., Lamb, P., Vardoulakis S. and Kamanyire, R., 2014. Review of the potential public health impacts of exposure to chemical and radioactive pollutants as a result of shale gas extraction process. PHE-CRCE-009. Public Health England, Chilton.
- Knowles, J.F., Smith, D.L. and Winpenny K., 1998. A comparative study of the uptake, clearance and metabolism of technetium in lobster (*Homarus gammarus*) and edible crab (*Cancer pagurus*). Radiat. Prot. Dosim., 75: 125 – 129.
- Kocher, D.C. and Eckerman, K.F., 1987. Electron dose-rate conversion factors for external exposure of the skin from uniformly deposited activity on the body surface. Hlth. Phys., 53: 135 – 141.
- Leonard, D.R.P., Camplin, W.C. and Tipple, J.R., 1990. The variability of radiocaesium concentrations in freshwater fish caught in the UK following the Chernobyl nuclear reactor accident: an assessment of potential doses to critical group consumers. pp. 247 – 256. In: 'Proc. Int. Symp. on Environmental Contamination Following a Major Nuclear Accident'. IAEA, Vienna, IAEA-SM-306/15.
- Leonard, K.S., McCubbin, D., Brown, J., Bonfield, R. and Brooks, T., 1997a. A summary report of the distribution of Technetium-99 in UK Coastal Waters. Radioprotection, 32: 109 – 114.
- Leonard, K.S., McCubbin, D., Brown, J., Bonfield, R. and Brooks, T., 1997b. Distribution of technetium-99 in UK coastal waters. Mar. Pollut. Bull., 34(8): 628 – 636.

- Leonard, K.S., McCubbin, D., Blowers, P. and Taylor, B.R., 1999. Dissolved plutonium and americium in surface waters of the Irish Sea, 1973 – 96. *J. Environ. Rad.*, 44: 129 – 158.
- Leonard, K.S., McCubbin, D. and Bailey, T.A., 2001. Organic forms of tritium in food chains. Project R01023/C0814. RL 6/01. Cefas, Lowestoft.
- Leonard, K.S., McCubbin, D., McDonald, P., Service, M., Bonfield, R. and Conney, S., 2004. Accumulation of technetium-99 in the Irish Sea. *Sci. Tot. Env.*, 322: 255 – 270.
- LLW Repository Ltd., 2015. Launch of the LLWR Plan. News and Updates 3<sup>rd</sup> July 2014. LLW Repository Ltd., Holmrook.
- Ly, V.E., Garrod, C.J., Clyne, F.J. and Rumney, P., 2012. Radiological Habits Survey: Aldermaston and Burghfield, 2011. RL 03/12. Cefas, Lowestoft.
- Ly, V.E., Clyne, F.J., Garrod, C.J. and Dewar, A., 2013. Radiological Habits Survey: Springfields, 2012. Project C2848. RL 03/13. Cefas, Lowestoft.
- Ly V.E., Cogan S.M., Camplin W.C., Peake L. and Leonard, K.S., 2015. Long Term Trends in far-field effects of marine radioactivity measured around Northern Ireland. ERA12: Proceedings of the International Symposium on Nuclear and Environmental Radiochemical Analysis (17-19 September 2014, Bath, UK), 134 – 143, Royal Society of Chemistry, Cambridge
- MAFF, 1995. Terrestrial radioactivity monitoring programme (TRAMP) report for 1994. Radioactivity in food and agricultural products in England and Wales. MAFF, London, TRAMP/9, 223pp.
- MAFF, 1996. Pesticides Safety Directorate's Handbook. Appendix IC. MAFF, London.
- MAFF and SEPA, 1998. Radioactivity in Food and the Environment, 1997. RIFE-3. MAFF and SEPA, London and Stirling.
- MAFF and SEPA, 1999. Radioactivity in Food and the Environment, 1998. RIFE-4. MAFF and SEPA, London and Stirling.
- McCubbin, D., Leonard, K.S., Bailey, T.A., Williams, J. and Tossell, P., 2001. Incorporation of organic tritium (<sup>3</sup>H) by marine organisms and sediment in the Severn Estuary/ Bristol Channel (UK). *Mar. Pollut. Bull.*, 42 (10): 852 – 863.
- McCubbin, D., Leonard, K.S., Brown, J., Kershaw, P.J., Bonfield, R.A. and Peak, T., 2002. Further studies of the distribution of <sup>99</sup>Tc and <sup>137</sup>Cs in UK and European coastal waters. *Cont. Shelf. Res.* 22/10: 1417 – 1445.
- McCubbin, D. and Vivian, C., 2006. Dose assessments in relation to disposal at sea under the London Convention 1972: judging *de minimis* radioactivity. FSA Project R01062. RL5/06. Cefas, Lowestoft.
- McCubbin, D., Jenkinson, S.B., Leonard, K.S., Bonfield, R.A. and McMeekan, I.T., 2008. An assessment of the availability of Tc-99 to marine foodstuffs from contaminated sediments. Project R01062. RL09/08. Cefas, Lowestoft.
- McKay, W.A., Barr H.M., Halliwell C.M., Spencer D., Adsley I. and Perks C.A., 1995. Site specific background dose rates in coastal areas. DoE/HMIP/RR/94/037. HMIP, London.
- McTaggart, K., 2003. Habits survey notebook, Unpublished. Cefas, Lowestoft.
- McTaggart, K.A., Tipple, J.R., Clyne, F.J. and McMeekan, I.T., 2004a. Radiological habits survey: Cardiff, 2003. RL 03/04. Cefas, Lowestoft.
- McTaggart, K.A., Tipple, J.R., Sherlock, M., Cogan, S.M., Joyce, A.E. and Clyne, F.J., 2004b. Radiological Habits Survey: Winfrith, 2003. Projects C1659, RB103 and C1666. RL02/04. Cefas, Lowestoft.
- Mobbs, S., Barraclough, I., Napier, I., Casey, A., Poynter, R. and Harvey, M., 1998. A review of the use and disposal of gaseous tritium light devices. Environment Agency, Lancaster.
- Mitchell, N., 2015. Environmental Safety Case: Disposal of low activity low-level radioactive waste at Northants Resource Management Facility. ENE-154/001. Eden Nuclear and Environment Ltd., Penrith.
- National Dose Assessment Working Group, 2004. Radiological Assessment Exposure Pathways Checklist (Common and Unusual). NDAWG/2/2004. Environment Agency, FSA, NRPB, NII, Chilton.
- Natural Scotland and SEPA, 2016. Alienated Land Former RAF Kinloss: Part IIA Inspection and Risk Assessment report. Scottish Government, Edinburgh.
- NDA, 2016a. Strategy effective from April 2016. NDA, Moor Row, Cumbria.
- NDA, 2016b. NDA Business Plan 2016/2019. NDA, Moor Row, Cumbria.
- NDA, 2016c. NDA Strategy. Integrated Impact Assessment Report. NDA, Moor Row, Cumbria.
- NDA and DECC, 2014. Radioactive Wastes in the UK: A Summary of the 2013 Inventory. NDA, Moor Row, Cumbria.

- NIEA, 2013. Shale gas regulators forum. Minutes of 2<sup>nd</sup> meeting, March 2013. NIEA, Belfast.
- Northern Ireland Assembly, 2003. Radioactive Substances (Basic Safety Standards) Regulations (Northern Ireland).
- NRPB, 1990. Gut transfer factors. Docs. NRPB 1(2). NRPB, Chilton, 26pp.
- NRPB, 2005. Guidance on the application of dose coefficients for the embryo and fetus from intakes of radionuclides by the mother. Docs NRPB 16(2). NRPB, Chilton, 41pp.
- NRW, SEPA and the Environment Agency, 2016. Summary brief of consultation and new Guidance on Requirements for Release of Nuclear Sites from Radioactive Substances Regulation. NRW, Cardiff.
- Oatway, W. and Brown, J., 2015. Health Risk to Seafood Consumers from Radioactive Particles in the marine environment in the vicinity of the Sellafield site. PHE-CRCE-021. Public Health England, Chilton.
- OECD, Nuclear Energy Agency, 1985. Review of the continued suitability of the dumping site for radioactive waste in the North-East Atlantic. OECD, Paris, 448pp.
- ONR, 2013. The decision on the application to carry out a decommissioning project at Wylfa nuclear power station under the Nuclear Reactors (Environmental Impact Assessment for Decommissioning) Regulations 1999 (as amended). November 2013. ONR, Bootle.
- ONR, 2015. The decision on the application to carry out a decommissioning project on the CONSORT reactor at Imperial College Silwood Park Campus under the Nuclear Reactors (Environmental Impact Assessment for Decommissioning) Regulations 1999 (as amended). Report ONR-DFW-PAR15-005, July 2015. ONR, Bootle.
- ONR, 2016. Hinkley Point C 9 March 2016 News. ONR, Bootle.
- OSPAR, 1998. SINTRA Statement. Summary Record OSPAR 98/14/1, Annex 45. OSPAR, London.
- OSPAR, 2000a. Convention for the protection of the marine environment of the North-East Atlantic. OSPAR, London.
- OSPAR, 2000b. Quality Status Report 2000. OSPAR, London.
- OSPAR, 2003. 2003 Strategies of the OSPAR Commission for the Protection of the Marine Environment of the North-East Atlantic. 2003-21. OSPAR, London.
- OSPAR, 2006. Agreement on a monitoring programme for concentrations of radioactive substances in the marine environment. Annex 14, ref 9.8c, Malahide (Ireland): 27 June-1 July 2005. OSPAR, London.
- OSPAR, 2009. Towards the Radioactive Substances Strategy Objectives. Third Periodic Evaluation, OSPAR, London.
- OSPAR, 2010a. The North-East Atlantic Environment Strategy. Strategy of the OSPAR Commission for the Protection of the Marine Environment of the North-East Atlantic 2010-2020. OSPAR Agreement 2010-3. OSPAR, London.
- OSPAR, 2010b. Quality Status Report 2010. OSPAR, London.
- OSPAR, 2011a. Liquid discharges from nuclear installations in 2009. OSPAR, London.
- OSPAR, 2011b. Discharges of radioactive substances from the non-nuclear sectors in 2009. OSPAR, London.
- OSPAR, 2013. UK report on application of Best Available Techniques (BAT) in civil nuclear facilities (2008-2011). OSPAR, London.
- OSPAR, 2016a. Summary Record. Meeting of the Radioactive Substances Committee (RSC), Brussels: 9-11 February 2016. OSPAR, London.
- OSPAR, 2016b. Towards the Radioactive Substances Strategy Objectives. Forth Periodic Evaluation, OSPAR, London.
- Ould-Dada, Z., 2000. Sea-to-Land transfer of radionuclides. How much do we know. Proceedings of the second RADREM-TEEC Workshop held in London on 21 January 1999. DETR/RADREM/00.001 DETR, London.
- Papworth, G.P., Garrod, C.J. and Clyne, F.J., 2014. Radiological Habits Survey: Dounreay, 2013. RL 06/14. Cefas, Lowestoft.
- PRAG (D), 2010. Annual report to SEPA and DSRL, March 2010. SEPA, Stirling.
- PRAG (D), 2011. Annual report to SEPA and DSRL, March 2011. SEPA, Stirling.
- PRAG(D), (2016). Annual report to SEPA and DSRL, SEPA, Stirling. March 2016. SEPA, Stirling.
- Povinec, P.P., Bailly Du Bois, P., Kershaw, P.J., Nies, H. and Scotto, P., 2003. Temporal and spatial trends in the distribution of <sup>137</sup>Cs in surface waters of Northern European Seas - a record of 40 years of investigations. Deep Sea Res. II, 50: 2785 – 2801.

- Povinec, P.P., Aarkrog, A., Buessler, K.O., Delfanti, R., Hirose, K., Hong, G.H., Ito, T., Livingston, H.D., Nies, H., Noshkin, V. E., Shima, S. and Togawa, O., 2005.  $^{90}\text{Sr}$ ,  $^{137}\text{Cs}$  and  $^{239,240}\text{Pu}$  concentration surface water time series in the Pacific and Indian Oceans - WOMARS results. *J. Environ. Rad.*, 81: 63 - 87.
- Preston, A., Mitchell, N.T. and Jefferies, D.F., 1974. Experience gained in applying the ICRP Critical Group concept to the assessment of public radiation exposure in control of liquid waste disposal. *Proc. Symp. IAEA Portoroz, IAEA-SM-184/10*, 131 – 146.
- Radioactive Waste Management, 2015. Providing information on geology. *Radioactive Waste Management*, Harwell.
- Rollo, S.F.N., Camplin, W.C., Allington, D.J. and Young, A.K., 1992. Natural radionuclides in the UK marine environment. In: 'Proceedings of the Fifth International Symposium on Natural Radiation Environment, Salzburg, September 22 – 28, 1991'. *Radiat. Prot. Dosim.*, 45(1/4): 203 – 210.
- Rollo, S.F.N., Camplin, W.C., Duckett, L., Lovett, M.B. and Young, A.K., 1994. Airborne radioactivity in the Ribble Estuary. pp277 – 280. In: 'Proc. IRPA Regional Congress on Radiological Protection, 6 – 10 June 1994, Portsmouth, UK'. Nuclear Technology Publishing.
- Rumney, P., Ly, V.E. and Clyne, F.J., 2013a. Radiological Habits Survey: Hunterston, 2012. RL 26/13. Cefas, Lowestoft.
- Rumney, P., Garrod, C.J. and Clyne, F.J., 2013b. Radiological Habits Survey: Faslane, 2011. RL 21/13. Cefas, Lowestoft.
- Scotland - Parliament, 2010. Marine (Scotland) Act, 2010. OQPS, Edinburgh.
- Scotland - Parliament, 2015. Food (Scotland) Act 2015. HMSO, London.
- Scottish Executive, 2000. Radioactive Substances (Basic Safety Standards) (Scotland) Direction 2000. Scottish Executive, Edinburgh.
- Scottish Government, 2008. Environment Act 1995. The UK Strategy for Radioactive Discharges. Statutory Guidance. Scottish Government, Edinburgh.
- Scottish Government, 2011. Scotland's Higher Activity Radioactive Waste Policy, 2011, Scottish Government, Edinburgh.
- Sellafield Limited, 2015. Particles in the Environment. Annual Report for 2014/15 and Forward programme. NDA, Seascale.
- Sellafield Limited, 2016. Particles in the Environment. Annual Report for 2015/16 and Forward programme. NDA, Seascale.
- SEPA, 2007. Strategy for the Assessment of the potential impact of Sellafield Radioactive Particles on Southwest Scotland, December 2007. SEPA, Stirling.
- SEPA, 2012a. Satisfying the ALARA requirement and the role of Best Practicable Means. SEPA, Stirling.
- SEPA, 2012b. SEPA Policy on the Regulation of Disposal of Radioactive Low Level Waste from Nuclear Sites. SEPA, Stirling.
- SEPA, 2013. Regulatory guidance: coal bed methane and shale gas. SEPA, Stirling.
- SEPA, in press. Radiological Habits Survey: Chapelcross. SEPA, Stirling.
- SEPA, in press. Radiological Habits Survey: Rosyth. SEPA, Stirling.
- Simmonds, J.R., Lawson, G. and Mayall, A., 1995. Radiation Protection 72; Methodology for assessing the radiological consequences of routine releases of radionuclides to the environment. Report EUR 15760 EN. Office for Official Publications of the European Community, Luxembourg.
- Sellafield Limited, 2016. Particles in the Environment. Annual Report for 2015/16 and Forward programme. NDA, Seascale.
- Smith, B.D. and Jeffs, T.M., 1999. Transfer of radioactivity from fishmeal in animal feeding stuffs to man. RL 8/99. Cefas, Lowestoft.
- Smith, K.R. and Jones, A.L. 2003. Generalised habit data for radiological assessments. NRPB-W41. NRPB, Chilton.
- Smith, J., Ottway, W., Brown, I. and Sherwood, J., 2009. PC Cream 08 User Guide. RPD-EA-9-2009. HPA, Chilton.
- Smith, D.L., Smith, B.D., Joyce, A.E. and McMeekan, I.T., 2002. An assessment of aquatic radiation exposure pathways in Northern Ireland. SR(02)14. RL 20/02. Scotland and Northern Ireland Forum for Environmental Research, Edinburgh.
- Smith, K.R., Mobbs, S.F. and Cooper, J.R., 2006. Dose criteria for the designation of radioactivity contaminated land. RCE-2. HPA, Chilton.
- Statutory Instruments, 2007. SI 2007 No 3236. The Radioactive Contaminated Land (Amendment) Regulations (Northern Ireland) 2007. HMSO, London.

- Statutory Instruments, 2010. SI 2010 No 2145. The Radioactive Contaminated Land (Amendment) Regulations (Northern Ireland) 2010. HMSO, London.
- Statutory Instruments, 2015. SI 2015 No 209. Justification Decision (UK ABWR) Regulations 2010. HMSO, London.
- Swift, D.J., 2001. Cardiff radiological survey of selected foodstuffs. Project C1003. RL 11/01. Cefas, Lowestoft.
- Swift, D.J. and Nicholson, M.D., 2001. Variability in the edible fraction content of  $^{60}\text{Co}$ ,  $^{99}\text{Tc}$ ,  $^{110\text{m}}\text{Ag}$ ,  $^{137}\text{Cs}$  and  $^{241}\text{Am}$  between individual crabs and lobsters from Sellafield (north eastern Irish Sea). *J. Environ. Radioact.*, 54, 311 – 326.
- Tipple, J.R., McTaggart, K., Clyne, F.J. and Sherlock, M., 2006. Radiological Habits Survey: Trawsfynydd, 2005. Projects C1659, RB103 and C1666. RL 02/06. Cefas, Lowestoft.
- Tipple, J.R., Jeffs, T.M., Clyne, F.J., Garrod, C.J. and Earl, T.J., 2009. Radiological Habits Survey: Capenhurst, 2008. Project C2848. RL 03/09. Cefas, Lowestoft.
- United Kingdom - Parliament, 1965. Nuclear Installations Act, 1965. HMSO, London.
- United Kingdom - Parliament, 1985. Food and Environment Protection Act, 1985. HMSO, London.
- United Kingdom - Parliament, 1989. The Sludge (Use in Agriculture) Regulations 1989. HMSO, London.
- United Kingdom - Parliament, 1993. Radioactive Substances Act, 1993. HMSO, London.
- United Kingdom - Parliament, 1995a. Environment Act, 1995. HMSO, London.
- United Kingdom - Parliament, 1995b. Review of Radioactive Waste Management Policy. HMSO, London, 55pp. (Cm 2919).
- United Kingdom - Parliament, 1999. The Ionising Radiations Regulations 1999. Stat. Inst. 1999 No 3232. HMSO, London, 67pp.
- United Kingdom - Parliament, 2004. Energy Act, 2004. HMSO, London.
- United Kingdom – Parliament, 2009. Marine and Coastal Access Act 2009. HMSO, London
- United Kingdom - Parliament, 2010a. Environmental Permitting (England and Wales) Regulations. Stat. Inst. 2010 No 675. HMSO, London.
- United Kingdom - Parliament, 2010b. The Marine Strategy Regulations 2010. Stat. Inst. 2010 No 1627. HMSO, London.
- Watson, S.J., Jones, A.L., Oatway, W.B. and Hughes, J.S., 2005. Ionising radiation exposure of the UK population: 2005 Review. HPA-RPD-001. HPA, Chilton.
- Welsh Government, 2015. Welsh Government Policy on the Management and Disposal of Higher Activity Radioactive Waste, 2015, WG23160. Welsh Government, Cardiff.
- WHO, 2004. Guidelines for drinking water quality. 3rd Edition. WHO, Geneva.
- Williams, J.L., Russ, R.M., McCubbin, D. and Knowles, J.F., 2001. An overview of tritium behaviour in the Severn estuary (UK). *J. Rad. Prot.*, 21: 337 – 344.
- Wood, M.D., Marshall, W.A., Beresford, N.A., Jones, S.R., Howard, B.J., Copplestone, D. and Leah, R.T., 2008. Application of the ERICA Integrated Approach to the Drigg coastal sand dunes. *J. Environ. Radioact.* (Special Issue: The ERICA Project), 99 (9) 1484 – 1495.
- Wood, M.D., Hall, P.J., Wittrick, S., Copplestone, D. and Leah, R.T., 2011. Survey of gamma dose rates in air around the Esk Estuary. Environment Agency, Bristol and London.
- Young, A.K., McCubbin, D. and Camplin, W.C., 2002. Natural radionuclides in seafood. Project R03010/C0808. RL 17/02. Cefas, Lowestoft.
- Young, A.K., McCubbin, D., Thomas, K., Camplin, W.C., Leonard, K.S., and Wood, N., 2003. Po Concentrations in UK Seafood. 9th International Symposium on Environmental Radiochemical Analysis, 18-20 September 2002, Oxford. ERA III. Royal Society of Chemistry, London.



## APPENDIX 1. Sampling, measurement, presentation and assessment methods and data

This Appendix contains information on the methods of sampling, measurement, presentation and assessment used in the Radioactivity in Food and the Environment report. It is provided in a separate file to the main report at <https://www.gov.uk/government/policies/radioactive-and-nuclear-substances-and-waste>

## APPENDIX 2. Disposals of radioactive waste

**Table A2.1. Principal discharges of gaseous radioactive wastes from nuclear establishments in the United Kingdom, 2015**

Establishment	Radioactivity	Discharge limit (annual equivalent) <sup>a</sup> , Bq	Discharges during 2015	
			Bq	% of annual limit <sup>b</sup>
<b>Nuclear fuel production and reprocessing</b>				
Capenhurst (CNS Ltd)	Alpha	BAT	1.50E+06	NA
Other authorised outlets	Beta	BAT	9.11E+05	NA
Capenhurst (Urenco UK Ltd)	Uranium	7.50E+06	2.01E+06	27
	Other alpha	2.40E+06	Nil	Nil
	Technetium-99	1.00E+08	Nil	Nil
	Others	2.25E+09	Nil	Nil
	Alpha (Incinerator)	2.00E+08	Nil	Nil
	Beta (Incinerator)	2.50E+08	Nil	Nil
Sellafield <sup>c</sup>	Alpha	8.80E+08	8.20E+07	9.3
	Beta	4.20E+10	6.20E+08	1.5
	Tritium	1.10E+15	8.40E+13	7.6
	Carbon-14	3.30E+12	4.20E+11	13
	Krypton-85	4.40E+17	6.80E+16	15
	Strontium-90	7.10E+08	2.90E+07	4.1
	Ruthenium-106	2.30E+10	7.30E+08	3.2
	Antimony-125	3.00E+10	1.20E+10	40
	Iodine-129	7.00E+10	1.10E+10	16
	Iodine-131	3.70E+10	4.40E+08	1.2
	Caesium-137	5.80E+09	9.00E+07	1.6
	Radon-222	5.00E+11	4.30E+10	8.6
	Plutonium alpha	1.90E+08	2.60E+07	14
	Plutonium-241	3.00E+09	2.90E+08	9.7
	Americium-241 and curium-242	1.20E+08	1.60E+07	13
Springfields	Uranium	5.30E+09	3.36E+07	<1
Springfields (National Nuclear Laboratory)	Tritium	1.00E+08	4.75E+06	4.8
	Carbon-14	1.00E+07	4.36E+04	<1
	Other alpha radionuclides	1.00E+06	Nil	Nil
	Other beta radionuclides	1.00E+07	1.44E+03	<1
<b>Research establishments</b>				
Dounreay <sup>d, 1</sup>	Alpha <sup>e</sup>	3.10E+07	8.01E+06	26
	Non-alpha <sup>f</sup>	1.70E+09	1.21E+08	7.1
	Tritium	1.72E+13	4.29E+10	<1
	Krypton-85 <sup>g</sup>	5.69E+14	2.30E+05	<1
	Iodine-129	1.08E+08	3.03E+07	28
Harwell	Alpha	8.00E+05	3.00E+04	3.8
	Beta	2.00E+07	5.90E+05	2.9
	Tritium	1.50E+13	3.60E+11	2.4
	Krypton-85	2.00E+12	Nil	Nil
	Radon-220	1.00E+14	5.60E+12	5.6
	Radon-222	3.00E+12	2.40E+11	8.2
	Iodines	1.00E+10	Nil	Nil
	Other radionuclides	1.00E+11	Nil	Nil
Winfrith (Inutec)	Alpha	1.00E+05	Nil	Nil
	Tritium	1.95E+13	2.99E+11	1.5
	Carbon-14	3.00E+10	Nil	Nil
	Other	1.00E+05	Nil	Nil
Winfrith (Research Sites Restoration Ltd)	Alpha	2.00E+06	1.05E+03	<1
	Tritium	4.95E+13	2.24E+11	<1
	Carbon-14	5.90E+09	1.29E+08	2.2
	Other	5.00E+06	1.94E+04	<1

**Table A2.1. continued**

Establishment	Radioactivity	Discharge limit (annual equivalent) <sup>a</sup> , Bq	Discharges during 2015	
			Bq	% of annual limit <sup>b</sup>
<b>Minor sites</b>				
Imperial College Reactor Centre Ascot	Tritium	3.00E+08	7.38E+06	2.5
	Argon-41	1.70E+12	Nil	Nil
<b>Nuclear power stations</b>				
Berkeley <sup>h</sup>	Beta	2.00E+07	1.16E+05	<1
	Tritium	2.00E+10	1.40E+10	70
	Carbon-14	5.00E+09	9.16E+08	18
Bradwell	Beta	6.00E+08	5.65E+05	<1
	Tritium	6.00E+12	1.71E+11	2.9
	Carbon-14	9.00E+11	1.37E+10	1.5
Chapelcross	Tritium	2.30E+14	2.58E+13	11
	All other nuclides	5.15E+09	7.88E+08	15
Dungeness A Station	Beta <sup>i</sup>	5.00E+08	8.69E+06	1.7
	Tritium	2.60E+12	1.24E+10	<1
	Carbon-14	5.00E+12	3.53E+08	<1
Dungeness B Station	Tritium	1.20E+13	1.68E+12	14
	Carbon-14	3.70E+12	8.56E+11	23
	Sulphur-35	3.00E+11	3.78E+10	13
	Argon-41	7.50E+13	1.26E+13	17
	Cobalt-60 <sup>i</sup>	1.00E+08	3.38E+06	3.4
	Iodine-131	1.50E+09	2.97E+07	2.0
Hartlepool	Tritium	1.00E+13	5.88E+11	5.9
	Carbon-14	4.50E+12	1.81E+12	40
	Sulphur-35	2.30E+11	1.82E+10	7.9
	Argon-41	1.50E+14	1.01E+13	6.7
	Cobalt-60 <sup>i</sup>	1.00E+08	2.37E+07	24
	Iodine-131	1.50E+09	1.82E+08	12
Heysham Station 1	Tritium	1.00E+13	8.27E+11	8.3
	Carbon-14	4.50E+12	1.00E+12	22
	Sulphur-35	2.00E+11	2.37E+10	12
	Argon-41	1.50E+14	3.62E+12	2.4
	Cobalt-60 <sup>i</sup>	1.00E+08	7.61E+06	7.6
	Iodine-131	1.50E+09	7.93E+07	5.3
Heysham Station 2	Tritium	1.00E+13	1.34E+12	13
	Carbon-14	3.70E+12	1.73E+12	47
	Sulphur-35	2.30E+11	9.91E+09	4.3
	Argon-41	7.50E+13	1.39E+13	19
	Cobalt-60 <sup>i</sup>	1.00E+08	8.38E+06	8.4
	Iodine-131	1.50E+09	7.40E+07	4.9
Hinkley Point A Station	Beta	5.00E+07	2.16E+05	<1
	Tritium	7.50E+11	1.87E+10	2.5
	Carbon-14	5.00E+10	4.57E+08	<1
Hinkley Point B Station	Tritium	1.20E+13	1.75E+12	15
	Carbon-14	3.70E+12	1.48E+12	40
	Sulphur-35	3.50E+11	7.00E+10	20
	Argon-41	1.00E+14	1.45E+13	15
	Cobalt-60 <sup>i</sup>	1.00E+08	8.38E+06	8.4
	Iodine-131	1.50E+09	5.22E+06	<1
Hunterston A Station <sup>2</sup>	Tritium	2.00E+10	6.29E+08	3.1
	Carbon-14	2.00E+09	5.77E+07	2.9
	All other radionuclides	3.00E+06	6.74E+05	22

**Table A2.1. continued**

Establishment	Radioactivity	Discharge limit (annual equivalent) <sup>a</sup> , Bq	Discharges during 2015	
			Bq	% of annual limit <sup>b</sup>
Hunterston B Station <sup>d</sup>	Particulate beta	5.00E+08	5.91E+07	12
	Tritium	1.50E+13	2.58E+12	17
	Carbon-14	4.50E+12	1.53E+12	34
	Sulphur-35	5.00E+11	9.89E+10	20
	Iodine-131	2.00E+09	4.04E+06	<1
Oldbury <sup>3</sup>	Beta	1.00E+08	4.80E+05	<1
	Tritium	9.00E+12	6.07E+10	<1
Sizewell A Station	Beta	8.50E+08	Nil	Nil
	Tritium	3.50E+12	9.63E+09	<1
	Carbon-14	1.00E+11	2.58E+09	2.6
Sizewell B Station	Noble gases	3.00E+13	3.08E+12	10
	Particulate Beta	1.00E+08	7.00E+06	7.0
	Tritium	3.00E+12	6.94E+11	23
	Carbon-14	5.00E+11	2.45E+11	49
Torness	Particulate Beta	4.00E+08	5.54E+06	1.4
	Tritium	1.10E+13	1.11E+12	10
	Carbon-14	4.50E+12	1.28E+12	28
	Argon-41	7.50E+13	6.11E+12	8.1
	Iodine-131	2.00E+09	1.00E+07	<1
Trawsfynydd	Particulate Beta	5.00E+07	1.10E+06	2.2
	Carbon-14	1.00E+10	7.00E+08	7.0
Wylfa	Particulate Beta	7.00E+08	1.35E+07	1.9
	Tritium	1.80E+13	9.30E+11	5.2
	Sulphur-35	4.50E+11	1.03E+11	23
	Argon-41	1.00E+14	1.27E+13	13
<b>Defence establishments</b>				
Aldermaston <sup>i</sup>	Alpha	1.65E+05	2.77E+04	17
	Particulate Beta	6.00E+05	5.58E+03	<1
	Tritium	3.90E+13	3.50E+11	<1
	Carbon-14	6.00E+06	Nil	Nil
	Volatile beta	4.40E+06	3.58E+06	81
Barrow <sup>l</sup>	Tritium	3.20E+06	Nil	Nil
	Argon-41	4.80E+10	Nil	Nil
Burghfield <sup>l</sup>	Tritium	1.00E+10	Nil	Nil
	Alpha	5.00E+03	1.52E+03	30
Coulport	Tritium	5.00E+10	4.23E+09	8.5
Derby <sup>i,m</sup>	Alpha <sup>n</sup>	3.00E+06	6.81E+05	23
	Alpha <sup>o,p</sup>	2.40E+04	3.90E+01	<1
	Beta <sup>o,p</sup>	1.80E+06	4.26E+04	2.4
	Tritium	4.00E+09	6.40E+08	16
	Carbon-14	4.30E+10	1.01E+09	2.3
	Argon-41	1.50E+10	3.09E+08	2.1
(Vulcan)	Noble gases	5.00E+09	6.48E+08	13

**Table A2.1. continued**

Establishment	Radioactivity	Discharge limit (annual equivalent) <sup>a</sup> , Bq	Discharges during 2015	
			Bq	% of annual limit <sup>b</sup>
Rosyth <sup>r</sup>	Beta (particulate)	1.00E+05	Nil	Nil
	Tritium	2.00E+08	Nil	Nil
	Carbon-14	5.00E+08	Nil	Nil
<b>Radiochemical production</b>				
Amersham (GE Healthcare)	Alpha	2.25E+06	8.57E+04	3.8
	Radionuclides T1/2<2hr	7.50E+11	5.97E+09	<1
	Tritium	2.00E+12	1.08E+06	<1
	Radon-222	1.00E+13	2.37E+12	24
	Other including selenium-75 and iodine-131	1.60E+10	5.83E+06	<1
Cardiff (GE Healthcare) <sup>4</sup>	Tritium	6.00E+12	1.15E+12	19
	Carbon-14	1.10E+12	1.31E+11	12
<b>Industrial and landfill sites</b>				
LLWR	Alpha	BAT	8.33E+03	NA
	Beta	BAT	6.15E+04	NA
Lillyhall (Studsvik)	Alpha (particulate)	5.00E+05	4.30E+03	<1
	Beta (particulate)	5.00E+05	1.49E+04	3.0

\* As reported to SEPA and the Environment Agency

<sup>a</sup> In some cases permits specify limits in greater detail than can be summarised in a single table; in particular, periods shorter than one year are specified at some sites

<sup>b</sup> Data quoted to 2 significant figures except where values are <1%

<sup>c</sup> Limits for tritium, carbon-14, krypton-85 and iodine-129 vary with the mass of uranium processed by THORP

<sup>d</sup> Some discharges are upper estimates because they include 'less than' data derived from analyses of effluents at limits of detection

<sup>e</sup> All alpha emitting nuclides taken together

<sup>f</sup> All non-alpha emitting radionuclides, not specifically listed, taken together

<sup>g</sup> Krypton-85 discharges are calculated

<sup>h</sup> Combined data for Berkeley Power Station and Berkeley Centre

<sup>i</sup> Particulate activity

<sup>j</sup> Discharges were made by AWE plc

<sup>k</sup> Argon-41 is reported under the Activation products total and the limit is the demonstration of Best Available Technique

<sup>l</sup> Discharges from Barrow are included with those from MoD sites because they are related to submarine activities. Discharges were made by BAE Systems Marine Ltd

<sup>m</sup> Discharges were made by Rolls Royce Marine Power Operations Ltd

<sup>n</sup> Discharge limit is for the Nuclear Fuel Production Plant site

<sup>o</sup> Annual limits on beta and alpha derived from monthly and weekly notification levels

<sup>p</sup> Discharge limit is for the Neptune Reactor Raynesway site

<sup>q</sup> Discharges were made by Devonport Royal Dockyard Ltd

<sup>r</sup> Discharges were made by Rosyth Royal Dockyard Ltd

<sup>1</sup> Discharge authorisation revised with effect from 1 May 2014

<sup>2</sup> Discharge Authorisation revised with effect 01 July 2014

<sup>3</sup> Discharge permit revised with effect 1 May 2014, sulphur-35 and argon-41 are no longer in the permit

<sup>4</sup> Discharge permit revised with effect 1 June 2015

NA Not applicable under permit

BAT Best available technology

**Table A2.2. Principal discharges of liquid radioactive waste from nuclear establishments in the United Kingdom, 2015**

Establishment	Radioactivity	Discharge limit (annual equivalent) <sup>a</sup> , Bq	Discharges during 2015	
			Bq	% of annual limit <sup>b</sup>
<b>Nuclear fuel production and reprocessing</b>				
Capenhurst (Urenco UK Ltd)	Uranium	7.50E+08	4.67E+06	<1
	Uranium daughters	1.36E+09	4.72E+06	<1
	Non-uranic alpha	2.20E+08	1.15E+07	5.2
	Technetium-99	1.00E+09	4.35E+06	<1
Sellafield <sup>c,l</sup>	Alpha	9.00E+11	1.87E+11	21
	Beta	2.00E+14	9.53E+12	4.8
	Tritium	1.80E+16	1.54E+15	8.6
	Carbon-14	2.10E+13	4.96E+12	24
	Cobalt-60	3.60E+12	5.13E+10	1.4
	Strontium-90	4.50E+13	1.59E+12	3.5
	Zirconium-95 + Niobium-95	2.80E+12	9.64E+10	3.4
	Technetium-99	1.00E+13	1.66E+12	17
	Ruthenium-106	5.10E+13	7.03E+11	1.4
	Iodine-129	2.00E+12	3.65E+11	18
	Caesium-134	1.60E+12	6.67E+10	4.2
	Caesium-137	3.40E+13	3.08E+12	9.1
	Cerium-144	4.00E+12	1.94E+11	4.9
	Neptunium-237	7.30E+11	3.65E+10	5.0
	Plutonium alpha	7.00E+11	1.39E+11	20
	Plutonium-241	2.50E+13	2.36E+12	9.4
	Americium-241	3.00E+11	2.87E+10	10
Curium-243+244	5.00E+10	1.19E+09	2.4	
Uranium (in kg) <sup>d</sup>	2.00E+03	3.28E+02	16	
Springfields	Alpha	1.00E+11	2.29E+10	23
	Beta	2.00E+13	1.84E+12	9.2
	Technetium-99	6.00E+11	6.90E+09	1.2
	Thorium-230	2.00E+10	1.08E+09	5.4
	Thorium-232	1.50E+10	1.54E+08	1.0
	Neptunium-237	4.00E+10	1.90E+08	<1
	Other transuranic radionuclides	2.00E+10	1.19E+09	6.0
	Uranium	4.00E+10	1.83E+10	46
<b>Research establishments</b>				
Dounreay <sup>e,2</sup>	Alpha <sup>f</sup>	3.40E+09	2.09E+08	6.1
	Non-alpha <sup>g</sup>	4.80E+10	1.22E+09	2.5
	Tritium	6.90E+12	5.36E+10	<1
	Strontium-90	1.77E+11	2.25E+10	13
	Caesium-137	6.29E+11	2.81E+09	<1
Harwell (Lydebank Brook)	Alpha	3.00E+07	3.10E+06	10
	Beta	3.00E+08	6.98E+06	2.3
	Tritium	2.00E+10	1.33E+09	6.6
Harwell (sewer)	Alpha	1.00E+07	4.16E+05	4.2
	Beta	6.00E+08	3.05E+07	5.1
	Tritium	1.00E+11	1.31E+08	<1
	Cobalt-60	5.00E+06	9.99E+05	20
	Caesium-137	2.00E+08	7.72E+06	3.9
Winfrith (inner pipeline) <sup>h</sup>	Alpha	2.00E+10	5.78E+06	<1
	Tritium	2.20E+14	2.29E+13	10
	Caesium-137	2.00E+12	1.64E+09	<1
	Other radionuclides	1.00E+12	1.06E+09	<1
Winfrith (outer pipeline)	Alpha	2.00E+09	2.90E+06	<1
	Tritium	1.50E+11	1.71E+09	1.1
	Other radionuclides	1.00E+09	1.06E+07	1.1
Winfrith (River Frome)	Tritium	7.50E+11	Nil	Nil

**Table A2.2. continued**

Establishment	Radioactivity	Discharge limit (annual equivalent) <sup>a</sup> , Bq	Discharges during 2015	
			Bq	% of annual limit <sup>b</sup>
<b>Minor sites</b>				
Imperial College Reactor Centre Ascot	Tritium	4.00E+07	Nil	Nil
	Other radioactivity	1.00E+07	Nil	Nil
<b>Nuclear power stations</b>				
Berkeley	Tritium	1.00E+12	6.93E+07	<1
	Caesium-137	2.00E+11	2.17E+08	<1
	Other radionuclides	2.00E+11	7.20E+07	<1
Bradwell	Tritium	7.00E+12	3.23E+10	<1
	Caesium-137	7.00E+11	2.20E+08	<1
	Other radionuclides	7.00E+11	4.39E+09	<1
Chapelcross <sup>3</sup>	Alpha	1.00E+09	2.00E+06	<1
	Non-alpha <sup>1</sup>	1.00E+12	1.24E+09	<1
	Tritium	6.50E+12	8.66E+08	<1
Dungeness A Station	Tritium	8.00E+12	3.27E+10	<1
	Caesium-137	1.10E+12	9.10E+09	<1
	Other radionuclides	8.00E+11	2.63E+09	<1
Dungeness B Station	Tritium	6.50E+14	2.92E+14	45
	Sulphur-35	2.00E+12	3.12E+11	16
	Cobalt-60	1.00E+10	6.75E+08	6.8
	Caesium-137	1.00E+11	1.49E+09	1.5
	Other radionuclides	8.00E+10	3.06E+09	3.8
Hartlepool	Tritium	6.50E+14	1.43E+14	22
	Sulphur-35	3.00E+12	7.01E+11	23
	Cobalt-60	1.00E+10	1.43E+08	1.4
	Caesium-137	1.00E+11	1.27E+09	1.3
	Other radionuclides	8.00E+10	2.36E+08	<1
Heysham Station 1	Tritium	6.50E+14	1.04E+14	16
	Sulphur-35	2.00E+12	1.78E+11	8.9
	Cobalt-60	1.00E+10	1.09E+08	1.1
	Caesium-137	1.00E+11	1.37E+09	1.4
	Other radionuclides	8.00E+10	2.02E+09	2.5
Heysham Station 2	Tritium	6.50E+14	3.90E+14	60
	Sulphur-35	2.00E+12	4.14E+10	2.1
	Cobalt-60	1.00E+10	6.40E+07	<1
	Caesium-137	1.00E+11	2.56E+09	2.6
	Other radionuclides	8.00E+10	1.27E+10	16
Hinkley Point A Station	Tritium	1.00E+12	3.18E+10	3.2
	Caesium-137	1.00E+12	2.08E+10	2.1
	Other radionuclides	7.00E+11	1.88E+11	27
Hinkley Point B Station	Tritium	6.50E+14	2.43E+14	37
	Sulphur-35	2.00E+12	2.70E+11	14
	Cobalt-60	1.00E+10	1.15E+08	1.2
	Caesium-137	1.00E+11	7.78E+08	0.8
	Other radionuclides	8.00E+10	3.85E+09	4.8
Hunterston A Station <sup>3</sup>	Alpha	2.00E+09	1.96E+08	10
	All other non-alpha	6.00E+10	7.17E+08	1.2
	Tritium	3.00E+10	3.03E+08	1.0
	Caesium-137	1.60E+11	2.62E+08	<1
	Plutonium-241	2.00E+09	7.50E+07	3.8

**Table A2.2. continued**

Establishment	Radioactivity	Discharge limit (annual equivalent) <sup>a</sup> , Bq	Discharges during 2015	
			Bq	% of annual limit <sup>b</sup>
Hunterston B Station	Alpha	1.00E+09	3.10E+07	3.1
	All other non-alpha	1.50E+11	8.10E+09	5.4
	Tritium	7.00E+14	2.55E+14	36
	Sulphur-35	6.00E+12	7.00E+11	12
	Cobalt-60	1.00E+10	4.70E+08	4.7
Oldbury	Tritium	1.00E+12	3.07E+11	31
	Caesium-137	7.00E+11	1.49E+11	21
	Other radionuclides	7.00E+11	7.17E+10	10
Sizewell A Station	Tritium	5.00E+12	1.98E+10	<1
	Caesium-137	1.00E+12	4.21E+10	4.2
	Other radionuclides	7.00E+11	1.96E+10	2.8
Sizewell B Station	Tritium	8.00E+13	1.95E+13	24
	Caesium-137	2.00E+10	1.00E+09	5.0
	Other radionuclides	1.30E+11	3.00E+09	2.3
Torness	Alpha	5.00E+08	1.87E+06	<1
	All other non-alpha	1.50E+11	2.35E+09	1.6
	Tritium	7.00E+14	2.84E+14	41
	Sulphur-35	3.00E+12	8.75E+11	29
	Cobalt-60	1.00E+10	2.65E+08	2.7
Trawsfynydd	Tritium	3.00E+11	5.70E+09	1.9
	Caesium-137	1.50E+10	3.00E+08	2.0
	Other radionuclides <sup>j</sup>	3.00E+10	9.00E+08	3.0
Wylfa	Tritium	1.50E+13	1.33E+12	8.9
	Other radionuclides	1.10E+11	2.76E+09	2.5
<b>Defence establishments</b>				
Aldermaston (Silchester) <sup>k</sup>	Alpha	1.00E+07	1.57E+06	16
	Other beta emitting radionuclides	2.00E+07	2.08E+06	10
	Tritium	2.50E+10	1.00E+08	<1
Aldermaston (to Stream) <sup>k, l</sup>	Tritium	NA	5.40E+08	NA
Barrow <sup>m</sup>	Tritium	1.20E+10	6.83E+07	<1
	Carbon-14	2.70E+07	1.57E+05	<1
	Other gamma emitting radionuclides	3.50E+06	5.55E+03	<1
Derby <sup>n</sup>	Alpha <sup>o</sup>	2.00E+09	4.83E+07	2.4
	Alpha <sup>p</sup>	3.00E+05	7.00E+03	2.3
	Beta <sup>p</sup>	3.00E+08	2.22E+05	<1
Devonport (sewer) <sup>q</sup>	Tritium	2.00E+09	5.10E+07	2.6
	Cobalt-60	3.50E+08	3.13E+06	<1
	Other radionuclides	6.50E+08	1.17E+08	18
Devonport (estuary) <sup>q</sup>	Tritium	7.00E+11	4.83E+10	6.9
	Carbon-14	1.70E+09	7.54E+07	4.4
	Cobalt-60	8.00E+08	3.16E+07	4.0
	Other radionuclides	3.00E+08	2.27E+07	7.6
Faslane	Alpha	2.00E+08	1.20E+05	<1
	Beta <sup>r</sup>	5.00E+08	1.47E+06	<1
	Tritium	1.00E+12	1.95E+10	2.0
	Cobalt-60	5.00E+08	6.30E+05	<1
Rosyth	Tritium	3.00E+09	7.66E+07	2.6
	Cobalt-60	3.00E+08	3.31E+06	1.1
	Other radionuclides	3.00E+08	1.74E+06	<1

**Table A2.2. continued**

Establishment	Radioactivity	Discharge limit (annual equivalent) <sup>a</sup> , Bq	Discharges during 2015	
			Bq	% of annual limit <sup>b</sup>
<b>Radiochemical production</b>				
Amersham (GE Healthcare) <sup>f</sup>	Alpha	3.00E+08	2.90E+06	<1
	Tritium	1.41E+11	1.00E+06	<1
	Other radionuclides	6.50E+10	2.73E+08	<1
Cardiff (GE Healthcare) (January - May) <sup>4</sup>	Tritium	1.30E+14	Nil	Nil
	Carbon-14	9.10E+11	2.36E+08	<1
	Phosphorus-32/33	8.50E+07	Nil	Nil
	Iodine-125	3.00E+08	Nil	Nil
	Others	1.20E+08	Nil	Nil
<b>Industrial and landfill sites</b>				
LLWR	Alpha	BAT	4.08E+07	NA
	Beta	BAT	8.18E+08	NA
	Tritium	BAT	4.27E+10	NA
Lillyhall (Studsvik)	Alpha	5.00E+05	8.92E+02	<1
	Beta	5.00E+05	9.62E+03	1.9

<sup>a</sup> In some cases permits specify limits in greater detail than can be summarised in a single table; in particular, periods shorter than one year are specified at some sites

<sup>b</sup> Data quoted to 2 significant figures except when values are less than 1%

<sup>c</sup> Includes discharges made via the sea pipelines, factory sewer and Calder interceptor sewer

<sup>d</sup> The limit and discharge data are expressed in kg

<sup>e</sup> Some discharges are upper estimates because they include 'less than' data derived from analyses of effluents at limits of detection. Data quoted to 2 decimal places

<sup>f</sup> All alpha emitting radionuclides taken together

<sup>g</sup> All non-alpha emitting radionuclides, not specifically listed, taken together

<sup>h</sup> Discharges reported include those from INUTECH

<sup>i</sup> Excluding tritium

<sup>j</sup> Including strontium

<sup>k</sup> Discharges were made by AWE plc

<sup>l</sup> The discharge limit has been replaced by an activity notification level of 30 Bq l<sup>-1</sup>

<sup>m</sup> Discharges from Barrow are included with those from MOD sites because they are related to submarine activities. Discharges were made by BAE Systems Marine Ltd

<sup>n</sup> Discharges were made by Rolls Royce Marine Power Operations Ltd

<sup>o</sup> Discharge limit is for Nuclear Fuel Production Plant

<sup>p</sup> Discharge limit is for Neptune Reactor Raynesway site

<sup>q</sup> Discharges were made by Devonport Royal Dockyard Ltd

<sup>r</sup> Excluding cobalt-60

<sup>s</sup> Discharges were made by Rosyth Royal Dockyard Ltd

<sup>1</sup> Discharge permit revised with effect 1 March 2015, limits for Alpha, Beta and Tritium were revised. The discharges for alpha, beta and tritium are reported for the whole calendar year, however strictly only the discharges from March to December 2015 are compared to the new limits for compliance.

<sup>2</sup> Discharge authorisation revised with effect from 1 May 2014

<sup>3</sup> Discharge authorisation revised with effect from 1 July 2014

<sup>4</sup> Discharge authorisation active during this period (% of annual limit is calculated from the discharge and the discharge limit over the appropriate time period). Discharge permit revoked with effect from 1 June 2015

NA Not applicable under permit

BAT Best available technology

**Table A2.3. Disposals of solid radioactive waste at nuclear establishments in the United Kingdom, 2015**

Establishment	Radioactivity	Disposal limit	Disposals during 2015	
		Bq	Bq	% of limit <sup>a</sup>
LLWR <sup>b</sup>	Tritium	1.00E+13	Nil	Nil
	Carbon-14	5.00E+10	Nil	Nil
	Cobalt-60	2.00E+12	Nil	Nil
	Iodine-129	5.00E+10	Nil	Nil
	Radium-226 plus Thorium-232	3.00E+10	Nil	Nil
	Uranium	3.00E+11	Nil	Nil
	Other alpha <sup>c</sup>	3.00E+11	Nil	Nil
	Others <sup>c,d</sup>	1.50E+13	Nil	Nil
Dounreay <sup>e</sup>	Alpha		Nil	NA
	Beta/gamma		Nil	NA

<sup>a</sup> Data quoted to 2 significant figures except where values are less than 1%

<sup>b</sup> Under current planning permission at the LLWR near to Drigg, certain wastes are temporarily stored, as opposed to being disposed, pending disposal/storage elsewhere or permission for disposal in-situ

<sup>c</sup> With half-lives greater than 3 months excluding uranium, radium-226 and thorium-232

<sup>d</sup> Iron-55 and beta-emitting radionuclides with half-lives greater than three months unless individually specified in this table

<sup>e</sup> The current authorisation includes limits on concentrations of activity. At no time did the concentrations exceed the limits

NA Not applicable

**Table A2.4. Summary of unintended leakages, spillages, emissions or unusual findings of radioactive substances from nuclear licensed sites in the UK in 2015**

Site	Month	Summary of incident	Consequences and action taken
Capenhurst UUK	Q4 (Oct-Dec) 2015	A higher-than-usual radioactive gaseous discharge was recorded by UUK in Q4 of 2015 from the A3 Ventilated Room.	The discharge was below the quarterly notification level in the UUK EPR-RSA permit and the total annual discharge for 2015 was below the permitted annual limit. The discharge was investigated by UUK, but the cause of the higher-than-usual activity recorded in the sample could not be identified.  The site plan to review the specific operating instructions relating to the sampling of uranium in gaseous effluents in A3 plant.
Dounreay	July 2015	DSRL notified SEPA of the discovery of holes in the ventilation ducting associated with the on-site laboratories. It was subsequently established that the ventilation system was in an extremely poor condition.	As the holes within the ducting were downstream of any abatement, no additional radioactivity would have been discharged to the environment. SEPA sent a Warning Letter to DSRL in relation to the failure to keep the ventilation system in good repair and associated failures in the maintenance and inspection regime, and the failure to review the implications on existing infrastructure when delays to programmes occur.
Hinkley Point A	May and June 2015	Discharges of radioactive effluent were made without the operator following the correct process, due to errors in data handling. This could have resulted in radioactive effluent being discharged to the environment that might have required further treatment. However, a post-event review concluded that no further treatment of the effluent would have been necessary.	The discharges were compliant with the permit but only by chance. There was no environmental impact from the event itself. The Environment Agency raised non-compliances (CCS 3 and CCS 4) against the site permit. The site operator undertook a number of actions such as refresher training, revising work specifications (where necessary), review of laboratory conditions and carried out an effectiveness review of the actions. It is highlighted that Hinkley Point A ponds are now drained and therefore discharges of radioactivity will be substantially lower in future.
Hunterston B	November 2015	Hunterston B exceeded its authorised tidal window by approximately 26 minutes on one occasion in November 2015 by approximately 26 minutes, due to human error.	The discharge was stopped, whilst the tide was still ebbing, and there has been no discernible environmental impact as a result of the exceedance. SEPA subsequently issued the station with a Warning Letter in response to this contravention of the station's authorisation.

## APPENDIX 3. Abbreviations and glossary

ABL	AWE plc, Babcock and Lockheed Martin UK	GDA	Generic Design Assessment
ABWR	Advanced Boiling Water Reactor	GDF	Geological Disposal Facility
AGIR	Advisory Group on Ionising Radiation	GDL	Generalised Derived Limit
AGR	Advanced Gas-cooled Reactor	GE	General Electric
AWE	Atomic Weapons Establishment	GES	Good Environmental Status
BAT	Best Available Techniques or Best Available Technology	GOCO	Government Owned Contractor Operator
BEIS	Department of Business, Energy and Industrial Strategy	HMIP	Her Majesty's Inspectorate of Pollution
BIP	Border Inspection Post	HMNB	Her Majesty's Naval Base
BNFL	British Nuclear Fuels plc	HMSO	Her Majesty's Stationery Office
BPEO	Best Practicable Environmental Option	HPA	Health Protection Agency
BPM	Best Practicable Means	HSE	Health & Safety Executive
BSS	Basic Safety Standards	IAEA	International Atomic Energy Agency
CCFE	Culham Centre for Fusion Energy	ICRP	International Commission on Radiological Protection
CEC	Commission of the European Communities	ID	Indicative Dose
CEDA	Consultative Exercise on Dose Assessments	IRPA	International Radiation Protection Association
Cefas	Centre for Environment, Fisheries & Aquaculture Science	ISO	International Standards Organisation
CNLS	Cardiff Nuclear Licensed Site	JET	Joint European Torus
CNS	Capenhurst Nuclear Services Limited	LGC	Laboratory of the Government Chemist
COMARE	Committee on Medical Aspects of Radiation in the Environment	LLLETP	Low Level Liquid Effluent Treatment Plant
COS	Carbonyl Sulphide	LLW	Low Level Waste
CoRWM	Committee on Radioactive Waste Management	LLWR	Low Level Waste Repository
DECC	Department of Energy and Climate Change	LoD	Limit of Detection
DAERA	Department of Agriculture Environment and Rural Affairs	MAC	Medium Active Concentrate
Defra	Department for Environment, Food and Rural Affairs	MAFF	Ministry of Agriculture, Fisheries & Food
DPE	Designated Port of Entry	MCAA	Marine and Coastal Act 2009
DETR	Department of the Environment, Transport and the Regions	MMO	Marine Management Organisation
DH	Department of Health	MoD	Ministry of Defence
DPAG	Dounreay Particles Advisory Group	MRF	Metals Recycling Facility
DSRL	Dounreay Site Restoration Limited	MRL	Minimum Reporting Level
DSTL	Defence Science and Technology Laboratory	MRWS	Managing Radioactive Waste Safely
Euratom	European Atomic Energy Community	ND	Not Detected
EA	Environment Agency	NDA	Nuclear Decommissioning Authority
EARP	Enhanced Actinide Removal Plant	NFPP	Nuclear Fuel Production Plant
EC	European Commission	NIEA	Northern Ireland Environment Agency
EDF	Electricité de France	NII	Nuclear Installations Inspectorate
ENRMF	East Northants Resource Management Facility	NMP	Nuclear Management Partners Limited
EPR 10	Environment Permitting (England and Wales) Regulations 2010	NNB Genco	NNB Generation Company Limited
ERICA	Environmental Risk from Ionising Contaminants: Assessment and Management	NORM	Naturally Occurring Radioactive Material
ESC	Environmental Safety Case	NRPB	National Radiological Protection Board
ESG	Environmental Scientifics Group	NRW	Natural Resources Wales
EU	European Union	NPS	National Policy Statement
FEPA	Food and Environment Protection Act	NRTE	Naval Reactor Test Establishment
FSA	Food Standards Agency	OBT	Organically Bound Tritium
FSS	Food Standards Scotland	OECD	Organisation for Economic Co-operation and Development
		OMAD	Old Main Active Drain
		ONR	Office for Nuclear Regulation
		OSPAR	Oslo and Paris Convention
		PBO	Parent Body Organisation
		PRAG (D)	Particles Retrieval Advisory Group (Dounreay)
		PHE	Public Health England
		PWR	Pressurised Water Reactor

RAPs	Reference Animals and Plants	SWIMMER	Sustainable Water Integrated Management and Ecosystem Research
REP	RSR Environmental Principle	THORP	Thermal Oxide Reprocessing Plant
RIFE	Radioactivity in Food and the Environment	TNORM	Technologically enhanced Naturally Occurring Radioactive Material
RRDL	Rosyth Royal Dockyard Limited	TRAMP	Terrestrial Radioactive Monitoring Programme
RRMPOL	Rolls-Royce Marine Power Operations Limited	UCP	Urenco ChemPlants Limited
RNAS	Royal Naval Air Station	UKAEA	United Kingdom Atomic Energy Authority
RSA 93	Radioactive Substances Act 1993	UKNWM	UK Nuclear Waste Management Limited
RSR	Radioactive Substances Regulation	UOC	Uranium Ore Concentrate
RSRL	Research Sites Restoration Limited	UUK	Urenco UK Limited
RSS	Radioactive Substances Strategy	VLLW	Very Low Level Waste
SEPA	Scottish Environment Protection Agency	WFD	Water Framework Directive
SFL	Springfields Fuels Limited	WHO	World Health Organisation
SIXEP	Site Ion Exchange Plant	WWTW	Waste Water Treatment Works
SLC	Site Licence Company	YP	Ystradyfodwg and Pontypridd
SRP	Society for Radiological Protection		
STW	Sewage Treatment Works		

Absorbed dose	The ionising radiation energy absorbed in a material per unit mass. The unit for absorbed dose is the gray (Gy) which is equivalent to J kg <sup>-1</sup> .
Authorised Premises	This is a premises that has been authorised by the environment agencies to discharge to the environment.
Becquerel	One radioactive transformation per second.
Bioaccumulation	Excretion may occur, however the rate of excretion is less than the rate of intake + accumulation.
Biota	Flora and fauna.
Committed effective dose	The sum of the committed equivalent doses for all organs and tissues in the body resulting from an intake (of a radionuclide), having been weighted by their tissue weighting factors. The unit of committed effective dose is the sievert (Sv). The 'committed' refers to the fact that the dose is received over a number of years but it is accounted for in the year of the intake of the activity.
Direct shine	Ionising radiation which arises directly from processes or operations on premises using radioactive substances and not as a result of discharges of those substances to the environment.
Dose	Shortened form of 'effective dose' or 'absorbed dose'.
Dose limits	Maximum permissible dose resulting from ionising radiation from practices covered by the Euratom Basic Safety Standards Directive, excluding medical exposures. It applies to the sum of the relevant doses from external exposures in the specified period and the 50 year committed doses (up to age 70 for children) from intakes in the same period. Currently, the limit has been defined as 1 mSv per year for the UK.
Dose rates	The radiation dose delivered per unit of time.
Effective dose	The sum of the equivalent doses from internal and external radiation in all tissue and organs of the body, having been weighted by their tissue weighting factors. The unit of effective dose is the sievert (Sv).
Environmental materials	Environmental materials include freshwater, grass, seawater, seaweed, sediment, soil and various species of plants.

Equivalent dose	The absorbed dose in a tissue or organ weighted for the type and quality of the radiation by a radiation-weighting factor. The unit of equivalent dose is the sievert (Sv).
External dose	Doses to humans from sources that do not involve ingestion or inhalation of the radionuclides.
Fragments	'Fragments' are considered to be fragments of irradiated fuel, which are up to a few millimetres in diameter.
Generalised Derived Limit	A convenient reference level against which the results of environmental monitoring can be compared. GDLs are calculated using deliberately cautious assumptions and are based on the assumption that the level of environmental contamination is uniform over the year. GDLs relate the concentrations of a single radionuclide in a single environmental material to the dose limit for members of the public.
Indicator materials	Environmental materials may be sampled for the purpose of indicating trends in environmental performance or likely impacts on the food chain. These include seaweed, soil and grass.
In-growth	Additional activity produced as a result of radioactive decay of parent radionuclides.
Kerma air rate	Air kerma is the quotient of the sum of the kinetic energies of all the charged particles liberated by indirectly ionising particles in a specified mass of air.
Millisievert	The millisievert is a 1/1000 of a sievert. A sievert is one of the International System of Units used for the measurement of dose equivalent.
Radiation exposure	Being exposed to radiation from which a dose can be received.
Radiation weighting	Factor used to weight the tissue or organ absorbed dose to take account of the type and quality of the radiation. Example radiation weighting factors: alpha particles = 20; beta particles = 1; photons = 1.
Radioactivity	The emission of alpha particles, beta particles, neutrons and gamma or x-radiation from the transformation of an atomic nucleus.
Radionuclide	An unstable form of an element that undergoes radioactive decay.
Representative person	A hypothetical individual receiving a dose that is representative of the most exposed individuals in the population.
TNORM	Naturally occurring radioactive materials that may have been technologically enhanced in some way. The enhancement has occurred when a naturally occurring radioactive material has its composition, concentration, availability, or proximity to people altered by human activity. The term is usually applied when the naturally occurring radionuclide is present in sufficient quantities or concentrations to require control for purposes of radiological protection of the public or the environment.
Tissue weighting factors	Factor used to weight the equivalent dose in a tissue or organ to take account of the different radiosensitivity of each tissue and organ. Example tissue weighting factors: lung = 0.12; bone marrow = 0.12; skin = 0.01.
<i>Total dose</i>	An assessment of dose that takes into account all exposure pathways such as radionuclides in food and the environment and direct radiation.

## APPENDIX 4. Research in support of the monitoring programmes

FSA and the environment agencies have programmes of special investigations and supporting research and development studies to complement the routine monitoring programmes. This additional work is primarily directed at the following objectives:

- To evaluate the significance of potential sources of radionuclide contamination of the food chain and the environment
- To identify and investigate specific topics or pathways not currently addressed by the routine monitoring programmes and the need for their inclusion in future routine monitoring
- To develop and maintain site-specific habit and agricultural practice data, in order to improve the realism of dose assessment calculations
- To develop more sensitive and/or efficient analytical techniques for measurement of radionuclides in natural matrices
- To evaluate the competence of laboratories' radiochemical analytical techniques for specific radionuclides in food and environmental materials
- To develop improved methods for handling and processing monitoring data

Other studies include projects relating to effects on wildlife, emergency response and planning and development of new environmental models and data.

The contents of the research programmes are regularly reviewed and open meetings are held to discuss ongoing, completed and potential future projects. Occasionally specific topics are the subject of dedicated workshops (for example, Ould-Dada, 2000). A summary of all the research and development undertaken by the Environment Agency between 1996 and 2001 was published in 2002 (Environment Agency, 2002b). A review of research funded by FSA was published in 2004 (FSA, 2004).

Information on recently completed extramural research is presented in Table A4.1. Those sponsored by the Environment Agency and FSA are also listed on their websites ([www.environment-agency.gov.uk](http://www.environment-agency.gov.uk), [www.food.gov.uk](http://www.food.gov.uk), respectively). Copies of the final reports for each of the projects funded by the FSA are available from Aviation House, 125 Kingsway, London WC2B 6NH. Further information on studies funded by SEPA and the Scotland and Northern Ireland Forum for Environmental Research is available from Greenside House, 25 Greenside Place, Edinburgh, EH1 3AA. Environment Agency reports are available from [www.environment-agency.gov.uk](http://www.environment-agency.gov.uk). A charge may be made to cover costs.

**Table A4.1. Extramural Projects**

Topic	Reference	Further details	Target completion date
Soil and herbage survey	UKRSR01 and SCO00027	E, S	In press

E *Environment Agency*

S *Scotland and Northern Ireland Forum for Environmental Research or SEPA*

## APPENDIX 5. Disposal of dredge material from Maryport Harbour, Cumbria

In England, the Marine Management Organisation (MMO) administers a range of statutory controls that apply to marine works on behalf of the Secretary of State for Environment, Food and Rural Affairs (Defra), this includes issuing licences under the Marine and Coastal Access Act (MCAA), 2009 (United Kingdom - Parliament, 2009) for the disposal of dredged material at sea. Licences for disposals made in Scottish waters and around the coast of Northern Ireland are the responsibility of the Scottish Government (Marine Scotland) and the Department of Agriculture, Environment and Rural Affairs (NIEA), respectively and licences for Welsh waters are the responsibility of the Welsh Government.

The protection of the marine environment is considered before a licence is issued. Since dredge material will contain radioactivity from natural and man-made sources at varying concentrations, assessments are undertaken when appropriate for assurance that there is no significant food chain or other risk from the disposal. Guidance on exemption criteria for radioactivity in relation to sea disposal is available from the International Atomic Energy Agency (IAEA) (IAEA, 1999). IAEA has published guidance that can be applied to dredge spoil disposal to assess doses to workers and members of the public (IAEA, 2003). More recently, the assessment procedure has incorporated an additional approach for also assessing doses to marine flora and fauna (IAEA, 2015). The assessment to determine doses to workers and members of the public has been adapted to reflect operational practices in England and Wales (McCubbin and Vivian, 2006).


In 2015, Maryport Harbour Authority lodged a licensing application to carry out a dredging programme involving the disposal at sea of approximately 75,600 tonnes of sediment. A specific assessment of doses to workers and members of the public was conducted for the disposal of the dredge material (Smedley *et al.*, 2015).

The dock sediments contained artificial radionuclides typical of sediments along the Cumbrian coastline, being significantly enhanced above background levels outside the Irish Sea. The contamination is a legacy of large discharges from the Sellafield Limited reprocessing plant (formally British Nuclear Fuels) at Sellafield in the 1970's. Samples of the material were taken and analysed, and the results are given in Table A5.1. The contributions from individual radionuclides to the doses for individual crew members and individual members of the public are given in Figures A5.1-2. Under the London Convention, only materials with *de minimis* levels of radioactivity may be considered for dumping. Using the conservative generic radiological assessment procedure developed by the IAEA (International Atomic Energy Agency, 2003) to convert radionuclide concentrations in dumped material into radiation doses due to dumping, the total dose (from artificial and naturally-occurring radionuclides) to individual members of the crew and public were within the IAEA *de minimis* criteria of 0.010 mSv per year, for each of the three locations.


**Table A5.1. Concentrations of radionuclides in sediment dredged from Maryport Harbour, Cumbria, 2015**

Sample number	Mean radioactivity concentration (dry), Bq kg <sup>-1</sup>					
	<sup>60</sup> Co	<sup>137</sup> Cs	<sup>226</sup> Ra (via <sup>214</sup> Pb) <sup>1</sup>	<sup>232</sup> Th (via <sup>228</sup> Ac) <sup>1</sup>	<sup>238</sup> U (via <sup>234</sup> Th) <sup>1</sup>	<sup>241</sup> Am
Senhouse Approach (0m)	1.5	165	20	28	84	379
Outer Approach (0m)	1.0	128	21	25	66	344
Elizababeth Approach (0m)	1.0	129	17	21	59	302
Elizababeth Basin (0m)	1.7	197	21	28	72	450
Elizababeth Basin (1m)	2.4	294	24	38	65	670
<b>Mean</b>	<b>1.5</b>	<b>183</b>	<b>21</b>	<b>28</b>	<b>69</b>	<b>429</b>

<sup>1</sup> Parent nuclides not directly detected by the method used. Instead, concentrations were estimated from levels of their daughter products



**Figure A5.1.** Radionuclide contribution to dose to individual crew members (refined dose estimate) due to dredging at Maryport Harbour, Cumbria, 2015



**Figure A5.2.** Radionuclide contribution to dose to individual members of the public due to dredging at Maryport Harbour, Cumbria, 2015





Environment Agency  
Reactor Assessment and Radiological Monitoring, Nuclear Regulation Group (North)  
Cumbria and Lancashire Area, Lutra House, Preston, Lancashire PR5 8BX



Food Standards Agency  
Chemical Safety Division  
Aviation House, 125 Kingsway, London WC2B 6NH



Food Standards Scotland  
4th Floor, Pilgrim House,  
Old Ford Road, Aberdeen AB11 5RL



Cyfoeth Naturiol Cymru / Natural Resources Wales  
Ty Cambria, 29 Newport Road, Cardiff CF29 0TP



Northern Ireland Environment Agency  
Industrial Pollution and Radiochemical Inspectorate  
Klondyke Building, Cromac Avenue, Lower Ormeau Road, Belfast BT7 2JA



Scottish Environment Protection Agency  
Radioactive Substances Unit  
Strathallan House, Castle Business Park, Stirling FK9 4TZ